

On Axiomatic Characterization of Information-Theoretic Measure of Type $\binom{\alpha, \beta}{\gamma, \delta}$

I. J. TANEJA, H. C. GUPTA

There are two information-theoretic measures (viz., Kullback's relative-information and Kerridge's inaccuracy) associated with a pair of probability distributions of a discrete random variable. These measures have found many applications in Statistics, Economics, Physics etc. Two parameter generalization of these measures has been studied by many authors. A generalized measure of type $\binom{\alpha, \beta}{\gamma, \delta}$ is characterized in this paper by taking a set of axioms. This measure in particular contains relative-information and inaccuracy and their generalized forms as the limiting cases. Cases of bivariate extensions of this generalized measure and relations between them have been also studied.

1. INTRODUCTION

Associated with a pair of discrete probability distributions $P = (p_1, \dots, p_n)$, $p_i \geq 0$, $\sum_{i=1}^n p_i = 1$ and $Q = (q_1, \dots, q_n)$, $q_i > 0$, $\sum_{i=1}^n q_i \leq 1$ an information-theoretic measure of type $\binom{\alpha, \beta}{\gamma, \delta}$ is given by

$$(1.1) \quad I_{(\gamma, \delta)}^{(\alpha, \beta)}(P; Q) = (2^{\alpha-\beta} - 2^{\gamma-\delta})^{-1} \sum_{i=1}^n (p_i^\alpha q_i^{\beta-\alpha} - p_i^\gamma q_i^{\delta-\gamma}),$$

$\alpha \neq \gamma$ ($\beta \neq \delta$) whenever $\beta = \delta$ ($\alpha = \gamma$).

The measure (1.1) has been studied by Sharma and Taneja [13] and Sharma and Gupta [11] by generalizing a functional equation considered by Chaundy and McLeod [1] and by Taneja and Gupta [15] by considering a functional equation which is a generalization of the one considered by Daróczy [2].

For $\gamma = \delta = 1$ (refer Sharma and Autar [9, 10]) the measure (1.1) reduces to

$$(1.2) \quad I^{(\alpha, \beta)}(P; Q) = (2^{\alpha-\beta} - 1)^{-1} \left(\sum_{i=1}^n p_i^\alpha q_i^{\beta-\alpha} - 1 \right), \quad \beta \neq \alpha, \quad \alpha > 0.$$

The measure (1.2) reduces to (i) Kullback's [7] relative-information when $\beta = 1$ and there is a limiting case $\alpha \rightarrow 1$, (ii) Kerridge's [5] inaccuracy when $\alpha = 1$ and there is a limiting case $\beta \rightarrow 1$.

Further when $Q \equiv P$ and $\alpha = \gamma = \delta = 1$, (1.1) reduces to

$$(1.3) \quad I^\beta(P; Q) = H^\beta(P) = (2^{1-\beta} - 1)^{-1} \left(\sum_{i=1}^n p_i^\beta - 1 \right), \quad \beta \neq 1, \quad \beta > 0,$$

which is entropy of type β introduced by Havrda and Charvát [4] and later differently studied by Daróczy [2] and Vajda [16].

In this communication we characterize the measure (1.1) by taking a set of axioms (cf. Havrda and Charvát [4]). We also study bivariate extensions of measure (1.1) and establish relations between them.

2. CHARACTERIZATION THEOREM

Recursivity plays a vital role in characterization of information-theoretic measures. It is well known that the most elegant characterization of Shannon's entropy given by Faddeev (refer Feinstein [3]) so as those given earlier by Shannon [8] and Khinchine [6] use the recursive relation

$$(2.1) \quad H(p_1, \dots, p_n) - H(p_1 + p_2, p_3, \dots, p_n) = p_i H(p_1/p_i, p_2/p_i),$$

where $p_1 + p_2 = p_i > 0$.

Certain changes in this property of additive measures lead to non-additive measures. Havrda and Charvát [4] also used a modification of this property by introducing a parameter β as

$$(2.2) \quad H^\beta(p_1, \dots, p_n) - H^\beta(p_1 + p_2, p_3, \dots, p_n) = p_i^\beta H^\beta(p_1/p_i, p_2/p_i), \quad \beta > 0,$$

where $p_1 + p_2 = p_i > 0$.

Further, Taneja [14] generalized (2.2) by taking a general continuous function $f(p_i)$ in place of p_i^β and established that such a change does not lead to new measures and the only measures that arise are those studied by Shannon [8] and Havrda and Charvát [4].

Also an axiomatic characterization of (1.2) which is a measure of a pair of probability distributions has been studied by Sharma and Taneja [13] by considering the recursive relation of type (α, β) given by

$$(2.3) \quad I^{(\alpha, \beta)}(p_1, \dots, p_n; q_1, \dots, q_n) - I^{(\alpha, \beta)}(p_1 + p_2, p_3, \dots, p_n; q_1 + q_2, q_3, \dots, q_n) = p_i^\alpha q_i^{\beta-\alpha} I^{(\alpha, \beta)}(p_1/p_i, p_2/p_i; q_1/q_i, q_2/q_i),$$

where $p_1 + p_2 = p_i > 0$, $q_1 + q_2 = q_i > 0$ and α, β are the parameters.

Here we consider a much different form of the recursive relation to characterize (1.1) axiomatically. Precisely, for a pair of probability distributions $P = (p_1, \dots, p_n)$, $p_i \geq 0$, $\sum_{i=1}^n p_i = 1$ and $Q = (q_1, \dots, q_n)$, $q_i > 0$, $\sum_{i=1}^n q_i \leq 1$ we consider the following axioms:

- (a) $I_{(\gamma, \delta)}^{(\alpha, \beta)}(p_1, \dots, p_n; q_1, \dots, q_n)$ is a continuous function of its arguments;
- (b) $I_{(\gamma, \delta)}^{(\alpha, \beta)}(1, 0; \frac{1}{2}, \frac{1}{2}) = 1$; $I_{(\gamma, \delta)}^{(\alpha, \beta)}(1, 0; 1, 0) = 0$;
- (c) $I_{(\gamma, \delta)}^{(\alpha, \beta)}(p_1, \dots, p_{i-1}, 0, p_{i+1}, \dots, p_n; q_1, \dots, q_{i-1}, 0, q_{i+1}, \dots, q_n) = I_{(\gamma, \delta)}^{(\alpha, \beta)}(p_1, \dots, p_{i-1}, p_{i+1}, \dots, p_n; q_1, \dots, q_{i-1}, q_{i+1}, \dots, q_n)$, for every $i = 1, 2, \dots, n$;
- (d) $I_{(\gamma, \delta)}^{(\alpha, \beta)}(p_1, \dots, p_{i-1}, v_i, v_{i_2}, p_{i+1}, \dots, p_n; q_1, \dots, q_{i-1}, w_i, w_{i_2}, q_{i+1}, \dots, q_n) = I_{(\gamma, \delta)}^{(\alpha, \beta)}(p_1, \dots, p_{i-1}, p_i, p_{i+1}, \dots, p_n; q_1, \dots, q_{i-1}, q_i, q_{i+1}, \dots, q_n) + \frac{A_{\alpha, \beta}}{A_{\alpha, \beta} - A_{\gamma, \delta}} p_i^\alpha q_i^{\beta - \alpha} I_{(1, 1)}^{(\alpha, \beta)}\left(\frac{v_{i_1}}{p_i}, \frac{v_{i_2}}{p_i}; \frac{w_{i_1}}{q_i}, \frac{w_{i_2}}{q_i}\right) + \frac{A_{\gamma, \delta}}{A_{\gamma, \delta} - A_{\alpha, \beta}} p_i^\gamma q_i^{\delta - \gamma} I_{(\gamma, \delta)}^{(1, 1)}\left(\frac{v_{i_1}}{p_i}, \frac{v_{i_2}}{p_i}; \frac{w_{i_1}}{q_i}, \frac{w_{i_2}}{q_i}\right)$,

for every $v_{i_1} + v_{i_2} = p_i > 0$, $w_{i_1} + w_{i_2} = q_i > 0$, $i = 1, 2, \dots, n$ and $A_{\alpha, \beta} = (2^{\alpha - \beta} - 1)$ and $A_{\gamma, \delta} = (2^{\gamma - \delta} - 1)$.

Theorem 2.1. Axioms (a)–(d) determine a measure given by

$$(2.4) \quad I_{(\gamma, \delta)}^{(\alpha, \beta)}(p_1, \dots, p_n; q_1, \dots, q_n) = (A_{\alpha, \beta} - A_{\gamma, \delta})^{-1} \sum_{i=1}^n (p_i^\alpha q_i^{\beta - \alpha} - p_i^\gamma q_i^{\delta - \gamma}),$$

$$A_{\alpha, \beta} \neq A_{\gamma, \delta}, \quad \alpha, \gamma > 0,$$

where* $A_{\alpha, \beta} = (2^{\alpha - \beta} - 1)$ and $A_{\gamma, \delta} = (2^{\gamma - \delta} - 1)$.

The proof of the above theorem is based on the following lemmas:

Lemma 1. If $v_k \geq 0$, $k = 1, 2, \dots, m$, $\sum_{k=1}^m v_k = p_i > 0$, $w_k > 0$, $k = 1, 2, \dots, m$, $\sum_{k=1}^m w_k = q_i > 0$, then

$$(2.5) \quad I_{(\gamma, \delta)}^{(\alpha, \beta)}(p_1, \dots, p_{i-1}, v_1, \dots, v_m, p_{i+1}, \dots, p_n; q_1, \dots, q_{i-1}, w_1, \dots, w_m, q_{i+1}, \dots, q_n) = I_{(\gamma, \delta)}^{(\alpha, \beta)}(p_1, \dots, p_n; q_1, \dots, q_n) +$$

* Throughout this paper, we shall adopt the notation $A_{\alpha, \beta}$ for $(2^{\alpha - \beta} - 1)$ and $A_{\gamma, \delta}$ for $(2^{\gamma - \delta} - 1)$.

$$\begin{aligned}
& + \frac{A_{\alpha,\beta}}{A_{\alpha,\beta} - A_{\gamma,\delta}} p_i^\alpha q_i^{\beta-\alpha} I_{(1,1)}^{(\alpha,\beta)} \left(\frac{v_1}{p_i}, \dots, \frac{v_m}{p_i}, \frac{w_1}{q_i}, \dots, \frac{w_m}{q_i} \right) + \\
& + \frac{A_{\gamma,\delta}}{A_{\gamma,\delta} - A_{\alpha,\beta}} p_i^\gamma q_i^{\delta-\gamma} I_{(\gamma,\delta)}^{(1,1)} \left(\frac{v_1}{p_i}, \dots, \frac{v_m}{p_i}, \frac{w_1}{q_i}, \dots, \frac{w_m}{q_i} \right).
\end{aligned}$$

Proof. We prove the lemma by induction. For $m = 2$ the result holds (cf. axiom (d)). Let us suppose that the result is true for $m = t$. We have (under the notation $\bar{V} = v_2 + \dots + v_{t+1}$, $\bar{W} = w_2 + \dots + w_{t+1}$, $v_1 + \bar{V} = p_i > 0$, $w_1 + \bar{W} = q_i > 0$)

$$\begin{aligned}
(2.6) \quad & I_{(\gamma,\delta)}^{(\alpha,\beta)}(p_1, \dots, p_{i-1}, v_1, \dots, v_{t+1}, p_{i+1}, \dots, p_n; q_1, \dots, q_{i-1}, \\
& \quad w_1, \dots, w_{t+1}, q_{i+1}, \dots, q_n) = \\
& = I_{(\gamma,\delta)}^{(\alpha,\beta)}(p_1, \dots, p_{i-1}, v_1, \bar{V}, p_{i+1}, \dots, p_n; q_1, \dots, q_{i-1}, w_1, \bar{W}, q_{i+1}, \dots, q_n) + \\
& + \frac{A_{\alpha,\beta}}{A_{\alpha,\beta} - A_{\gamma,\delta}} \bar{V}^\alpha \bar{W}^{\beta-\alpha} I_{(1,1)}^{(\alpha,\beta)} \left(\frac{v_2}{\bar{V}}, \dots, \frac{v_{t+1}}{\bar{V}}, \frac{w_2}{\bar{W}}, \dots, \frac{w_{t+1}}{\bar{W}} \right) + \\
& + \frac{A_{\gamma,\delta}}{A_{\gamma,\delta} - A_{\alpha,\beta}} \bar{V}^\gamma \bar{W}^{\delta-\gamma} I_{(\gamma,\delta)}^{(1,1)} \left(\frac{v_2}{\bar{V}}, \dots, \frac{v_{t+1}}{\bar{V}}, \frac{w_2}{\bar{W}}, \dots, \frac{w_{t+1}}{\bar{W}} \right) = \\
& = I_{(\gamma,\delta)}^{(\alpha,\beta)}(p_1, \dots, p_n; q_1, \dots, q_n) + \\
& + \frac{A_{\alpha,\beta}}{A_{\alpha,\beta} - A_{\gamma,\delta}} p_i^\alpha q_i^{\beta-\alpha} I_{(1,1)}^{(\alpha,\beta)} \left(\frac{v_1}{p_i}, \frac{\bar{V}}{p_i}; \frac{w_1}{q_i}, \frac{\bar{W}}{q_i} \right) + \\
& + \frac{A_{\gamma,\delta}}{A_{\gamma,\delta} - A_{\alpha,\beta}} p_i^\gamma q_i^{\delta-\gamma} I_{(\gamma,\delta)}^{(1,1)} \left(\frac{v_1}{p_i}, \frac{\bar{V}}{p_i}; \frac{w_1}{q_i}, \frac{\bar{W}}{q_i} \right) + \\
& + \frac{A_{\alpha,\beta}}{A_{\alpha,\beta} - A_{\gamma,\delta}} \bar{V}^\alpha \bar{W}^{\beta-\alpha} I_{(1,1)}^{(\alpha,\beta)} \left(\frac{v_2}{\bar{V}}, \dots, \frac{v_{t+1}}{\bar{V}}; \frac{w_2}{\bar{W}}, \dots, \frac{w_{t+1}}{\bar{W}} \right) + \\
& + \frac{A_{\gamma,\delta}}{A_{\gamma,\delta} - A_{\alpha,\beta}} \bar{V}^\gamma \bar{W}^{\delta-\gamma} I_{(\gamma,\delta)}^{(1,1)} \left(\frac{v_2}{\bar{V}}, \dots, \frac{v_{t+1}}{\bar{V}}; \frac{w_2}{\bar{W}}, \dots, \frac{w_{t+1}}{\bar{W}} \right) = \\
& = I_{(\gamma,\delta)}^{(\alpha,\beta)}(p_1, \dots, p_n; q_1, \dots, q_n) + \\
& + \frac{A_{\alpha,\beta}}{A_{\alpha,\beta} - A_{\gamma,\delta}} p_i^\alpha q_i^{\beta-\alpha} I_{(1,1)}^{(\alpha,\beta)} \left(\frac{v_1}{p_i}, \frac{\bar{V}}{p_i}; \frac{w_1}{q_i}, \frac{\bar{W}}{q_i} \right) + \\
& + \frac{A_{\alpha,\beta}}{A_{\alpha,\beta} - A_{\gamma,\delta}} \bar{V}^\alpha \bar{W}^{\beta-\alpha} I_{(1,1)}^{(\alpha,\beta)} \left(\frac{v_2}{\bar{V}}, \dots, \frac{v_{t+1}}{\bar{V}}; \frac{w_2}{\bar{W}}, \dots, \frac{w_{t+1}}{\bar{W}} \right) + \\
& + \frac{A_{\gamma,\delta}}{A_{\gamma,\delta} - A_{\alpha,\beta}} p_i^\gamma q_i^{\delta-\gamma} I_{(\gamma,\delta)}^{(1,1)} \left(\frac{v_1}{p_i}, \frac{\bar{V}}{p_i}; \frac{w_1}{q_i}, \frac{\bar{W}}{q_i} \right) + \\
& + \frac{A_{\gamma,\delta}}{A_{\gamma,\delta} - A_{\alpha,\beta}} \bar{V}^\gamma \bar{W}^{\delta-\gamma} I_{(\gamma,\delta)}^{(1,1)} \left(\frac{v_2}{\bar{V}}, \dots, \frac{v_{t+1}}{\bar{V}}; \frac{w_2}{\bar{W}}, \dots, \frac{w_{t+1}}{\bar{W}} \right).
\end{aligned}$$

$$\begin{aligned}
 (2.7) \quad I_{(\gamma, \delta)}^{(\alpha, \beta)} \left(\frac{v_1}{p_i}, \dots, \frac{v_{t+1}}{p_i}; \frac{w_1}{q_i}, \dots, \frac{w_{t+1}}{q_i} \right) &= \\
 &= I_{(\gamma, \delta)}^{(\alpha, \beta)} \left(\frac{v_1}{p_i}, \frac{\bar{V}}{p_i}; \frac{w_1}{q_i}, \frac{\bar{W}}{q_i} \right) + \\
 &+ \frac{A_{\alpha, \beta}}{A_{\alpha, \beta} - A_{\gamma, \delta}} \left(\frac{\bar{V}}{p_i} \right)^\alpha \left(\frac{\bar{W}}{q_i} \right)^{\beta - \alpha} I_{(1, 1)}^{(\alpha, \beta)} \left(\frac{v_2}{\bar{V}}, \dots, \frac{v_{t+1}}{\bar{V}}; \frac{w_2}{\bar{W}}, \dots, \frac{w_{t+1}}{\bar{W}} \right) + \\
 &+ \frac{A_{\gamma, \delta}}{A_{\gamma, \delta} - A_{\alpha, \beta}} \left(\frac{\bar{V}}{p_i} \right)^\gamma \left(\frac{\bar{W}}{q_i} \right)^{\delta - \gamma} I_{(\gamma, \delta)}^{(1, 1)} \left(\frac{v_2}{\bar{V}}, \dots, \frac{v_{t+1}}{\bar{V}}; \frac{w_2}{\bar{W}}, \dots, \frac{w_{t+1}}{\bar{W}} \right).
 \end{aligned}$$

For $\gamma = \delta = 1$ (2.7) gives

$$\begin{aligned}
 (2.8) \quad I_{(1, 1)}^{(\alpha, \beta)} \left(\frac{v_1}{p_i}, \dots, \frac{v_{t+1}}{p_i}; \frac{w_1}{q_i}, \dots, \frac{w_{t+1}}{q_i} \right) &= I_{(1, 1)}^{(\alpha, \beta)} \left(\frac{v_1}{p_i}, \frac{\bar{V}}{p_i}; \frac{w_1}{q_i}, \frac{\bar{W}}{q_i} \right) + \\
 &+ \left(\frac{\bar{V}}{p_i} \right)^\alpha \left(\frac{\bar{W}}{q_i} \right)^{\beta - \alpha} I_{(1, 1)}^{(\alpha, \beta)} \left(\frac{v_2}{\bar{V}}, \dots, \frac{v_{t+1}}{\bar{V}}; \frac{w_2}{\bar{W}}, \dots, \frac{w_{t+1}}{\bar{W}} \right).
 \end{aligned}$$

For $\alpha = \beta = 1$ (2.7) gives

$$\begin{aligned}
 (2.9) \quad I_{(\gamma, \delta)}^{(1, 1)} \left(\frac{v_1}{p_i}, \dots, \frac{v_{t+1}}{p_i}; \frac{w_1}{q_i}, \dots, \frac{w_{t+1}}{q_i} \right) &= I_{(\gamma, \delta)}^{(1, 1)} \left(\frac{v_1}{p_i}, \frac{\bar{V}}{p_i}; \frac{w_1}{q_i}, \frac{\bar{W}}{q_i} \right) + \\
 &+ \left(\frac{\bar{V}}{p_i} \right)^\gamma \left(\frac{\bar{W}}{q_i} \right)^{\delta - \gamma} I_{(\gamma, \delta)}^{(1, 1)} \left(\frac{v_2}{\bar{V}}, \dots, \frac{v_{t+1}}{\bar{V}}; \frac{w_2}{\bar{W}}, \dots, \frac{w_{t+1}}{\bar{W}} \right).
 \end{aligned}$$

Expression (2.6) together with (2.8) and (2.9) gives the desired result.

Lemma 2. If $v_{ij} \geq 0, j = 1, 2, \dots, m_i; \sum_{j=1}^{m_i} v_{ij} = p_i > 0, i = 1, 2, \dots, n, \sum_{i=1}^n p_i = 1; w_{ij} > 0, j = 1, 2, \dots, m_i, \sum_{j=1}^{m_i} w_{ij} = q_i > 0, i = 1, 2, \dots, n, \sum_{i=1}^n q_i \leq 1$, then

$$\begin{aligned}
 (2.10) \quad I_{(\gamma, \delta)}^{(\alpha, \beta)}(v_{11}, \dots, v_{1m_1}, \dots, v_{n1}, \dots, v_{nm_n}; w_{11}, \dots, w_{1m_1}, \dots, w_{n1}, \dots, w_{nm_n}) &= \\
 &= I_{(\gamma, \delta)}^{(\alpha, \beta)}(p_1, \dots, p_n; q_1, \dots, q_n) + \\
 &+ \frac{A_{\alpha, \beta}}{A_{\alpha, \beta} - A_{\gamma, \delta}} \sum_{i=1}^n p_i^\alpha q_i^{\beta - \alpha} I_{(1, 1)}^{(\alpha, \beta)} \left(\frac{v_{i1}}{p_i}, \dots, \frac{v_{im_i}}{p_i}; \frac{w_{i1}}{q_i}, \dots, \frac{w_{im_i}}{q_i} \right) + \\
 &+ \frac{A_{\gamma, \delta}}{A_{\gamma, \delta} - A_{\alpha, \beta}} \sum_{i=1}^n p_i^\gamma q_i^{\delta - \gamma} I_{(\gamma, \delta)}^{(1, 1)} \left(\frac{v_{i1}}{p_i}, \dots, \frac{v_{im_i}}{p_i}; \frac{w_{i1}}{q_i}, \dots, \frac{w_{im_i}}{q_i} \right).
 \end{aligned}$$

The proof of this lemma directly follows from the Lemma 1.

209

Lemma 3. If $F_{(\gamma,\delta)}^{(\alpha,\beta)}(m; r) = f_{(\gamma,\delta)}^{(\alpha,\beta)}(1/m, \dots, 1/m; 1/r, \dots, 1/r)$ $1 \leq m \leq r$, then

$$(2.11) \quad F_{(\gamma,\delta)}^{(\alpha,\beta)}(m; r) = \frac{A_{\alpha,\beta}}{A_{\alpha,\beta} - A_{\gamma,\delta}} F_{(1,1)}^{(\alpha,\beta)}(m; r) + \frac{A_{\gamma,\delta}}{A_{\gamma,\delta} - A_{\alpha,\beta}} F_{(\gamma,\delta)}^{(1,1)}(m; r),$$

where

$$(2.12) \quad F_{(1,1)}^{(\alpha,\beta)}(m; r) = A_{\alpha,\beta}^{-1}(m^{1-\alpha}r^{\alpha-\beta} - 1), \quad A_{\alpha,\beta} \neq 0$$

and

$$(2.13) \quad F_{(\gamma,\delta)}^{(1,1)}(m; r) = A_{\gamma,\delta}^{-1}(m^{1-\gamma}r^{\gamma-\delta} - 1), \quad A_{\gamma,\delta} \neq 0.$$

Proof. Replacing in Lemma 2 m_i by m , $v_{ij} = 1/mn$, $w_{ij} = 1/rs$, $q_i = 1/s$, $i = 1, \dots, n$, $j = 1, \dots, m$ where m, n, r and s are positive integers such that $1 \leq m \leq r$, $1 \leq n \leq s$ we obtain

$$(2.14) \quad F_{(\gamma,\delta)}^{(\alpha,\beta)}(mn; rs) = F_{(\gamma,\delta)}^{(\alpha,\beta)}(n; s) + \frac{A_{\alpha,\beta}}{A_{\alpha,\beta} - A_{\gamma,\delta}} n^{1-\alpha} s^{\alpha-\beta} F_{(1,1)}^{(\alpha,\beta)}(m; r) + \\ + \frac{A_{\gamma,\delta}}{A_{\gamma,\delta} - A_{\alpha,\beta}} n^{1-\gamma} s^{\gamma-\delta} F_{(\gamma,\delta)}^{(1,1)}(m; r).$$

$$(2.15) \quad F_{(\gamma,\delta)}^{(\alpha,\beta)}(mn; rs) = F_{(\gamma,\delta)}^{(\alpha,\beta)}(m; r) + \frac{A_{\alpha,\beta}}{A_{\alpha,\beta} - A_{\gamma,\delta}} m^{1-\alpha} r^{\alpha-\beta} F_{(1,1)}^{(\alpha,\beta)}(n; s) + \\ + \frac{A_{\gamma,\delta}}{A_{\gamma,\delta} - A_{\alpha,\beta}} m^{1-\gamma} r^{\gamma-\delta} F_{(\gamma,\delta)}^{(1,1)}(n; s).$$

Putting $n = s = 1$ in (2.14) and using $F_{(\gamma,\delta)}^{(\alpha,\beta)}(1; 1) = 0$ (for all $\alpha, \beta, \gamma, \delta > 0$) we get

$$F_{(\gamma,\delta)}^{(\alpha,\beta)}(m; r) = \frac{A_{\alpha,\beta}}{A_{\alpha,\beta} - A_{\gamma,\delta}} F_{(1,1)}^{(\alpha,\beta)}(m; r) + \frac{A_{\gamma,\delta}}{A_{\gamma,\delta} - A_{\alpha,\beta}} F_{(\gamma,\delta)}^{(1,1)}(m; r)$$

which is (2.11)

Equating (2.14) and (2.15) we get

$$(2.16) \quad F_{(\gamma,\delta)}^{(\alpha,\beta)}(n; s) + \frac{A_{\alpha,\beta}}{A_{\alpha,\beta} - A_{\gamma,\delta}} n^{1-\alpha} s^{\alpha-\beta} F_{(1,1)}^{(\alpha,\beta)}(m; r) + \\ + \frac{A_{\gamma,\delta}}{A_{\gamma,\delta} - A_{\alpha,\beta}} n^{1-\gamma} s^{\gamma-\delta} F_{(\gamma,\delta)}^{(1,1)}(m; r) = \\ = F_{(\gamma,\delta)}^{(\alpha,\beta)}(m; r) + \frac{A_{\alpha,\beta}}{A_{\alpha,\beta} - A_{\gamma,\delta}} m^{1-\alpha} r^{\alpha-\beta} F_{(1,1)}^{(\alpha,\beta)}(n; s) + \\ + \frac{A_{\gamma,\delta}}{A_{\gamma,\delta} - A_{\alpha,\beta}} m^{1-\gamma} r^{\gamma-\delta} F_{(\gamma,\delta)}^{(1,1)}(n; s).$$

210 Expression (2.16) together with (2.11) gives

$$(2.17) \quad A_{\alpha,\beta}[(1 - m^{1-\alpha}r^{\alpha-\beta})F_{(1,1)}^{(\alpha,\beta)}(n; s) + (n^{1-\alpha}s^{\alpha-\beta} - 1)F_{(1,1)}^{(\alpha,\beta)}(m; r)] = \\ = A_{\gamma,\delta}[(1 - m^{1-\gamma}r^{\gamma-\delta})F_{(\gamma,\delta)}^{(1,1)}(n; s) + (n^{1-\gamma}s^{\gamma-\delta} - 1)F_{(\gamma,\delta)}^{(1,1)}(m; r)].$$

Putting $n = 1, s = 2$ in (2.17) and using

$$F_{(\gamma,\delta)}^{(\alpha,\beta)}(1; 2) = I_{(\gamma,\delta)}^{(\alpha,\beta)}(1, 0; \frac{1}{2}, \frac{1}{2}) = 1 \quad \text{for all } \alpha, \beta, \gamma, \delta > 0$$

we get

$$A_{\alpha,\beta}[(1 - m^{1-\alpha}r^{\alpha-\beta}) + (2^{\alpha-\beta} - 1)F_{(1,1)}^{(\alpha,\beta)}(m; r)] = \\ = A_{\gamma,\delta}[(1 - m^{1-\gamma}r^{\gamma-\delta}) + (2^{\gamma-\delta} - 1)F_{(\gamma,\delta)}^{(1,1)}(m; r)] = C \quad (\text{say}).$$

For $m = 1, r = 1$ we get $C = 0$. Thus we have

$$F_{(1,1)}^{(\alpha,\beta)}(m; r) = A_{\alpha,\beta}^{-1}(m^{1-\alpha}r^{\alpha-\beta} - 1), \quad A_{\alpha,\beta} \neq 0$$

and

$$F_{(\gamma,\delta)}^{(1,1)}(m; r) = A_{\gamma,\delta}^{-1}(m^{1-\gamma}r^{\gamma-\delta} - 1), \quad A_{\gamma,\delta} \neq 0$$

which are (2.12) and (2.13) respectively. This completes the proof of the lemma.

Now (2.11) together with (2.12) gives

$$(2.18) \quad F_{(\gamma,\delta)}^{(\alpha,\beta)}(m; r) = (A_{\alpha,\beta} - A_{\gamma,\delta})^{-1}(m^{1-\alpha}r^{\alpha-\beta} - m^{1-\gamma}r^{\gamma-\delta}).$$

Proof of the Theorem. We prove the theorem for rational values of p 's and q 's and then the continuity axiom (a) extends the result for all real values. Therefore let m, a_i and b_i be positive integers such that $a_i \leq b_i$ for every $i = 1, 2, \dots, n$ and if we put $p_i = a_i/m, q_i = b_i/r, i = 1, 2, \dots, n$ where $\sum_{i=1}^n a_i = m, \sum_{i=1}^n b_i \leq r$, then the application of the Lemma 2 gives

$$I_{(\gamma,\delta)}^{(\alpha,\beta)}\left(\underbrace{\frac{1}{m}, \dots, \frac{1}{m}}_{a_1}, \dots, \underbrace{\frac{1}{m}, \dots, \frac{1}{m}}_{a_n}; \underbrace{\frac{1}{r}, \dots, \frac{1}{r}}_{b_1}, \dots, \underbrace{\frac{1}{r}, \dots, \frac{1}{r}}_{b_n}\right) = \\ = I_{(\gamma,\delta)}^{(\alpha,\beta)}(p_1, \dots, p_n; q_1, \dots, q_n) + \\ + \frac{A_{\alpha,\beta}}{A_{\alpha,\beta} - A_{\gamma,\delta}} \sum_{i=1}^n p_i^\alpha q_i^{\beta-\alpha} I_{(1,1)}^{(\alpha,\beta)}\left(\frac{1}{a_i}, \dots, \frac{1}{a_i}; \frac{1}{b_i}, \dots, \frac{1}{b_i}\right) + \\ + \frac{A_{\gamma,\delta}}{A_{\gamma,\delta} - A_{\alpha,\beta}} \sum_{i=1}^n p_i^\gamma q_i^{\delta-\gamma} I_{(\gamma,\delta)}^{(1,1)}\left(\frac{1}{a_i}, \dots, \frac{1}{a_i}; \frac{1}{b_i}, \dots, \frac{1}{b_i}\right)$$

or

$$F_{(\gamma,\delta)}^{(\alpha,\beta)}(m; r) = I_{(\gamma,\delta)}^{(\alpha,\beta)}(p_1, \dots, p_n; q_1, \dots, q_n) +$$

$$+ \frac{A_{\alpha,\beta}}{A_{\alpha,\beta} - A_{\gamma,\delta}} \sum_{i=1}^n p_i^\alpha q_i^{\beta-\alpha} F_{(1,1)}^{(\alpha,\beta)}(a_i; b_i) + \frac{A_{\gamma,\delta}}{A_{\gamma,\delta} - A_{\alpha,\beta}} \sum_{i=1}^n p_i^\gamma q_i^{\delta-\gamma} F_{(\gamma,\delta)}^{(1,1)}(a_i; b_i)$$

i.e.

$$(2.19) \quad \begin{aligned} I_{(\gamma,\delta)}^{(\alpha,\beta)}(p_1, \dots, p_n; q_1, \dots, q_n) &= \\ &= F_{(\gamma,\delta)}^{(\alpha,\beta)}(m; r) - \frac{A_{\alpha,\beta}}{A_{\alpha,\beta} - A_{\gamma,\delta}} \sum_{i=1}^n p_i^\alpha q_i^{\beta-\alpha} F_{(1,1)}^{(\alpha,\beta)}(a_i; b_i) - \\ &\quad - \frac{A_{\gamma,\delta}}{A_{\gamma,\delta} - A_{\alpha,\beta}} \sum_{i=1}^n p_i^\gamma q_i^{\delta-\gamma} F_{(\gamma,\delta)}^{(1,1)}(a_i; b_i). \end{aligned}$$

Expression (2.19) together with (2.12), (2.13) and (2.18) gives (2.4).

3. BIVARIATE DISTRIBUTIONS

Let $X = (x_1, \dots, x_n)$ and $Y = (y_1, \dots, y_m)$ be two discrete random variables. The information-theoretic measure of type $\begin{pmatrix} \alpha, \beta \\ \gamma, \delta \end{pmatrix}$ for two probability distributions $P(X)$ and $Q(X)$ of X is given by

$$(3.1) \quad \begin{aligned} I_{(\gamma,\delta)}^{(\alpha,\beta)}(P(X); Q(X)) &= I_{(\gamma,\delta)}^{(\alpha,\beta)}(p(x_1), \dots, p(x_n); q(x_1), \dots, q(x_n)) = \\ &= (A_{\alpha,\beta} - A_{\gamma,\delta})^{-1} \sum_{i=1}^n (p^{\alpha}(x_i) q^{\beta-\alpha}(x_i) - p^{\gamma}(x_i) q^{\delta-\gamma}(x_i)), \end{aligned}$$

where $p(x_i) = P(X = x_i)$, $q(x_i) = Q(X = x_i)$, $i = 1, 2, \dots, n$ and $\sum_{i=1}^n p(x_i) = \sum_{i=1}^n q(x_i) = 1$.

An analogous formula holds for the random variable Y .

Next if $p(x_i, y_j) = P(X = x_i, Y = y_j)$ and $q(x_i, y_j) = Q(X = x_i, Y = y_j)$ are the joint probabilities of (x_i, y_j) then the joint information-theoretic measure of type $\begin{pmatrix} \alpha, \beta \\ \gamma, \delta \end{pmatrix}$ is given by

$$(3.2) \quad \begin{aligned} I_{(\gamma,\delta)}^{(\alpha,\beta)}(P(X, Y); Q(X, Y)) &= \\ &= I_{(\gamma,\delta)}^{(\alpha,\beta)}(p(x_1, y_1), \dots, p(x_1, y_m), \dots, p(x_n, y_1), \dots, p(x_n, y_m); \\ &\quad q(x_1, y_1), \dots, q(x_1, y_m), \dots, q(x_n, y_1), \dots, q(x_n, y_m)) = \\ &= (A_{\alpha,\beta} - A_{\gamma,\delta})^{-1} \sum_{i=1}^n \sum_{j=1}^m (p^{\alpha}(x_i, y_j) q^{\beta-\alpha}(x_i, y_j) - p^{\gamma}(x_i, y_j) q^{\delta-\gamma}(x_i, y_j)), \end{aligned}$$

where $\sum_{i=1}^n \sum_{j=1}^m p(x_i, y_j) = \sum_{i=1}^n \sum_{j=1}^m q(x_i, y_j) = 1$.

Further $P(Y/x_i) = \{p(y_1/x_i), \dots, p(y_m/x_i)\}$ and $Q(Y/x_i) = \{q(y_1/x_i), \dots, q(y_m/x_i)\}$ are complete distributions of Y given $X = x_i$, therefore the conditional information-theoretic measure of type $\begin{pmatrix} \alpha, \beta \\ \gamma, \delta \end{pmatrix}$ of Y given X may be defined as follows:

$$\begin{aligned}
 (3.3) \quad & I_{(\gamma, \delta)}^{(\alpha, \beta)}(P(Y/X); Q(Y/X)) = \\
 & = \frac{A_{\alpha, \beta}}{A_{\alpha, \beta} - A_{\gamma, \delta}} \sum_{i=1}^n p^\alpha(x_i) q^{\beta-\alpha}(x_i) I_{(1,1)}^{(\alpha, \beta)}(P(Y/x_i); Q(Y/x_i)) + \\
 & + \frac{A_{\gamma, \delta}}{A_{\gamma, \delta} - A_{\alpha, \beta}} \sum_{i=1}^n p^\gamma(x_i) q^{\delta-\gamma}(x_i) I_{(\gamma, \delta)}^{(1,1)}(P(Y/x_i); Q(Y/x_i)),
 \end{aligned}$$

where

$$(3.4) \quad I_{(1,1)}^{(\alpha, \beta)}(P(Y/x_i); Q(Y/x_i)) = A_{\alpha, \beta}^{-1} \left[\sum_{j=1}^m p^\alpha(y_j/x_i) q^{\beta-\alpha}(y_j/x_i) - 1 \right]$$

and

$$(3.5) \quad I_{(\gamma, \delta)}^{(1,1)}(P(Y/x_i); Q(Y/x_i)) = A_{\gamma, \delta}^{-1} \left[\sum_{j=1}^m p^\gamma(y_j/x_i) q^{\delta-\gamma}(y_j/x_i) - 1 \right],$$

$i = 1, 2, \dots, n$.

Similarly we can obtain the information-theoretic measure of type $\begin{pmatrix} \alpha, \beta \\ \gamma, \delta \end{pmatrix}$ of X given Y .

The interdependence and relationships among these bivariate measures are studied in the following theorems:

Theorem 3.1. If X and Y are two discrete random variables then

$$\begin{aligned}
 (3.6) \quad & I_{(\gamma, \delta)}^{(\alpha, \beta)}(P(X, Y); Q(X, Y)) = I_{(\gamma, \delta)}^{(\alpha, \beta)}(P(X); Q(X)) + I_{(\gamma, \delta)}^{(\alpha, \beta)}(P(Y/X); Q(Y/X)) = \\
 & = I_{(\gamma, \delta)}^{(\alpha, \beta)}(P(Y); Q(Y)) + I_{(\gamma, \delta)}^{(\alpha, \beta)}(P(X/Y); Q(X/Y)),
 \end{aligned}$$

where $p(x_i, y_j) = p(x_i) p(y_j/x_i) = p(y_j) p(x_i/y_j)$, $i = 1, 2, \dots, n$, $j = 1, 2, \dots, m$.

Proof. From the definitions given above we have

$$\begin{aligned}
 & I_{(\gamma, \delta)}^{(\alpha, \beta)}(P(X); Q(X)) + I_{(\gamma, \delta)}^{(\alpha, \beta)}(P(Y/X); Q(Y/X)) = \\
 & = (A_{\alpha, \beta} - A_{\gamma, \delta})^{-1} \sum_{i=1}^n (p^\alpha(x_i) q^{\beta-\alpha}(x_i) - p^\gamma(x_i) q^{\delta-\gamma}(x_i)) + \\
 & + (A_{\alpha, \beta} - A_{\gamma, \delta})^{-1} \sum_{i=1}^n p^\alpha(x_i) q^{\beta-\alpha}(x_i) \left[\sum_{j=1}^m p^\alpha(y_j/x_i) q^{\beta-\alpha}(y_j/x_i) - 1 \right] + \\
 & + (A_{\gamma, \delta} - A_{\alpha, \beta})^{-1} \sum_{i=1}^n p^\gamma(x_i) q^{\delta-\gamma}(x_i) \left[\sum_{j=1}^m p^\gamma(y_j/x_i) q^{\delta-\gamma}(y_j/x_i) - 1 \right] =
 \end{aligned}$$

$$\begin{aligned}
&= (A_{\alpha, \beta} - A_{\gamma, \delta})^{-1} \sum_{i=1}^n \sum_{j=1}^m [p^\alpha(x_i) p^\alpha(y_j/x_i) q^{\beta-\alpha}(x_i) q^{\beta-\alpha}(y_j/x_i) - \\
&\quad - p^\gamma(x_i) p^\gamma(y_j/x_i) q^{\delta-\gamma}(x_i) q^{\delta-\gamma}(y_j/x_i)] = \\
&= (A_{\alpha, \beta} - A_{\gamma, \delta})^{-1} \sum_{i=1}^n \sum_{j=1}^m [p^\alpha(x_i, y_j) q^{\beta-\alpha}(x_i, y_j) - p^\gamma(x_i, y_j) q^{\delta-\gamma}(x_i, y_j)] = \\
&\quad = I_{(\gamma, \delta)}^{(\alpha, \beta)}(P(X, Y); Q(X, Y)).
\end{aligned}$$

Theorem 3.2. If X and Y are statistically independent discrete random variables then

(a)

$$\begin{aligned}
(3.7) \quad I_{(\gamma, \delta)}^{(\alpha, \beta)}(P(X, Y); Q(X, Y)) &= G_{(\gamma, \delta)}^{(\alpha, \beta)}(P(X); Q(X)) I_{(\gamma, \delta)}^{(\alpha, \beta)}(P(Y); Q(Y)) + \\
&\quad + G_{(\gamma, \delta)}^{(\alpha, \beta)}(P(Y); Q(Y)) I_{(\gamma, \delta)}^{(\alpha, \beta)}(P(X); Q(X)),
\end{aligned}$$

where

$$(3.8) \quad G_{(\gamma, \delta)}^{(\alpha, \beta)}(P(X); Q(X)) = \frac{1}{2} \sum_{i=1}^n (p^\alpha(x_i) q^{\beta-\alpha}(x_i) + p^\gamma(x_i) q^{\delta-\gamma}(x_i)),$$

(b)

$$(3.9) \quad I_{(\gamma, \delta)}^{(\alpha, \beta)}(P(X, Y); Q(X, Y)) \leq I_{(\gamma, \delta)}^{(\alpha, \beta)}(P(X); Q(X)) + I_{(\gamma, \delta)}^{(\alpha, \beta)}(P(Y); Q(Y)),$$

for all $\alpha, \gamma \geq 1, \beta - \alpha \geq 1$ and $\delta - \gamma \geq 1$.

Proof. The proof of the part (a) follows by simple computation.

(b) For $\alpha, \gamma \geq 1, \beta - \alpha \geq 1$ and $\delta - \gamma \geq 1$ we have from (3.8)

$$(3.10) \quad G_{(\gamma, \delta)}^{(\alpha, \beta)}(P(X); Q(X)) \leq 1.$$

Expression (3.7) together with (3.10) gives (3.9).

ACKNOWLEDGEMENT

Authors are thankful to Dr. B. D. Sharma, Department of Mathematics, University of Delhi for guidance in the preparation of this paper and for discussions at various stages.

(Received May 5, 1976.)

- [1] T. W. Chaundy, J. B. McLeod: On a functional equation. Proc. Edin. Math. Soc. Edin. Math. Notes 43 (1960–61), 6–7.
- [2] Z. Daróczy: Generalized information functions. Information and Control 11 (1970), 36–51.
- [3] A. Feinstein: Foundations of Information Theory. McGraw-Hill, New York 1958.
- [4] J. Havrda, F. Charvát: Quantification method of classification processes, concept of structural α -entropy. Kybernetika 3 (1967), 30–35.
- [5] D. F. Kerridge: Inaccuracy and inference. J. Royal Stat. Soc. Ser. B 23 (1961), 1, 184–194.
- [6] A. I. Khinchine: Mathematical Foundations in Information Theory. Dover Publication, Inc., New York 1958.
- [7] S. Kullback: Information Theory and Statistics. Dover Pub. Inc., New York 1968.
- [8] C. E. Shannon: A mathematical theory of communication. Bell System Tech. J. 27 (1948), 379–423, 623–656.
- [9] B. D. Sharma, R. Autar: On characterization of a generalized inaccuracy measure in information theory. J. Appl. Prob. 10 (1973), 464–468.
- [10] B. D. Sharma, R. Autar: Relative-information functions and their type (α, β) generalizations. Metrika 21 (1974), 41–50.
- [11] B. D. Sharma, H. C. Gupta: Sub-additive measures of relative-information, and inaccuracy. Metrika 23 (1976), 155–165.
- [12] B. D. Sharma, I. J. Taneja: On axiomatic characterization of information-theoretic measures. J. Stat. Phys. 10 (1974), 337–346.
- [13] B. D. Sharma, I. J. Taneja: Entropy of type (α, β) and other generalized measures in information theory. Metrika 22 (1975), 205–215.
- [14] I. J. Taneja: On the branching property of entropy. Annales Polonici Mathematici 35 (1977), 67–75.
- [15] I. J. Taneja, H. C. Gupta: On generalized measures of relative-information and inaccuracy. Aplikace matematiky 23 (1978).
- [16] I. Vajda: Axioms for α -entropy of generalized probability scheme. Kybernetika 4 (1968), 2, 105–112.

Dr. I. J. Taneja, Departamento de Matematica, Universidade Federal de Santa Catarina, 88000 – Florianopolis, SC, Brazil.

Dr. H. C. Gupta, Department of Mathematics, Rajdhani College (University of Delhi), Raja Garden, Ring Road, New Delhi - 110015, India.