

## OPTIMAL DISCRETE APPROXIMATION OF CONTINUOUS LINEAR OPERATORS APPLICABLE TO CONTROL PROBLEMS

ANTONÍN TUZAR

This article is devoted to some problems of optimal discrete approximation of continuous linear operators. It presents continuation of our work [18] and brings a more general theorem of approximation (Theorem 2.1), some examples of the form of cardinal spline-functions (in Sec. 3) and an example of solution of a simple optimum control problem (Sec. 4), on which we try to compare the current and new optimal formulae for derivatives. Some necessary facts and notations are introduced in Section 1. Our results are based on a theory which was originally developed for purposes of numerical quadratures and cubatures and for the interpolation [11], [13], [14], [15]. Bibliographical notes see e.g. in [7], [10]. In the following we deal mostly with approximation of derivatives.

### 1. THEORETICAL BACKGROUND

Let  $H$  be a real Hilbert space with the scalar product  $\langle \cdot, \cdot \rangle$  and the norm  $\| \cdot \|$ . Let  $\langle h, \cdot \rangle$  and  $\langle \varphi_1, \cdot \rangle, \dots, \langle \varphi_n, \cdot \rangle$  be some bounded linear functionals on  $H$ . We shall approximate  $h$  by the expression  $\sum_{i=1}^n c_i \varphi_i$  with  $c_i$  real in such a way that the norm of error functional will be minimized:

$$(1.1) \quad \left\| h - \sum_{i=1}^n c_i^0 \varphi_i \right\| = \min_c \left\| h - \sum_{i=1}^n c_i \varphi_i \right\|, \quad \text{where } c \in \mathbb{R}^n.$$

The coordinates of  $c^0 = (c_1^0, \dots, c_n^0)$  are called the optimal coefficients. The elements  $\varphi_i$ ,  $i = 1, \dots, n$  are supposed to be linearly independent. Then they form a *natural* basis of a linear  $n$ -dimensional subspace  $S \subset H$ . It is easy to prove the existence of the unique set of linearly independent elements  $\sigma_j \in S$ ,  $j = 1, \dots, n$ , for which the following conditions hold

$$(1.2) \quad \langle \varphi_i, \sigma_j \rangle = \delta_{ij}; \quad i, j = 1, \dots, n.$$

Here  $\delta_{ij}$  is Kronecker's symbol. We call this set  $\{\sigma_1, \dots, \sigma_n\}$  the *cardinal basis* of  $S$ . Our following concept is based on

**Theorem 1.1.** The real numbers  $c_1^0, \dots, c_n^0$  are optimal coefficients in the approximation of bounded linear functional  $\langle h, \cdot \rangle$  on  $H$  by means of  $\langle \sum_{i=1}^n c_i \varphi_i, \cdot \rangle$ , iff

$$(1.3) \quad c_j^0 = \langle h, \sigma_j \rangle, \quad j = 1, \dots, n,$$

where  $\sigma_1, \dots, \sigma_n$  are given as the solution of system (1.2). For  $h \in S$  (1.1) is equal to zero.

*Proof.* The nearest element in  $S$  to the given  $h \in H$  is the orthogonal projection of  $h$  to  $S$ . Therefore  $h - \sum_{i=1}^n c_i^0 \varphi_i$  is orthogonal to all elements  $\sigma_j$  of the cardinal basis

$$\langle \sigma_j, h - \sum_{i=1}^n c_i^0 \varphi_i \rangle = 0, \quad j = 1, \dots, n,$$

which, using the property (1.2), immediately leads to (1.3).  $\square$

## 2. APPROXIMATION OF DERIVATIVES ON SOBOLEV SPACE

We shall investigate the optimal coefficients of formulae, which approach the linear bounded operators on Sobolev spaces  $H^q(a, b)$  with integer  $q \geq 2$ . The elements of  $H^q(a, b)$  are absolutely continuous functions on  $[a, b]$  with quadratic integrable generalized (in sense of theory of distributions) derivatives of order  $q$ . The scalar product and the corresponding norm are introduced in the following manner (see, e.g. [3] or [15], Chapter XIV, § 4). Let be given  $q$  arbitrary points

$$(2.1) \quad -\infty < a < x_1 < \dots < x_q < b < +\infty.$$

and put

$$(2.2) \quad \langle f, g \rangle_H = \int_a^b f^{(q)}(x) g^{(q)}(x) dx + \sum_{i=1}^q f(x_i) g(x_i),$$

$$(2.3) \quad \|f\|_H^2 = \langle f, f \rangle_H.$$

This norm is equivalent to the usual norm

$$(2.4) \quad \|f\|^2 = \sum_{i=0}^q \int_a^b [f^{(i)}(x)]^2 dx.$$

The main advantage of the norm given by (2.2)–(2.3) is connected with the fact that we obtain the elements of the linear subspace  $S$  in a very simple form. Namely we will show that the functionals of the form

$$\langle \varphi_{p,y}, f \rangle_H = f^{(p)}(y), \quad a < y < b, \quad 0 \leq p < q$$

are generated by spline functions, i.e.  $\varphi_{p,y}$  are piecewise polynomials. Introducing another norm, e.g. (2.4), this will not be true.

The following inequalities are well known as the Sobolev imbedding theorem:

$$(2.5) \quad \|f^{(p)}\|_{C(a,b)} \leq K \|f\|_H, \quad p = 0, 1, \dots, q-1,$$

where  $q > 1$  and  $K$  does not depend on  $f \in H^q(a, b)$ . Therefore the values of function and its derivatives of order less than  $q$  in given interior points of  $]a, b[$  are linear bounded functionals on  $H^q(a, b)$  and so there is reasonable to find the optimal approximations for them using Theorem 1.1. We will find the approximative formulae for evaluation of some derivative using the values of function and its derivatives of lower order in the points of a given net. Generally, let be given the points

$$(2.6) \quad \begin{aligned} a < x_{01} < \dots < x_{0n_0} < b \\ a < x_{11} < \dots < x_{1n_1} < b \\ \dots \dots \dots \dots \dots \dots \dots \\ a < x_{p1} < \dots < x_{pn_p} < b, \end{aligned}$$

where  $0 \leq p \leq q - 1$ ,  $q > 1$ . Moreover, suppose that  $n_0 \geq q$  and that the points (2.1) are some arbitrary, but fixed of  $x_{01}, \dots, x_{0n_0}$ . We introduce the bounded linear functionals on  $H^q(a, b)$ , for which

$$(2.7) \quad \begin{aligned} \langle \varphi_{kl}, f \rangle_H = f^{(k)}(x_{kl}), \quad \forall f \in H^q(a, b), \\ k = 0, \dots, p; \quad l = 1, \dots, n_k. \end{aligned}$$

Those functionals are linearly independent. Indeed, in  $H^q(a, b)$  there evidently exist elements  $\psi_{kl}$  such that  $\psi_{kl}^{(k)}(x_{kl}) = 1$  and  $\psi_{st}^{(s)}(x_{st}) = 0$  for all remaining ordered pairs  $(s, t) \neq (k, l)$ . Let us suppose

$$\sum_{s,t} \lambda_{st} \varphi_{st} = 0,$$

where not all  $n_0 + n_1 + \dots + n_p$  constants  $\lambda_{st}$  are equal to zero. Multiplying the left side scalarly in  $H^q(a, b)$  by  $\psi_{kl}$ , we obtain

$$\langle \psi_{kl}, \sum_{s,t} \lambda_{st} \varphi_{st} \rangle_H = \lambda_{kl},$$

therefore all  $\lambda_{kl} = 0$  and this contradiction proves the linear independency of  $\varphi_{kl}$ . We denote by  $S$  the linear subspace spanned on all  $\varphi_{kl}$  defined by (2.7). This subspace  $S$  is, as we have just proved,  $n$ -dimensional, where

$$(2.8) \quad n = n_0 + n_1 + \dots + n_p.$$

The set of all elements  $\varphi_{kl}$  is the natural basis of  $S$ . The cardinal basis in  $S$  consists of elements  $\sigma_{kl}$  with the following properties:

$$(2.9) \quad \langle \varphi_{kl}, \sigma_{st} \rangle = \delta_{ks} \delta_{lt},$$

where

$$\begin{aligned} k, s &= 0, 1, \dots, p, \\ l &= 1, \dots, n_k, \\ t &= 1, \dots, n_s. \end{aligned}$$

The system of  $n$  equations (2.9), where  $n$  is given by (2.8), has exactly one solution

of the form

$$(2.10) \quad \sigma_{st} = \sum_{u,v} \lambda_{stuv} \varphi_{uv}$$

where  $\lambda_{stuv}$  are real constants. This fact follows from linear independency of the elements  $\varphi_{uv}$ , as the matrix of linear system (2.9) after substitution from (2.10) is the Gramm matrix  $[\langle \varphi_{ki}, \varphi_{st} \rangle]_{k,i,s,t}$  which is necessarily regular. The elements  $\varphi_{ki}$  with properties (2.9) and generally the elements of the linear subspace  $S$  can be found using the following

**Theorem 2.1.** Let the points (2.1) be given, let  $y \in ]a, b[$  and let  $p$  be an integer such that  $0 \leq p < q, q > 1$ . Then the unique element  $\psi \in H^q(a, b)$ , such that

$$(2.11) \quad \langle \psi, f \rangle_H = f^{(p)}(y), \quad \forall f \in H^q,$$

is the solution of the following boundary value problem:

$$(2.12) \quad \psi^{(2q)}(x) = 0, \quad x \in ]a, b[ - \bigcup_{i=1}^q \{x_i\} - \{y\},$$

$$(2.13) \quad (-1)^{q-p} [\psi^{(2q-p-1)}(y_+) - \psi^{(2q-p-1)}(y_-)] = \begin{cases} 1 - \psi(y), & \text{if } p = 0 \\ \text{and } y = x_i & \text{for some} \\ i = 1, \dots, q, \\ 1 & \text{otherwise} \end{cases}$$

$$(2.14) \quad (-1)^{q+1} [\psi^{(2q-1)}(x_{i+}) - \psi^{(2q-1)}(x_{i-})] = \psi(x_i), \quad i = 1, \dots, q, \quad x_i \neq y,$$

$$(2.15) \quad \psi^{(q)}(a) = \psi^{(q)}(b) = \psi^{(q+1)}(a) = \psi^{(q+1)}(b) = \dots = \psi^{(2q-1)}(a) = \psi^{(2q-1)}(b) = 0,$$

$$(2.16) \quad \psi(\cdot), \dots, \psi^{(2q-p-2)}(\cdot) \text{ are continuous on } [a, b].$$

Denoting by  $(z)_+ = \max(0, z), z \in \mathbb{R}^1$  we can express the function  $\psi(\cdot)$  in the form

$$(2.17) \quad \psi(x) = \eta_{q-1}(x) + \frac{(-1)^{q-p}}{(2q-p-1)!} (x-y)_+^{2q-p-1} + \sum_{i=1}^q \frac{(-1)^{q-1} \psi(x_i)}{(2q-1)!} (x-x_i)_+^{2q-1},$$

where  $\eta_{q-1}(\cdot)$  is a suitable polynomial of order less than  $q$ .

**Proof.** According to the Sobolev imbedding theorem (2.5) for an arbitrary  $y \in ]a, b[$  and  $0 \leq p < q, q \geq 2$  the linear functional (2.11) is bounded. The existence and uniqueness of  $\psi$  in (2.11) follows from the theorem by F. Riesz.

On the other hand it is easy to verify that the solution of boundary value problem (2.12)  $\div$  (2.16) can be found in the form (2.17). Notice that the unknown  $q$  coefficients of the polynomial  $\eta_{q-1}$  and  $q$  values  $\psi(x_i), i = 1, \dots, q$  are determined from the equations  $\psi^{(q)}(b) = \psi^{(q+1)}(b) = \dots = \psi^{(2q-1)}(b) = 0$  ( $q$  independent equations) and from the substitution of  $x_i, i = 1, \dots, q$  into (2.17) (also  $q$  independent equations). Let  $\psi(\cdot)$  be the function (2.17). Using integration by parts and denoting  $a = x_0$ ,

$b = x_{q+1}$ , we express the scalar product as follows:

$$(2.18) \quad \langle \psi, f \rangle_H = \int_{x_0}^{x_{q+1}} \psi^{(q)}(x) f^{(q)}(x) dx + \sum_{i=1}^q \psi(x_i) f(x_i) = \\ \psi^{(q)}(x) f^{(q-1)}(x) \Big|_{x_0}^{x_{q+1}} - \psi^{(q+1)}(x) f^{(q-2)}(x) \Big|_{x_0}^{x_{q+1}} + \dots \\ \dots + (-1)^{q-p-1} \psi^{(2q-p-1)}(x) f^{(p)}(x) \Big|_{x_0}^{x_q} + (-1)^{q-p-1} \psi^{(2q-p-1)}(x) \cdot \\ \cdot f^{(p)}(x) \Big|_{x_q}^{x_{q+1}} + \dots + \sum_{i=0}^q (-1)^{q-1} \psi^{(2q-1)}(x) f(x) \Big|_{x_i}^{x_{i+1}} + \\ + (-1)^q \int_{x_0}^{x_{q+1}} \psi^{(2q)}(x) f(x) dx + \sum_{i=1}^q \psi(x_i) f(x_i).$$

Using properties (2.12) + (2.16) the equality (2.11) can be verified.  $\square$

**Remark.** From the previous theorem it follows that the elements of linear subspace  $S$  are spline-functions of order equal or less than  $2q - 1$ . Moreover, in the intervals  $]a, \min(x_1, y)[$  and  $]\max(x_q, y), b[$  they are equal to polynomials of order  $r \leq q - 1$ . For  $x \in ]a, \min(x_1, y)[$  this assertion follows directly from (2.17) and for  $x \in ]\max(x_q, y), b[$  from the conditions (2.15) in  $b$ .

**Example.** Let be  $a = 0, b = 1, p = 1, q = 2$  and  $x_1 = \frac{1}{4}, x_2 = \frac{3}{4}, y = \frac{1}{2}$ . In this case using expression (2.17) there is

$$(2.19) \quad \psi(x) = a_1 x + a_0 - \frac{1}{2}(x - \frac{1}{2})_+^2 - \frac{1}{6}[\psi(\frac{1}{4})(x - \frac{1}{4})_+^3 + \psi(\frac{3}{4})(x - \frac{3}{4})_+^3].$$

Denoting by  $\mathbf{1}(\cdot)$  the unit jump and by  $\delta(\cdot)$  the Dirac delta distribution, we express

$$\psi''(x) = -\mathbf{1}(x - \frac{1}{2}) - \psi(\frac{1}{4})(x - \frac{1}{4})_+ - \psi(\frac{3}{4})(x - \frac{3}{4})_+, \\ \psi'''(x) = -\delta(x - \frac{1}{2}) - \psi(\frac{1}{4})\mathbf{1}(x - \frac{1}{4}) - \psi(\frac{3}{4})\mathbf{1}(x - \frac{3}{4}),$$

and applying conditions (2.15) at  $b = 1$ , we obtain the solution of equations  $\psi''(1) = \psi'''(1) = 0$  as  $\psi(\frac{1}{4}) = -2, \psi(\frac{3}{4}) = 2$ . Now it is possible to determine the coefficients  $a_0, a_1$  using the substitution  $x = \frac{1}{4}$  and  $x = \frac{3}{4}$  into (2.19):

$$-2 = \frac{1}{4}a_1 + a_0, \\ 2 = \frac{3}{4}a_1 + a_0 + \frac{1}{2}(\frac{3}{4} - \frac{1}{2})^2 - \frac{1}{6}\{2(\frac{3}{4} - \frac{1}{4})^3\}.$$

Solving this system, we obtain

$$(2.20) \quad \psi(x) = \frac{383}{48}x - \frac{767}{192} - \frac{1}{2}(x - \frac{1}{2})_+^2 + \frac{1}{3}[(x - \frac{1}{4})_+^3 - (x - \frac{3}{4})_+^3].$$

There is not difficult to verify that for  $x \in ]0, \frac{1}{4}[$  and for  $x \in ]\frac{3}{4}, 1[$  the function (2.20) its equal to a polynomial of order  $q - 1 = 1$ .

Using Theorem 2.1 it is possible in every concrete case to evaluate the corresponding elements of the natural basis. As their linear combinations have obligatory equal

structure we can modify this method for computation of elements of cardinal basis. According to the previous remark we shall call this elements cardinal splines.

Let be  $0 \leq p < s < q$  and let  $y \in ]a, b[$ . From Theorems 1.1 and 2.1 we can compute the optimal approximation of derivative  $f^{(s)}(y)$  for arbitrary  $y \in ]a, b[$ , namely

$$(2.21) \quad f^{(s)}(y) \doteq \sum_{i=0}^p \sum_{j=1}^{n_i} \sigma_{ij}^{(s)}(y) f^{(i)}(x_{ij}).$$

Analogously the optimal quadrature formula for  $f \in H^q(a, b)$  is of the form

$$(2.22) \quad \int_a^b f(x) dx \doteq \sum_{i=0}^p \sum_{j=1}^{n_i} \left( \int_a^b \sigma_{ij}(x) dx \right) f^{(i)}(x_{ij}).$$

### 3. CARDINAL SPLINES FOR EQUIDISTANT PARTITION

Since now we shall consider the simplest case  $p = 0$  and

$$(3.1) \quad \langle \varphi_i, f \rangle_H = f(x_i), \quad i = 1, \dots, n, \quad \forall f \in H^q(a, b),$$

where  $n \geq q > 1$  and

$$(3.2) \quad x_i = a + \frac{i}{n+1}(b-a), \quad i = 0, 1, \dots, n+1.$$

Especially,  $x_0 = a$ ,  $x_{n+1} = b$ .  $S$  is the linear subspace of  $H^q(a, b)$  spanned on  $\{\varphi_1, \dots, \varphi_n\}$ .

From Theorem 2.1 it follows

**Proposition 3.1.** The cardinal splines  $\sigma_i \in S$ , where  $S$  is the linear subspace in  $H^q(a, b)$  consisting of elements  $\varphi_i$ ,  $i = 1, \dots, n$ , with properties (3.1), are the functions of the form

$$(3.3) \quad \sigma_i(x) = \begin{cases} \sum_{j=0}^{q-1} a_{ij} x^j + \sum_{j=1}^{n-1} \lambda_{ij} (x - x_j)_+^{2q-1}, & x \in [x_0, x_n[ \\ \sum_{j=0}^{q-1} b_{ij} x^j, & x \in [x_n, x_{n+1}] \end{cases}$$

with the properties

$$(3.4) \quad \sigma_i(x_j) = \delta_{ij}, \quad i, j = 1, \dots, n.$$

There is  $\sigma_i \in C^{(2q-2)}(a, b)$  and the  $2q + n - 1$  coefficients  $a_{10}, \dots, a_{iq}, \lambda_{i1}, \dots, \lambda_{i,n-1}, b_{i0}, \dots, b_{iq}$  are the (unique) solution of the linear system of independent  $n$  equations (3.4) and  $2q - 1$  equations

$$(3.5) \quad \sigma_i^{(k)}(x_n -) = \sigma_i^{(k)}(x_n +), \quad k = 0, 1, \dots, 2q - 2.$$

**Remark.** The assertion about the independency of equations (3.4), (3.5) can be verified ([2]) also directly, if we analyse the matrix of the corresponding linear

system. We present here some numerical results, based on Proposition 3.1. In our examples we suppose the interval  $[a, b]$  transformed to the special form  $[0, (n + 1)h]$ . In the tables there are values of coefficients in the expression (3.3).

**Example 3.1.**

$n = 3, q = 2$

$i$	$a_{i0}$	$a_{i1}$	$\lambda_{i1}$	$\lambda_{i2}$	$b_{i0}$	$b_{i1}$
1	$\frac{9}{4}$	$-\frac{5}{4h}$	$\frac{1}{4h^3}$	$-\frac{2}{4h^3}$	$-\frac{3}{4}$	$\frac{1}{4h}$
2	$-\frac{3}{2}$	$\frac{3}{2h}$	$-\frac{1}{2h^3}$	$\frac{2}{2h^3}$	$\frac{9}{2}$	$-\frac{3}{2h}$
3	$\frac{1}{4}$	$-\frac{1}{4h}$	$\frac{1}{4h^3}$	$-\frac{2}{4h^3}$	$-\frac{11}{4}$	$\frac{5}{4h}$

**Example 3.2.**

$n = 4, q = 2$

$i$	$a_{i0}$	$a_{i1}$	$\lambda_{i1}$	$\lambda_{i2}$	$\lambda_{i3}$	$b_{i0}$	$b_{i1}$
1	$\frac{84}{15}$	$-\frac{19}{15h}$	$\frac{4}{15h^3}$	$-\frac{9}{15h^3}$	$\frac{6}{15h^3}$	$\frac{4}{15}$	$-\frac{1}{15h}$
2	$\frac{8}{5}$	$\frac{8}{5h}$	$-\frac{3}{5h^3}$	$\frac{8}{5h^3}$	$-\frac{7}{5h^3}$	$\frac{8}{5}$	$\frac{2}{5h}$
3	$\frac{2}{5}$	$-\frac{2}{5h}$	$\frac{2}{5h^3}$	$-\frac{7}{5h^3}$	$\frac{8}{5h^3}$	$\frac{32}{5}$	$-\frac{8}{5h}$
4	$-\frac{1}{15}$	$\frac{1}{15h}$	$-\frac{1}{15h^3}$	$\frac{6}{15h^3}$	$-\frac{9}{15h^3}$	$-\frac{61}{15}$	$\frac{19}{15h}$

**Example 3.3.**

$n = 5, q = 2$

$i$	$a_{i0}$	$a_{i1}$	$\lambda_{i1}$	$\lambda_{i2}$	$\lambda_{i3}$	$\lambda_{i4}$	$b_{i0}$	$b_{i1}$
1	$\frac{127}{56}$	$-\frac{71}{56h}$	$\frac{15}{56h^3}$	$-\frac{34}{56h^3}$	$\frac{24}{56h^3}$	$-\frac{6}{56h^3}$	$\frac{5}{56}$	$\frac{1}{56h}$
2	$\frac{45}{28}$	$\frac{45}{28h}$	$-\frac{17}{28h^3}$	$\frac{46}{28h^3}$	$-\frac{44}{28h^3}$	$\frac{18}{28h^3}$	$\frac{15}{28}$	$-\frac{3}{28h}$
3	$\frac{3}{7}$	$-\frac{3}{7h}$	$\frac{3}{7h^3}$	$-\frac{11}{7h^3}$	$\frac{16}{7h^3}$	$-\frac{11}{7h^3}$	$\frac{15}{7}$	$\frac{3}{7h}$
4	$-\frac{3}{28}$	$\frac{3}{28h}$	$-\frac{3}{28h^3}$	$\frac{18}{28h^3}$	$-\frac{44}{28h^3}$	$\frac{46}{28h^3}$	$\frac{225}{28}$	$-\frac{45}{28h}$
5	$\frac{1}{56}$	$-\frac{1}{56h}$	$\frac{1}{56h^3}$	$-\frac{6}{56h^3}$	$\frac{24}{56h^3}$	$-\frac{34}{56h^3}$	$-\frac{355}{56}$	$\frac{71}{56h}$

**Example 3.4.**

$n = 4, q = 3$

$i$	$a_{i0}$	$a_{i1}$	$a_{i2}$	$\lambda_{i1}$	$\lambda_{i2}$	$\lambda_{i3}$	$b_{i0}$	$b_{i1}$	$b_{i2}$
1	$\frac{448}{132}$	$-\frac{409}{132h}$	$\frac{93}{132h^2}$	$\frac{2}{132h^5}$	$\frac{6}{132h^5}$	$\frac{6}{132h^5}$	$\frac{332}{132}$	$\frac{191}{132h}$	$-\frac{27}{132h^2}$
2	$-\frac{187}{44}$	$\frac{255}{44h}$	$-\frac{71}{44h^2}$	$\frac{2}{44h^5}$	$-\frac{6}{44h^5}$	$\frac{6}{44h^5}$	$\frac{596}{44}$	$-\frac{345}{44h}$	$\frac{49}{44h^2}$
3	$\frac{96}{44}$	$-\frac{145}{44h}$	$\frac{49}{44h^2}$	$-\frac{2}{44h^5}$	$\frac{6}{44h^5}$	$\frac{6}{44h^5}$	$-\frac{684}{44}$	$\frac{445}{44h}$	$-\frac{71}{44h^2}$
4	$-\frac{52}{132}$	$\frac{79}{132h}$	$-\frac{27}{132h^2}$	$\frac{2}{132h^5}$	$\frac{6}{132h^5}$	$\frac{6}{132h^5}$	$\frac{728}{132}$	$-\frac{521}{132h}$	$\frac{93}{132h^2}$

This concrete cardinal bases can be used as in (2.21), (2.22) for determining of optimal formulae for different linear operators.

**4. EXAMPLE OF NUMERICAL SOLUTION OF OPTIMUM CONTROL PROBLEM**

Let us minimize the integral cost function

$$\mathcal{J}(u) = \frac{1}{2} \int_0^T (x^2(t) + u^2(t)) dt,$$

where  $T > 0$  is given and

$$\begin{aligned} \dot{x}(t) &= -a x(t) + u(t), \quad a > 0, \\ x(0) &= x_0. \end{aligned}$$

Using the Pontrjagin principle, the Hamilton function is

$$\mathcal{H}(\psi, x, u) = \frac{1}{2}\psi_0(x^2 + u^2) + \psi_1(-ax + u),$$

where we can put  $\psi_0 = -1$ .

The optimal control is therefore

$$u^*(t) = \psi_1(t).$$

The function  $\psi_1(t)$  is the solution of the following boundary value problem:

$$\begin{aligned} \dot{x}(t) &= -a x(t) + u(t), \quad x(0) = x_0, \\ \dot{\psi}_1(t) &= x(t) + a \psi_1(t), \quad \psi_1(T) = 0, \end{aligned}$$

which solved can be analytically and there is

$$\psi_1(t) = x_0 \frac{e^{\lambda t} - e^{\lambda(2T-t)}}{\lambda - a + (\lambda + a)e^{2\lambda T}}, \quad \lambda = \sqrt{(a^2 + 1)}$$



This boundary value problem was discretized by means of the following formulae:

- (1) Optimal formula for  $f \in H^2$ , using the values of  $f(\cdot)$  in 3 points with  $h = x_{i+1} - x_i$ ,  $i = 0, 1, \dots, n-1$

$$f'(x_j) \doteq \frac{1}{4h} [-5f(x_j) + 6f(x_{j+1}) - f(x_{j+2})], \quad j = 1, \dots, n-1,$$

$$f'(x_j) \doteq \frac{1}{2h} [f(x_{j+1}) - f(x_{j-1})], \quad j = 1, \dots, n,$$

$$f'(x_j) \doteq \frac{1}{4h} [f(x_{j-2}) - 6f(x_{j-1}) + 5f(x_j)], \quad j = 2, \dots, n.$$

This formula was derived using the cardinal splines from Example 3.1. It is precise for polynomials of order 1.

- (2) Classical three-points formula derived using the Lagrange interpolation, precise for polynomials of order 2

$$f'(x_j) \doteq \frac{1}{2h} [-3f(x_j) + 4f(x_{j+1}) - f(x_{j+2})], \quad j = 1, \dots, n-1,$$

$$f'(x_j) \doteq \frac{1}{2h} [f(x_{j+1}) - f(x_{j-1})], \quad j = 1, \dots, n,$$

$$f'(x_j) \doteq \frac{1}{2h} [f(x_{j-2}) - 4f(x_{j-1}) + 3f(x_j)], \quad j = 2, \dots, n.$$

- (3) The common Euler method.

The results of computation (due by Z. Beran in [2]) for  $T = 1$ ,  $a = 1$ ,  $x_0 = 1$  with the step  $h = 0, 2$  on  $[0, 1]$  are given in the following table, where  $e(\cdot)$  is the error in comparison with the analytic solution. For  $u_i^* = \psi_{1i}$  we used the first, for  $x_i$  the third formula in (1) and (2). As the conditions for  $x_i$  are given at the initial point and those for  $u_i^*$  at the final point of the interval, this choice seems to be the simplest one, although evidently not the only possible way of application of discretization formulas.

(Received April 13, 1984.)

#### REFERENCES

- [1] H. C. Бахванов: Численные методы, Наука, Москва 1973.  
 [2] Z. Beran: Optimalizace diskrétního popisu spojitéch systémů v teorii regulace. Candidate of Sciences Thesis, ÚTIA ČSAV, Praha 1980.  
 [3] M. Golomb: Optimal approximating manifolds in  $L_2$ -spaces. J. Math. Anal. Appl. 12 (1965), 505–512.  
 [4] M. Golomb, and H. Weinberger: Optimal approximation and error bounds. In: On Numerical Approximation (R. E. Langer, ed.), Univ. of Wisconsin Press, Madison 1959, pp. 117–190.  
 [5] A. S. B. Holland and B. N. Sahney: The General Problem of Approximation and Spline Functions. Krieger Publishing Co., Huntington, N. Y. 1979.

$t_i$	Optimal formula					Lagrange's formula					Euler's method					
	$x_i$	$e(x_i)$	$u_i$	$e(u_i)$	$x_i$	$e(x_i)$	$u_i$	$e(u_i)$	$x_i$	$e(x_i)$	$u_i$	$e(u_i)$	$x_i$	$e(x_i)$	$u_i$	$e(u_i)$
0	1	0	-0.406	-0.020	1	0	-0.359	0.027	1	0	-0.334	0.052	1	0	-0.334	0.052
0.2	0.766	$6.9 \cdot 10^{-3}$	-0.293	-0.016	0.707	-0.052	-0.246	0.031	0.733	-0.026	-0.234	0.043	0.733	-0.026	-0.234	0.043
0.4	0.592	0.0124	-0.202	-0.012	0.509	-0.071	-0.163	0.027	0.539	-0.040	-0.158	0.032	0.539	-0.040	-0.158	0.032
0.6	0.463	0.016	-0.127	$-7.8 \cdot 10^{-3}$	0.367	-0.080	-0.099	0.019	0.400	-0.046	-0.097	0.022	0.400	-0.046	-0.097	0.022
0.8	0.368	0.018	-0.061	$-3.6 \cdot 10^{-3}$	0.267	-0.084	-0.048	$8.9 \cdot 10^{-3}$	0.301	-0.049	-0.046	0.011	0.301	-0.049	-0.046	0.011
1	0.302	0.019	—	—	0.206	-0.076	—	—	0.231	-0.050	—	—	0.231	-0.050	—	—

- [6] T. A. Kilgore: A characterization of the Lagrange interpolating projection with minimal Tchebycheff norm. *J. Approx. Theory* 24 (1978), 273–288.
- [7] P. J. Laurent: Approximation et optimisation. Hermann, Paris 1972.
- [8] T. Lyche and L. L. Schumaker: Local spline approximation methods. *J. Approx. Theory* 15 (1975), 294–325.
- [9] G. Meinardus: Approximation der Funktionen, Theorie und numerische Methoden. Springer-Verlag, Berlin–Göttingen–Heidelberg–New York 1964.
- [10] M. J. D. Powell: Approximation Theory and Methods. Cambridge University Press, Cambridge 1981.
- [11] A. Sard: Best approximate integration formulae; best approximation formulae. *Amer. J. Math.* 71 (1949), 80–91.
- [12] A. Sard: Optimal approximation. *J. Funct. Anal.* 1, 2 (1967), 222–244.
- [13] I. J. Schoenberg: Spline interpolation and best quadrature formulae. *Bull. Amer. Math. Soc.* 70 (1964), 143–148.
- [14] С. Л. Соболев, И. Бабушка: Оптимизация численных методов. *Аplikace matematiky* 10 (1965), 2, 96–129.
- [15] С. Л. Соболев: Введение в теорию кубатурных формул. Наука, Москва 1974.
- [16] J. Stoer and R. Bulirsch: Introduction to Numerical Analysis. Springer-Verlag, Berlin–Heidelberg–New York 1980.
- [17] A. Tuzar: Optimal Approximation of Linear Functionals in Hilbert Space with Applications to Numerical Methods. Research Report No. 988, ÚTIA ČSAV, 1979.
- [18] A. Tuzar and Z. Beran: Optimization of the discrete description for continuous systems. *Problems Control Inform. Theory* 10 (1981), 2, 83–94.
- [19] J. F. Traub and H. Wozniakowski: A General Theory of Optimal Algorithms. Academic Press, New York 1980.
- [20] R. Varga: Functional Analysis and Approximation Theory in Numerical Analysis. Society for Industrial and Applied Mathematics, Philadelphia, Pennsylvania 1971.

*RNDr. Antonín Tuzar, CSc., Ústav teorie informace a automatizace ČSAV (Institute of Information Theory and Automation – Czechoslovak Academy of Sciences), Pod vodárenskou věží 4, 182 08 Praha 8, Czechoslovakia.*