

Kybernetika

VOLUME 45 (2009), NUMBER 3

The Journal of the Czech Society for
Cybernetics and Information Sciences

Published by:

Institute of Information Theory and Automation of the AS CR

Editorial Office:

Pod Vodárenskou věží 4, 182 08 Praha 8

Editor-in-Chief:

Milan Mareš

Managing Editors:

Lucie Fajfrová
Karel Sladký

Editorial Board:

Jiří Anděl, Sergej Čelikovský, Marie Demlová, Jan Flusser, Petr Hájek, Vladimír Havlena, Didier Henrion, Yiguang Hong, Zdeněk Hurák, Martin Janžura, Jan Ježek, George Klir, Ivan Kramosil, Tomáš Kroupa, Petr Lachout, Friedrich Liese, Jean-Jacques Loiseau, František Matúš, Radko Mesiar, Karol Mikula, Jiří Outrata, Jan Seidler, Karel Sladký, Jan Štecha, Olga Štěpánková, František Turnovec, Igor Vajda, Jiřina, Vojnarová, Milan Vlach, Miloslav Vošvrda, Pavel Zítek

Kybernetika is a bi-monthly international journal dedicated for rapid publication of high-quality, peer-reviewed research articles in fields covered by its title.

Kybernetika traditionally publishes research results in the fields of Control Sciences, Information Sciences, System Sciences, Statistical Decision Making, Applied Probability Theory, Random Processes, Fuzziness and Uncertainty Theories, Operations Research and Theoretical Computer Science, as well as in the topics closely related to the above fields.

The Journal has been monitored in the Science Citation Index since 1977 and it is abstracted/indexed in databases of Mathematical Reviews, Zentralblatt für Mathematik, Current Mathematical Publications, Current Contents ISI Engineering and Computing Technology.

Kybernetika. Volume 45 (2009)

ISSN 0023-5954, MK ČR E 4902.

Published bimonthly by the Institute of Information Theory and Automation of the Academy of Sciences of the Czech Republic, Pod Vodárenskou věží 4, 182 08 Praha 8. — Address of the Editor: P. O. Box 18, 182 08 Prague 8, e-mail: kybernetika@utia.cas.cz. — Printed by PV Press, Pod vrstevnicí 5, 140 00 Prague 4. — Orders and subscriptions should be placed with: MYRIS TRADE Ltd., P. O. Box 2, V Štíhlách 1311, 142 01 Prague 4, Czech Republic, e-mail: myris@myris.cz. — Sole agent for all “western” countries: Kubon & Sagner, P. O. Box 34 01 08, D-8 000 München 34, F.R.G.

Published in June 2009.

© Institute of Information Theory and Automation of the AS CR, Prague 2009.

PURE FILTERS AND STABLE TOPOLOGY ON BL-ALGEBRAS

ESFANDIAR ESLAMI AND FARHAD KH. HAGHANI

In this paper we introduce stable topology and F -topology on the set of all prime filters of a BL-algebra A and show that the set of all prime filters of A , namely $\text{Spec}(A)$ with the stable topology is a compact space but not T_0 . Then by means of stable topology, we define and study pure filters of a BL-algebra A and obtain a one to one correspondence between pure filters of A and closed subsets of $\text{Max}(A)$, the set of all maximal filters of A , as a subspace of $\text{Spec}(A)$. We also show that for any filter F of BL-algebra A if $\sigma(F) = F$ then $U(F)$ is stable and F is a pure filter of A , where $\sigma(F) = \{a \in A \mid y \wedge z = 0 \text{ for some } z \in F \text{ and } y \in a^\perp\}$ and $U(F) = \{P \in \text{Spec}(A) \mid F \not\subseteq P\}$.

Keywords: BL-algebra, prime filters, maximal filters, pure filters, stable topology, F -topology

AMS Subject Classification: 03G25, 06F99, 08A72

1. INTRODUCTION

L.P. Belluce and S. Sessa studied in [2] stable topology and pure ideals in the framework of MV-algebras. They defined the stable topology for MV-algebras as follows: let A be an MV-algebra. The set of all prime ideals of A is denoted by $\text{Spec}(A)$. The open sets in $\text{Spec}(A)$ are of the form $U(I) = \{P \in \text{Spec}(A) \mid I \not\subseteq P\}$ where I is an ideal of A . The set $U(I)$ is stable under ascent if $P \in U(I)$ and $Q \in \text{Spec}(A)$ with $P \subseteq Q$, then $Q \in U(I)$. The set $U(I)$ is stable under descent if $P \in U(I)$ and $Q \in \text{Spec}(A)$ with $Q \subseteq P$, then $Q \in U(I)$.

$U(I)$ is said to be stable if it is stable under ascent and under descent. The stable topology for A is the collection of stable open subsets of $\text{Spec}(A)$.

In 1998 Petr Hájek introduced in [4] the variety of BL-algebras and showed that the variety of MV-algebras actually is a subvariety of the variety of BL-algebras. In other words, any MV-algebra can be easily viewed as a special BL-algebra. Thus it makes sense to generalize the notion of stable topology to BL-algebras. But in fact since the multiplication (\odot) is a fundamental operation and filters are basic notions in BL-algebras defined in terms of \odot (see the Definitions 2.1, 2.2 below) as well as a dual notion of ideals, we prefer to present stable topology based on filters. Therefore the generalization does not work easily and we face some related difficulties towards

this approach. Although we get similar results as in [2], we also prove some more theorems regarding different properties of this topology on BL-algebras.

This paper consists of four sections. In the second section we recall the definition of a BL-algebra A , a filter F and $\text{Spec}(A)$ with more preliminary facts that we need in the sequel. In the third section we define F -topology which is actually the same as spectral topology but in terms of filters (letter F comes from the word filter) and introduce the stable topology on $\text{Spec}(A)$. We show that the topological space $\text{Spec}(A)$ with the stable topology is compact but not T_0 and hence neither T_1 nor T_2 .

In the fourth section, we define pure filters of A and prove some important results. In fact let $\text{Max}(A)$ be the set of all maximal filters of A . Since $\text{Max}(A) \subseteq \text{Spec}(A)$, we consider the topology induced by F -topology on $\text{Max}(A)$ and show that F -topology and stable topology coincide on subspace $\text{Max}(A)$. We show that pure filters of A are in one to one correspondence with closed subsets of $\text{Max}(A)$. We also investigate some conditions for purity of a filter F by considering $\sigma(F) = \{a \in A \mid y \wedge z = 0 \text{ for some } z \in F \text{ and } y \in a^\perp\}$ and stability of $U(F)$ where $U(F)$ is an open set in $\text{Spec}(A)$ with F -topology.

2. PRELIMINARIES

Definition 2.1. (Hájek [6]) A BL-algebra is an algebra $A = (A, \vee, \wedge, \odot, \longrightarrow, 0, 1)$ of type $(2,2,2,2,0,0)$ satisfying the following properties:

1. $(A, \vee, \wedge, 0, 1)$ is a lattice with 0 as the least element and 1 as the greatest element.
2. $(A, \odot, 1)$ is a commutative monoid.
3. The following statements hold for every $a, b, c \in A$:
 - (i) $c \leq a \longrightarrow b$ iff $a \odot c \leq b$ (Residuation);
 - (ii) $a \wedge b = a \odot (a \longrightarrow b)$ (Divisibility);
 - (iii) $(a \longrightarrow b) \vee (b \longrightarrow a) = 1$ (Prelinearity).

A BL-algebra A is nontrivial iff $0 \neq 1$. We also define a unary operation “ $-$ ” on A by $a \longrightarrow 0 = \bar{a}$

Definition 2.2. (Hájek [6]) A filter of a BL-algebra A is a nonempty subset F of A such that:

- (i) $a, b \in F$ implies $a \odot b \in F$;
- (ii) $a \in F$ and $a \leq b$ imply $b \in F$.

Definition 2.3. (Hájek [6]) A filter F of a BL-algebra A is proper if $F \neq A$. A proper filter P of A is called prime provided that $a \vee b \in P$ implies that $a \in P$ or $b \in P$, for every $a, b \in A$.

A proper filter M of A is called maximal, if it is not contained in any other proper filter, that is for any filter E such that $M \subseteq E \subseteq A$, either $E = M$ or $E = A$.

A BL-algebra A is local if it has a unique maximal filter.

It is easy to see that F is a proper filter iff $0 \notin F$.

Remark 2.1. The usual notions of morphisms can be defined on BL-algebras (see for example [4, 11]).

Proposition 2.1. (Georgescu and Leustean [5]) Let $h : A \rightarrow B$ be a BL-morphism. Then

- (i) If G is a (proper, prime, maximal) filter of B , then $h^{-1}(G)$ is a (proper, prime, maximal) filter of A .
- (ii) If h is surjective and F is a filter of A , then $h(F)$ is a filter of B .
- (iii) If h is surjective and M is a maximal filter of A such that $h(M)$ is proper, then $h(M)$ is a maximal filter of B .

We denote the lattice reduct of a BL-algebra A by $L(A)$, and it is easy to see that any (prime) filter of A is a (prime) filter of $L(A)$.

From now on, in this paper we consider $\text{Spec}(A)$, $\text{Max}(A)$ and $\text{Min}(A)$ as the set of all prime filters, maximal filters and minimal prime filters of a BL-algebra A , respectively.

Proposition 2.2. (Di Nola et al. [4], Leustean [8], Turunen [10]) Let A be a BL-algebra. Then the followings hold.

- (i) If F is a filter of A and S is a nonempty \vee -closed subset of A , (i.e. if $a, b \in S$ then $a \vee b \in S$) such that $F \cap S = \emptyset$, then there exists a prime filter P of A such that $F \subseteq P$ and $P \cap S = \emptyset$.
- (ii) Any maximal filter of A is a prime filter.
- (iii) If A is nontrivial, then any proper filter F of A is the intersection of all prime filters containing F .
- (iv) If A is nontrivial, then any prime filter of A is contained in a unique maximal filter.
- (v) If A is nontrivial, then any proper filter A can be extended to a prime, maximal filter.

Proposition 2.3. (Di Nola et al. [4]) If A is a nontrivial BL-algebra and M a proper filter of A , then the following are equivalent:

- (i) M is maximal,
- (ii) For any $x \in A$, $x \notin M$ implies that $\overline{x^n} \in M$ for some $n \in \omega$, where ω is the set of natural numbers.

Definition 2.4. (Leustean [8]) Let $X \subseteq A$. The filter generated by X will be denoted by $\langle X \rangle$. If $X = \emptyset$ then $\langle \emptyset \rangle = \{1\}$ and if $X \neq \emptyset$ then we have $\langle X \rangle = \{y \in A \mid x_1 \odot x_2 \odot \cdots \odot x_n \leq y \text{ for some } n \in \omega \text{ and some } x_1, x_2, \dots, x_n \in X\}$.

It is easy to see that if $a \in A$ then we have $\langle a \rangle = \{b \in A \mid a^n \leq b \text{ for some } n \in \omega\}$.

Proposition 2.4. (Leustean [8]) If $F(A)$ is the set of all filters of A , then $(F(A), \subseteq)$ is a complete lattice and for every family $\{F_i\}$ of filters of A , we have $\bigvee_{i \in I} F_i = \langle \bigcup_{i \in I} F_i \rangle$ and $\bigwedge_{i \in I} F_i = \bigcap_{i \in I} F_i$.

From [6] it follows that to every filter F of a BL-algebra A we can associate a congruence relation \sim_F on A by defining $a \sim_F b$ iff $a \longrightarrow b \in F$ and $b \longrightarrow a \in F$ iff $(a \longrightarrow b) \odot (b \longrightarrow a) \in F$. For element $a \in A$, let $\frac{a}{F}$ be the congruence class $\frac{a}{\sim_F}$. If we denote by $\frac{A}{F}$ the quotient set $\frac{A}{\sim_F}$, then $\frac{A}{F}$ becomes a BL-algebra with the natural operations induced by those of A .

We recall that a BL-chain is a totally ordered BL-algebra, i.e. a BL-algebra whose lattice order is total [6].

Proposition 2.5. (Hájek [6]) Let F be a filter of A and $a, b \in A$. Then

- (i) $\frac{a}{F} = \frac{1}{F}$ iff $a \in F$.
- (ii) $\frac{a}{F} = \frac{0}{F}$ iff $\bar{a} \in F$.
- (iii) $\frac{a}{F} \leq \frac{b}{F}$ iff $a \longrightarrow b \in F$.
- (iv) $\frac{A}{F}$ is a BL-chain iff F is a prime filter of A .

Definition 2.5. (Busneage and Piciu [3]) Let A be a BL-algebra. An element $a \in A$ is called archimedean if there is $n \in \omega$, $n \geq 1$ such that $a \vee \bar{a}^n = 1$. A BL-algebra A is called hyperarchimedean if all its elements are archimedean.

Proposition 2.6. (Busneage and Piciu [3]) A BL-algebra A is hyperarchimedean iff $\text{Spec}(A) = \text{Max}(A)$.

Based on the definitions and propositions in this section, we define our main notion of stable topology on BL-algebras.

3. STABLE TOPOLOGY

Let A be a non trivial BL-algebra. Denote by $\text{Spec}(A)$ the set of all its prime filters. Consider the spectral topology (Zariski topology) on $\text{Spec}(A)$, i.e. the topology in which its closed sets are exactly the sets of the form $V(X) = \{P \in \text{Spec}(A) \mid X \subseteq P\}$ for each subset X of A . Then $\text{Spec}(A)$ equipped with this topology is called the prime spectrum of A .

Now we are planning to introduce F -topology on $\text{Spec}(A)$.

Proposition 3.1. Let A be a nontrivial BL-algebra and F be a filter of A . Define $V(F) = \{P \in \text{Spec}(A) \mid F \subseteq P\}$. Then the following hold:

- (i) $V(\{1\}) = \text{Spec}(A)$, $V(A) = \emptyset$.
- (ii) If $\{F_i\}_{i \in I}$ is a family of filters of A , then $\bigcap_{i \in I} V(F_i) = V(\langle \bigcup_{i \in I} F_i \rangle)$.
- (iii) If F_1, F_2 are filters of A then $V(F_1) \cup V(F_2) = V(F_1 \cap F_2)$.

Proof.

- (i) Follows from the fact that 1 belongs to any filter F and every prime filter P is proper.
- (ii) Since each $F_i \subseteq \bigcup_{i \in I} F_i \subseteq \langle \bigcup_{i \in I} F_i \rangle$, $V(\langle \bigcup_{i \in I} F_i \rangle) \subseteq V(F_i)$ for each $i \in I$. Then $V(\langle \bigcup_{i \in I} F_i \rangle) \subseteq \bigcap_{i \in I} V(F_i)$. Now let $P \in \bigcap_{i \in I} V(F_i)$ then $P \in V(F_i)$ and $F_i \subseteq P$ for each $i \in I$. We claim that $\langle \bigcup_{i \in I} F_i \rangle \subseteq P$. Let $t \in \langle \bigcup_{i \in I} F_i \rangle$. Then $t \geq f_1 \odot f_2 \odot \dots \odot f_k$ for some $k \in \omega$ and $f_1, f_2, \dots, f_k \in \bigcup_{i \in I} F_i$. But for each f_i there exists F_{k_i} such that $f_i \in F_{k_i}$. Therefore, $f_1, f_2, \dots, f_k \in P$ and since P is a filter, $t \in P$. Thus, we conclude that $P \in V(\langle \bigcup_{i \in I} F_i \rangle)$.
- (iii) Since $F_1 \cap F_2 \subseteq F_1, F_2$ we have $V(F_1), V(F_2) \subseteq V(F_1 \cap F_2)$. Thus $V(F_1) \cup V(F_2) \subseteq V(F_1 \cap F_2)$. Now let $P \in V(F_1 \cap F_2)$ but $P \notin V(F_1) \cup V(F_2)$. Then $P \not\subseteq V(F_1)$, $P \not\subseteq V(F_2)$, i. e. $F_1 \not\subseteq P$, $F_2 \not\subseteq P$. There exist $x \in F_1$, $y \in F_2$ such that $x, y \notin P$. Since $x, y \leq x \vee y$, $x \vee y \in F_1, F_2$ and hence $x \vee y \in F_1 \cap F_2$. But since $F_1 \cap F_2 \subseteq P$, $x \vee y \in P$. This implies $x \in P$ or $y \in P$ which contradicts the assumption. \square

Based on Proposition 3.1, we have

Corollary 3.1. The collection $\{V(F) \mid F \text{ is a filter of } A\}$ defines a topology on $\text{Spec}(A)$ whose closed sets are of the form $V(F)$ for some filter F in A .

We call the resulting topology in Corollary 3.1, *F-topology*.

Remark 3.1. From [8] since the family $\{U(a)\}_{a \in A}$ where $U(a) = \{P \in \text{Spec}(A) \mid a \notin P\}$, is a basis for the spectral topology on $\text{Spec}(A)$, this family is also a basis for *F-topology*. For let F be a filter of A . Then by [8, Proposition 2.2], we have $U(F) = U(\bigcup_{f \in F} \{f\}) = \bigcup_{f \in F} U(f)$ and hence any open subset of $\text{Spec}(A)$ with *F-topology* is the union of subsets from the family $\{U(a)\}_{a \in A}$. Thus by [8, Theorem 2.7], $\text{Spec}(A)$ with *F-topology* is also a compact T_0 topological space.

It is obvious to see that if F is a filter of A , then $V(F)$ is *stable under ascent*, that is if $P \in V(F)$, $Q \in \text{Spec}(A)$ and $P \subseteq Q$, then $Q \in V(F)$.

We also know that $U(F) = \text{Spec}(A) - V(F) = \{P \in \text{Spec}(A) \mid F \not\subseteq P\}$ is *stable under descent*, i. e. if $P \in U(F)$, $Q \in \text{Spec}(A)$ and $Q \subseteq P$, then $Q \in U(F)$.

Let A be a BL-algebra and F be a filter of A . We say that $U(F)$ is *stable* if $U(F)$ is stable under ascent and descent. Since $U(F)$ is always stable under descent, being stable it is enough that $U(F)$ is stable under ascent.

Remark 3.2. If F, G are filters of a nontrivial BL-algebra A . Then $F = G$ iff $U(F) = U(G)$.

Proof. If $F = G$, then obviously $U(F) = U(G)$. Now let $U(F) = U(G)$. Thus $V(F) = V(G)$ and $\bigcap_{P \in \text{Spec}(A), F \subseteq P} P = \bigcap_{P \in \text{Spec}(A), G \subseteq P} P$. Therefore by Proposition 2.2 (iii), $F = G$. \square

In the following proposition we introduce the stable topology on $\text{Spec}(A)$.

Proposition 3.2. The collection of all stable open subsets of $\text{Spec}(A)$ satisfies the axioms for open sets in a topological space. The resulting topology is called stable topology on $\text{Spec}(A)$. In other words, $\{U \mid U \text{ is open with } F\text{-topology and stable}\}$ is the collection of open sets for stable topology.

Proof. Let T be the set of all stable open subsets of $\text{Spec}(A)$. It is obvious that \emptyset and $\text{Spec}(A) \in T$. Now let T_1, T_2 be in T . Then $T_1 = U(F_1)$ and $T_2 = U(F_2)$ for some $F_1, F_2 \in F(A)$. Since $U(F_1) \cap U(F_2) = U(\langle F_1 \cup F_2 \rangle)$, $T_1 \cap T_2$ is open. For stability, It is enough to show that $T_1 \cap T_2$ is stable under ascent. Let $P \in U(\langle F_1 \cup F_2 \rangle)$, $Q \in \text{Spec}(A)$ and $P \subseteq Q$. Then $P \in U(F_1), U(F_2)$ and hence by stability of $U(F_1), U(F_2)$ we have, $Q \in U(F_1), U(F_2)$. Thus $Q \in U(F_1) \cap U(F_2)$.

Let $\{T_i\}_{i \in I}$ be a family of stable open subsets of $\text{Spec}(A)$. Then for each $i \in I$, there exists filter F_i of A such that $T_i = U(F_i)$. Thus $\bigcup_i T_i = \bigcup_i U(F_i) = U(\bigcap_i F_i)$. For stability, let $P \in U(\bigcap_i F_i)$, $Q \in \text{Spec}(A)$ and $P \subseteq Q$, then $P \in U(F_i)$ for some $i \in I$ and by stability of $U(F_i)$ we have $Q \subseteq U(F_i) \subseteq \bigcup_i U(F_i)$ and hence $Q \in \bigcup_i U(F_i)$. \square

In the next corollary, we see that there is a distinction between topological property of $\text{Spec}(A)$ with stable topology and F -topology. In fact, $\text{Spec}(A)$ with stable topology is a T_0 topological space but with F -topology is not.

Corollary 3.2. With the stable topology, $\text{Spec}(A)$ is a compact topological space but not T_0 and hence neither T_1 nor T_2 .

Proof. We know that every stable open set is also open in F -topology. Therefore, since by Remark 3.1, $\text{Spec}(A)$ is compact in F -topology, it is also compact in stable topology. Now let $P, Q \in \text{Spec}(A)$ such that $P \not\subseteq Q$. Since all open sets ($U(F)$ for some filter F of A) are stable under descent, every $U(F)$ that contains Q , will contain P . Now suppose that $U(F)$ is stable, and $P \in U(F)$. Then since Q contains P , and $U(F)$ is stable, we have $Q \in U(F)$. Hence we see that P and Q can not be separated by stable open sets, so the stable topology is not T_0 and therefore is neither T_1 nor T_2 . \square

Lemma 3.1. Let A be a nontrivial BL-algebra, $M \in \text{Max}(A)$ and F be a proper filter of A . Suppose that $\widehat{W}_M = \{P \in \text{Spec}(A) \mid P \subseteq M\}$ and $W_M = \bigcap \widehat{W}_M = \bigcap_P P$ for $P \in \widehat{W}_M$ and $W_M \subseteq F$. Then $F \subseteq M$.

Proof. Let $W_M \subseteq F$ but $F \not\subseteq M$. Therefore by Proposition 2.3, there exists $x \in F$ such that $x \notin M$ and $\overline{x^n} \in M$ for some $n \in \omega$. Let $P \in \widehat{W}_M$ be arbitrary. Then $\frac{M}{P} \in \text{Max}\left(\frac{A}{P}\right)$. Since P is a prime filter, by Proposition 2.5, $\frac{A}{P}$ is linear ordered and we have $\frac{x^n}{P} < \frac{\overline{x^n}}{P}$ or $\frac{\overline{x^n}}{P} < \frac{x^n}{P}$. If $\frac{x^n}{P} < \frac{\overline{x^n}}{P}$, then $\frac{x^n}{P} \odot \frac{x^n}{P} < \frac{\overline{x^n}}{P} \odot \frac{x^n}{P} = \frac{0}{P}$, i.e. $\frac{x^{2n}}{P} = \frac{0}{P}$. Thus $(x^{2n} \rightarrow 0) \in P$, i.e. $\overline{x^{2n}} \in P$. But since P is arbitrary in \widehat{W}_M , $\overline{x^{2n}} \in \bigcap_P P = W_M$ for $P \in \widehat{W}_M$. Since $W_M \subseteq F$, $\overline{x^{2n}} \in F$. But by assumption $x \in F$. Therefore $x^{2n} \in F$ and hence $0 \in F$ which is a contradiction. Now if $\frac{\overline{x^n}}{P} < \frac{x^n}{P}$, since $\overline{x^n} \in M$, $\frac{\overline{x^n}}{P} \in \frac{M}{P} \in \text{Max}\left(\frac{A}{P}\right)$ and hence $\frac{x^n}{P} \in \frac{M}{P}$, i.e. $x^n \in M$. Thus $x^n \odot \overline{x^n} = 0 \in M$ which is a contradiction. \square

Proposition 3.3. With the notation of Lemma 3.1 the following hold:

- (i) $V(W_M) = \widehat{W}_M$ and hence \widehat{W}_M is closed with respect to F -topology.
- (ii) $\text{Spec}(A)$ is the disjoint union of subspaces \widehat{W}_M , $M \in \text{Max}(A)$.

Proof.

- (i) Let $P \in V(W_M)$. Then $W_M \subseteq P$ and by Lemma 3.1, $P \subseteq M$, i.e. $P \in \widehat{W}_M$. Obviously $\widehat{W}_M \subseteq V(W_M)$ and hence \widehat{W}_M is a closed set.
- (ii) The proof follows from [1, p. 333]. \square

4. PURE FILTERS AND SOME RESULTS

In this section we introduce pure filters and study some of their properties.

Definition 4.1. Let F be a filter of A . We say that F is *pure* if $U(F)$ is stable.

It is easy to see that if A is a BL-algebra, then A and $\{1\}$ are pure filters.

Lemma 4.1. Let F be a pure filter of A , $P \in \text{Spec}(A)$, $M \in \text{Max}(A)$ and $P, F \subseteq M$. Then $F \subseteq P$.

Proof. Assume on the contrary that $F \not\subseteq P$. Then $P \in U(F)$. Since $P \subseteq M$, $M \in \text{Max}(A)$ and $\text{Max}(A) \subseteq \text{Spec}(A)$, by stability of $U(F)$, we conclude that $M \in U(F)$, i.e. $F \not\subseteq M$ and this is a contradiction. \square

Corollary 4.1. Let F be a pure filter of A , $M \in \text{Max}(A)$ and $F \subseteq M$. Then $F \subseteq W_M$.

Proof. Let P be an arbitrary prime filter such that $P \subseteq M$. Since $F \subseteq M$ and $U(F)$ is stable, by Lemma 4.1, $F \subseteq P$. Thus $F \subseteq \bigcap_P P$ for each $P \subseteq M$ and hence $F \subseteq W_M$. \square

Theorem 4.1. Let F be a pure filter of A . Then $F = \bigcap_M \{W_M \mid F \subseteq M \in \text{Max}(A)\}$.

Proof. Let $U(F)$ be stable. Then by Corollary 4.1, for each $M \in \text{Max}(A)$ such that $F \subseteq M$, we have $F \subseteq W_M$ and hence $F \subseteq \bigcap_{M \supseteq F} W_M$. Now let $x \in \bigcap_M \{W_M \mid F \subseteq M \in \text{Max}(A)\}$. Then we have $x \in W_M$ for each $M \in \text{Max}(A)$ such that $F \subseteq M$. Let P be any prime filter of A such that $F \subseteq P$. By Proposition 2.2 there exists a unique maximal filter M_P over P , i.e. $P \subseteq M_P$. Therefore $F \subseteq M_P$ and by assumption $x \in W_{M_P}$. Since $V(W_M) = \widehat{W}_M$ and $P \in \widehat{W}_{M_P}$, $P \in V(W_M)$, i.e. $W_{M_P} \subseteq P$ and hence $x \in P$. Since P is an arbitrary prime filter such that $F \subseteq P$, $x \in \bigcap_{F \subseteq P} P$ and by Proposition 2.2 since $F = \bigcap_{F \subseteq P} P$, $x \in F$. \square

We are planning to obtain a relation between subsets of $\text{Max}(A)$ and stable open sets.

Theorem 4.2. Let F be a filter of A such that $U(F)$ is stable. Let $T = \{M \in \text{Max}(A) \mid F \subseteq M\}$. Then $U(F) = \text{Spec}(A) - \bigcup_{M \in T} \widehat{W}_M$.

Proof. Let $P \in U(F)$. Then $F \not\subseteq P$ and by Lemma 4.1, $F \not\subseteq M$. Let M_P be the unique maximal filter such that $P \subseteq M_P$. Then $F \not\subseteq M_P$, i.e. $M_P \notin T$ and hence $P \in \widehat{W}_{M_P} \subseteq \bigcup_{M \in \text{Max}(A)-T} \widehat{W}_M$. Since $\text{Spec}(A) - \bigcup_{M \in T} \widehat{W}_M = \bigcup_{M \in \text{Max}(A)} \widehat{W}_M - \bigcup_{M \in T} \widehat{W}_M = \bigcup_{M \in \text{Max}(A)-T} \widehat{W}_M$, we have $U(F) \subseteq \text{Spec}(A) - \bigcup_{M \in T} \widehat{W}_M$. Now let $P \in \text{Spec}(A) - \bigcup_{M \in T} \widehat{W}_M$ but $P \notin U(F)$. Then $P \notin \bigcup_{M \in T} \widehat{W}_M$, i.e. $P \notin \widehat{W}_M$ and $P \not\subseteq M$ for each $M \in T$. Since $P \notin U(F)$, $F \subseteq P$ and hence $F \subseteq M_P$, i.e. $M_P \in T$ but by above $P \not\subseteq M_P$ therefore, this is impossible. \square

Remark 4.1. For any $T \subseteq \text{Max}(A)$, $X_T = \text{Spec}(A) - \bigcup_{M \in T} \widehat{W}_M$ is stable.

Proof. It is enough to show that X_T is stable under ascent. Let $P \in X_T$, $Q \in \text{Spec}(A)$, $P \subseteq Q$. Then, $P \notin \bigcup_{M \in T} \widehat{W}_M$, i.e. $P \notin \widehat{W}_M$ for each $M \in T$. This means that $P \not\subseteq M$ for each $M \in T$. Thus $Q \not\subseteq M$ for each $M \in T$. That is equivalent to $Q \notin \widehat{W}_M$ for each $M \in T$. Hence $Q \in X_T$. \square

Theorem 4.3. Let T be a finite subset of $\text{Max}(A)$ and F be a filter of A such that $F = \bigcap_{M \in T} W_M$. Then $U(F)$ is stable, i.e. F is a pure filter.

Proof. Let $X = \text{Spec}(A) - \bigcup_{M \in T} \widehat{W}_M$. Since each \widehat{W}_M is closed and T is finite, the finite union of \widehat{W}_M is closed and hence X is an open set. By Remark 4.1, it is enough to show that $X = U(F)$. Let $P \in X$ then $P \notin \widehat{W}_M$ for each $M \in T$. Since $V(W_M) = \widehat{W}_M$, $P \notin V(W_M)$, i.e. $W_M \not\subseteq P$ for each $M \in T$ and hence $\bigcap_{M \in T} W_M \not\subseteq P$, which implies that $P \in U(\bigcap_{M \in T} W_M) = U(F)$. Now let $P \in U(F)$. Then $F \not\subseteq P$, i.e. $\bigcap_{M \in T} W_M \not\subseteq P$. This implies that $W_M \not\subseteq P$ for each $M \in T$. But since $P \notin V(W_M)$, $P \notin \widehat{W}_M$ for each $M \in T$ and hence $P \in \text{Spec}(A) - \bigcup_{M \in T} \widehat{W}_M$. \square

We recall that a BL-algebra A is semilocal iff $\text{Max}(A)$ is a finite set [5], then:

Corollary 4.2. A BL-algebra A is semilocal iff the stable topology on $\text{Spec}(A)$ is finite.

Proof. By Theorems 4.2 and 4.3, there is a relation between subsets of $\text{Max}(A)$ and stable open sets. Let A be semilocal. Then, there are only finitely many maximal filters and hence only finitely many stable open sets. Thus the stable topology is finite. Now let the stable topology on $\text{Spec}(A)$ be finite. Then, there can be only finitely many maximal filters and hence A is semilocal. \square

In the following theorems we provide needed facts to obtain a one to one correspondence between pure filters and closed subsets of $\text{Max}(A)$.

Theorem 4.4. The map $\varphi : \text{Spec}(A) \longrightarrow \text{Max}(A)$ by $P \longmapsto M_P$ is a continuous retraction.

Proof. From [8], it is known that associated to each BL-algebra A , there is a bounded distributive lattice $\beta(A)$ such that the topological space $\text{Spec}(A)$ and $\text{Spec}(\beta(A))$ are homeomorphic. On the other hand, by ([7] p.68), if A is a normal distributive lattice, then the map $\varphi : \text{Spec}(A) \longrightarrow \text{Max}(A)$ is a continuous retraction. Thus the Theorem holds for BL-algebras. \square

Corollary 4.3. The stable topology and F -topology coincide on subspace $\text{Max}(A)$.

Proof. We know that if $T_1 = \{U_i \mid i \in I\}$ is stable topology for $\text{Spec}(A)$ then $T_2 = \{U_i \cap \text{Max}(A) \mid i \in I\}$ is stable topology for $\text{Max}(A)$. Let G be an open set of $\text{Max}(A)$ and $Q = \{P \in \text{Spec}(A) \mid M_P \in G\}$ where M_P is the unique maximal filter of A such that $P \subseteq M_P$. Since $\varphi : \text{Spec}(A) \longrightarrow \text{Max}(A)$ by $P \longmapsto M_P$ is a continuous retraction, $\varphi^{-1}(G)$ is open in $\text{Spec}(A)$. But $\varphi^{-1}(G) = Q$. Hence Q is open in $\text{Spec}(A)$. Now we claim that Q is stable. Let $P_1 \in Q$, $P_2 \in \text{Spec}(A)$ such that $P_1 \subseteq P_2$. Let M_{P_2} be the unique maximal filter such that $P_2 \subseteq M_{P_2}$. Since $P_1 \subseteq M_{P_1}$ and $P_2 \subseteq M_{P_2}$, $M_{P_1} = M_{P_2}$. But $M_{P_1} \in G$ implies $M_{P_2} \in G$, i. e. $P_2 \in Q$. Therefore Q is stable and clearly $G = Q \cap \text{Max}(A)$. \square

Corollary 4.4. Let A be a nontrivial BL-algebra. Then every stable open subset G has the form $G = \bigcup_{M \in Y} \widehat{W}_M$ for some open subset $Y \subseteq \text{Max}(A)$.

Proof. Let G be a stable open set. We take $Y = G \cap \text{Max}(A)$. Then if we consider the map $\varphi : \text{Spec}(A) \longrightarrow \text{Max}(A)$ by $P \longmapsto M_P$, then it is trivial that $\varphi^{-1}(Y) = G$. But $\text{Spec}(A) = \bigcup_{M \in \text{Max}(A)} \widehat{W}_M$. Therefore, $G = \bigcup_{M \in Y} \widehat{W}_M$. \square

Remark 4.2. If G is open in $\text{Spec}(\widehat{A})$ and G is a union of closed sets in $\text{Spec}(A)$, then G is stable.

Proof. Let $G = \bigcup_{M \in T \subseteq \text{Max}(A)} \widehat{W}_M$, $P \in G$, $Q \in \text{Spec}(A)$ and $P \subseteq Q$. Since $Q \subseteq M_Q$ (unique maximal filter over Q), $Q \in \widehat{W}_{M_Q} \subseteq \bigcup_{M \in \text{Max}(A)} \widehat{W}_M$. Thus G is stable. \square

Theorem 4.5. Let A be a nontrivial BL-algebra, $T \subseteq \text{Max}(A)$, $Y = \bigcup_{M \in T} \widehat{W}_M$ be closed in $\text{Spec}(A)$. Then $E = \bigcap_{M \in T} W_M$ is a pure filter.

Proof. Let $E = \bigcap_{M \in T} W_M$. We know that $Y \cap \text{Max}(A) = \bigcup_{M \in T} \widehat{W}_M \cap \text{Max}(A) = \bigcup_{M \in T} \{P \mid P \in \text{Max}(A), P \subseteq M\} = \bigcup_{M \in T} M = T$. Since Y is closed, $\text{Spec}(A) - Y$ is open and stable. Thus there exists a pure filter F of A such that $\text{Spec}(A) - Y = U(F)$. It is enough to show that $E = F$. Since $U(F)$ is stable, by Theorem 4.1, $F = \bigcap_{M \supseteq F, M \in \text{Max}(A)} W_M = \bigcap_{M \supseteq F} (\bigcap_{P \subseteq M} P) \subseteq \bigcap_{P \subseteq M} P = W_M$ for each $M \in Y \cap \text{Max}(A) = T$. Then $F \subseteq \bigcap_{M \in F} M$, i. e. $F \subseteq E$. On the other hand, since $\text{Spec}(A) - Y = U(F)$, $\text{Spec}(A) = U(F) \cup V(F)$ and $U(F) \cap V(F) = \emptyset$ this implies that $Y = V(F)$. Thus $M \in V(F) = Y$ iff $F \subseteq M$, i. e. $M \in Y \cap \text{Max}(A) = T$ iff $F \subseteq M$. Therefore $E = \bigcap_{M \in T} W_M \subseteq W_M$ for each $M \in T$, that is for each $M \supseteq F$. Hence $E \subseteq \bigcap_{F \subseteq M} W_M = F$. \square

Corollary 4.5. There is a one to one correspondence between pure filters and subsets $T \subseteq \text{Max}(A)$ where $\bigcup_{M \in T} \widehat{W}_M$ is closed in $\text{Spec}(A)$.

In the next theorem we prove a good relation between closed subsets of $\text{Max}(A)$ and closed subsets of $\text{Spec}(A)$.

Theorem 4.6. Let T be a subset of $\text{Max}(A)$. Then T is closed in $\text{Max}(A)$ iff $\bigcup_{M \in T} \widehat{W}_M$ is closed in $\text{Spec}(A)$.

Proof. Let T be closed in $\text{Max}(A)$. Then $G = \text{Max}(A) - T$ is open in $\text{Max}(A)$. By Corollary 4.3, there is a stable open subset $U(I)$ such that $U(I) \cap \text{Max}(A) = G$. We claim that $\text{Spec}(A) - Y = U(I)$. Since $U(I)$ is stable, by Theorem 4.2, we have $U(I) = \text{Spec}(A) - \bigcup_{I \subseteq M} \widehat{W}_M$. It is enough to show that $M \in T$ iff $I \subseteq M$ (since $Y = \bigcup_{I \subseteq M} \widehat{W}_M$ iff $\bigcup_{I \subseteq M} \widehat{W}_M = \bigcup_{M \in T} \widehat{W}_M$). Now let $I \subseteq M$. Then $M \notin U(I)$, hence $M \notin G$, i. e. $M \in T$. Conversely, let $M \in T$. Since $M \subseteq M$ and $M \in \text{Max}(A) \subseteq \text{Spec}(A)$, $M \in \widehat{W}_M \subseteq \bigcup_{M \in T} \widehat{W}_M = Y$. Then $M \in Y$. It is enough to show that $M \in V(I)$. Suppose that $M \notin V(I)$, i. e. $M \in U(I)$. Then $M \in G$. Thus $M \in \text{Max}(A) - T$ and hence $M \notin T$ which is a contradiction.

Let $Y = \bigcup_{M \in T} \widehat{W}_M$ be closed in $\text{Spec}(A)$. Then $Y \cap \text{Max}(A)$ is closed in $\text{Max}(A)$. But $Y \cap \text{Max}(A) = T$. Therefore T is closed in $\text{Max}(A)$. \square

Theorem 4.7. Let P be a prime filter and T a closed subset of $\text{Max}(A)$. If $\bigcap_{M \in T} W_M \subseteq P$ then $W_M \subseteq P$ for some $M \in T$.

Proof. Suppose that $F = \bigcap_{M \in T} W_M$. Then we have $U(F) = \text{Spec}(A) - \bigcup_{M \in T} \widehat{W}_M$. Let $F \subseteq P$ but $W_M \not\subseteq P$ for all $M \in T$, i.e. $P \notin V(W_M)$. In other words, $P \notin \widehat{W}_M$ for all $M \in T$. Thus $P \notin \bigcup_{M \in T} \widehat{W}_M$, i.e. $P \in U(F)$ and hence $F \not\subseteq P$ which is a contradiction.

The converse of the above Theorem is also true.

Theorem 4.8. Let $T \subseteq \text{Max}(A)$, $F = \bigcap_{M \in T} W_M$ and suppose that each prime filter P , $F \subseteq P$ implies that $W_M \subseteq P$ for some $M \in T$. Then F is a pure filter and T is a closed set.

Proof. Let U be an open set in $\text{Spec}(A)$. We must show that $U(F)$ is stable. Let $P \in U(F)$, $Q \in \text{Spec}(A)$ and $P \subseteq Q$. Suppose that $Q \notin U(F)$, i.e. $F \not\subseteq Q$. Then $\bigcap_{M \in T} W_M \not\subseteq Q$. Thus by Theorem 4.7, $W_M \not\subseteq Q$ for some $M \in T$, i.e. $Q \in V(W_M)$. Therefore $Q \in \widehat{W}_M$ and $Q \subseteq M$. But $P \subseteq Q \subseteq M$ implies that $P \in \widehat{W}_M$ and hence $P \in V(W_M)$. Therefore $W_M \subseteq P$. On the other hand $F \subseteq W_M \subseteq P$. Thus we have $F \subseteq P$. This means that $P \notin U(F)$ which is a contradiction. Therefore we have $Q \in U(F)$. \square

Corollary 4.6. From Theorems 4.7 and 4.8, pure filters correspond to close subsets of $\text{Max}(A)$.

In the next proposition we prove that a prime filter having a certain condition is a pure filter and vice versa.

Proposition 4.1. Let P be a prime filter of A . Then P is pure iff \widehat{W}_{M_P} is a chain and $P = W_{M_P}$.

Proof. Suppose that P is a pure filter. Then $U(P)$ is stable and by Theorem 4.2, $U(P) = \text{Spec}(A) - \bigcup_{P \subseteq M} \widehat{W}_M$. Since $\text{Max}(A) \cap V(P) = M_P$, $U(P) = \text{Spec}(A) - \widehat{W}_{M_P}$. We show that for each $Q \in \widehat{W}_{M_P}$, $P \subseteq Q$. Let $Q \in \widehat{W}_{M_P}$. Then $Q \subseteq M_P$. Since $U(P)$ is stable and $P \subseteq M_P$ by Lemma 4.1, $P \subseteq Q$, i.e. \widehat{W}_{M_P} is a chain and P is a minimal prime filter of A . But $W_{M_P} = \bigcap_{Q \in \widehat{W}_{M_P}} Q = \bigcap_{Q \subseteq M_P} Q = P$ (since $P \subseteq M_P$ and P is minimal prime). Therefore $W_{M_P} = P$.

Now let \widehat{W}_{M_P} be a chain and $P = W_{M_P}$. Thus $U(P) = U(W_{M_P}) = \text{Spec}(A) - V(W_{M_P}) = \text{Spec}(A) - \widehat{W}_{M_P}$ and by Remark 4.1, $U(P)$ is stable and hence P is a pure filter. \square

Now based on the above we get an equivalent statement for a BL-algebra to be hyperarchimedean, that is

Corollary 4.7. A BL-algebra A is hyperarchimedean iff every maximal filter of A is pure.

Proof. Let A be hyperarchimedean. Then by Proposition 2.6, $\text{Spec}(A) = \text{Max}(A)$. Suppose that N is a maximal filter of A . Thus $M_N = N$ and $\widehat{W}_{M_N} = \widehat{W}_N = N$. Since $W_{M_N} = W_N = \bigcap_{\{P \in \text{Spec}(A), P \in \widehat{W}_N\}} P = N$, by Proposition 4.1, we get that N is a pure filter. Conversely, we know that $\text{Max}(A) \subseteq \text{Spec}(A)$. Let $P \in \text{Spec}(A)$ and $M \in \text{Max}(A)$ such that $P \subseteq M$ but $M \not\subseteq P$. This means that $P \in U(M)$. Since $P \subseteq M$ and every maximal filter is pure, $U(M)$ is stable and hence $M \in U(M)$ which is a contradiction. Thus $M \subseteq P$ and A is hyperarchimedean. \square

Proposition 4.2. Let F be a filter of A such that $U(F)$ is stable and $G = U(F) \cap \text{Max}(A)$. Then $U(F)$ is minimal among all $U(E)$ such that $G = U(E) \cap \text{Max}(A)$.

Proof. Suppose that F and E are filters of A such that $U(F)$ is stable and $G = U(E) \cap \text{Max}(A)$. Let $P \in U(F)$ and M_P be the unique maximal filter over P , i.e. $P \subseteq M_P$. By stability of $U(F)$ we have $M_P \in U(F)$. Thus $M_P \in G$ and $M_P \in U(E)$, i.e. $E \not\subseteq M_P$. Hence $E \not\subseteq P$ and $P \in U(E)$. Therefore $U(F) \subseteq U(E)$. \square

Proposition 4.3. Let S be a nonempty \vee -closed subset of A and F be a filter of A such that $F \cap S = \emptyset$. Then there exists a minimal prime filter Q of A such that $F \subseteq Q$ and $Q \cap S = \emptyset$.

Proof. Let F be a proper filter of A . Consider $T = \{P \in \text{Spec}(A) \mid F \subseteq P, F \cap S = \emptyset\}$. By Proposition 2.2, T is nonempty and by Zorn's Lemma, T has a minimal element. \square

Corollary 4.8. Let S be a nonempty \vee -closed subset of A such that $1 \notin S$ then there exists a minimal prime filter Q such that $Q \cap S = \emptyset$.

Proof. Take $F = \{1\}$ and apply Proposition 4.3. \square

We recall that if F is a filter of A and $x \in F$ then $x^\perp = \{y \in A \mid x \vee y = 1\}$ is a filter of A [9].

Theorem 4.9. Let F be a filter of A such that $U(F)$ is stable and let $x \in F$. Then we have $x^\perp \vee F = A$.

Proof. Let $U(F)$ be stable but $x^\perp \vee F \neq A$ for some $x \in F$, i.e. $x^\perp \vee F \subset A$. In other words, $x^\perp \vee F$ is a proper filter of A . Then by Proposition 2.2, there exists a maximal filter M of A such that $x^\perp \vee F \subseteq M$. Then we have $x^\perp \subseteq M$. We define $T = \{x \vee y \mid y \notin M\}$. Since $x = x \vee 0$, $x \in T$ and T is nonempty. Now let $x \vee y$ and $x \vee z$ be two elements of T . Then we have $(x \vee y) \vee (x \vee z) = x \vee (y \vee z) \in T$. Since, $M \in \text{Max}(A) \subseteq \text{Spec}(A)$ then M is a prime filter and $y \notin M$ and $z \notin M$. Thus

we have $y \vee z \notin M$. Now we claim that $1 \notin T$ (otherwise, if $1 \in T$, $x \vee y = 1$ for $y \notin M$, i. e. $y \in x^\perp$ and since $x^\perp \subseteq M$, $y \in M$ which is a contradiction). Therefore by Corollary 4.8, there exists a minimal prime filter Q such that $Q \cap T = \emptyset$. Then $Q \subseteq M$. On the other hand, since $x = x \vee 0$, we have $F \not\subseteq Q$ which means $Q \in U(F)$. Since $Q \in U(F)$ and $Q \subseteq M$ by stability of $U(F)$, we conclude that $M \in U(F)$, i. e. $F \not\subseteq M$. This is a contradiction since $F \subseteq x^\perp \vee F \subseteq M$. \square

The converse of Theorem 4.9 is true, that is

Theorem 4.10. Let F be a filter of A such that for each $x \in A$, $x^\perp \vee F = A$. Then $U(F)$ is stable and hence F is a pure filter.

Proof. Let $P \in U(F)$, $P \subseteq Q$ and $Q \in \text{Spec}(A)$. We must show that $Q \in U(F)$. Assume on the contrary that $Q \notin U(F)$ which implies $F \subseteq Q$. Choose $J \in \text{Min}(A)$ such that $J \subseteq P$. Then $F \not\subseteq J$. In fact if $F \subseteq J \subseteq P$, then $F \subseteq P$ and hence $P \notin U(F)$ which is a contradiction. Hence $F \not\subseteq J$. Thus there exists $x \in F - J$ such that $x^\perp \subseteq J$ (otherwise, if for each $x \in F - J$, $x^\perp \not\subseteq J$, i. e. there exists $t \in x^\perp$, $t \notin J$, we have $t \vee x = 1 \in J$ but $x \notin J$ and $t \notin J$ which is impossible). But $J \subseteq P \subseteq Q$. Then $x^\perp \subseteq Q$. Since $F \subseteq Q$, $\langle x^\perp \cup F \rangle \subseteq Q$, i. e. $x^\perp \vee F \subseteq Q$. Thus $A = Q$. Hence by this contradiction, we conclude that $Q \in U(F)$. \square

Corollary 4.9. Let F be a filter of A . Then $U(F)$ is stable iff $U(F) = \bigcup_{x \in F} V(x^\perp)$.

Proof. Let $U(F) = \bigcup_{x \in F} V(x^\perp)$, $P \in U(F)$, $Q \in \text{Spec}(A)$ and $P \subseteq Q$. Then $F \not\subseteq P$. Therefore there exists $x_1 \in F$ such that $x_1^\perp \subseteq P \subseteq Q$ and hence $P \in V(x_1^\perp)$, that is $x_1^\perp \subseteq Q$. This implies that $Q \in V(x_1^\perp) \subseteq \bigcup_{x_1 \in F} V(x_1^\perp) = U(F)$. Thus $U(F)$ is stable.

Now let $P \in U(F)$, i. e. $F \not\subseteq P$. Therefore, there exists an element $x_1 \in F - P$ such that $x_1^\perp \subseteq P$, that is $P \in V(x_1^\perp) \subseteq \bigcup_{x_1 \in F} V(x_1^\perp)$.

Conversely let $P \in \bigcup_{x_1 \in F} V(x_1^\perp)$. Then there is an element $y \in F$ such that $P \in V(y^\perp)$, i. e. $y^\perp \subseteq P$. We see that $F \not\subseteq P$. In fact if $F \subseteq P$, since $y^\perp \subseteq P$, we conclude that $y^\perp \vee F \subseteq P$. By the stability of $U(F)$ and Theorem 4.9, we have $A \subseteq P$ which is a contradiction and hence $P \in U(F)$. \square

Proposition 4.4. Let F be a filter of A such that $U(F)$ is stable and $P_1, P_2 \in \text{Min}(A)$, $M \in \text{Max}(A)$ such that $P_1, P_2 \subseteq M$. Then we have $F \subseteq P_1$ iff $F \subseteq P_2$.

Proof. Let P_1 and P_2 be two minimal prime filters contained in a some maximal filter M . Suppose that $F \subseteq P_1$ but $F \not\subseteq P_2$. Since $U(F)$ is stable, $M \in U(F)$ i. e. $F \not\subseteq M$, which is a contradiction with $F \subseteq P_1 \subseteq M$. \square

Corollary 4.10. Let F be a pure filter of A and $M \in \text{Max}(A)$. Then for all $P \in \widehat{W}_M$ either $F \subseteq P$ or $F \not\subseteq P$.

Proof. An immediate consequence of Proposition 4.4. \square

We recall that $\sigma(F) = \{a \in A \mid x \wedge y = 0 \text{ for some } x \in F \text{ and } y \in a^\perp\}$ where F is a filter of $L(A)$ [9].

Proposition 4.5. (Leustean [9]) Let F be a filter of $L(A)$. Then $\sigma(F)$ is a filter of A and $\sigma(F) \subseteq F$.

Based on Proposition 4.5, since any filter of A is a filter of $L(A)$, $\sigma(F)$ is a filter of A for every filter F of A .

Corollary 4.11. Let F be a filter of A such that $\sigma(F) = F$. Then $U(F)$ is stable and F is a pure filter.

Proof. By Corollary 4.9, It is enough to show that $U(F) = \bigcup_{x \in F} V(x^\perp)$. Let $P \in U(F)$. Then $F \not\subseteq P$, i.e. there exists $x \in F - P$. Therefore $x^\perp \subseteq P$. Thus $P \in V(x^\perp)$ for some $x \in F$ and hence $P \in \bigcup_{x \in F} V(x^\perp)$. Conversely, we must show that $\bigcup_{x \in F} V(x^\perp) \subseteq U(F)$. Let $P \notin U(F)$ but $P \in V(x^\perp)$ for some $x \in F$, i.e. $x^\perp \subseteq P$ for some $x \in F$. Since $\sigma(F) = F$, $x \in \sigma(F)$. Thus there exists $y \in x^\perp$ and $z \in F$ such that $y \wedge z = 0$. Since $x^\perp \subseteq P$ and $P \notin U(F)$, i.e. $F \not\subseteq P$, we conclude that $y \in P$ and $z \in P$. Therefore $y \odot z \in P$. But $y \odot z \leq y \wedge z = 0$. Hence $z \odot y = 0$, i.e. $0 \in P$ which is a contradiction. \square

It is easy to see that $\sigma(\{1\}) = \{1\}$ and $\sigma(A) = A$. Then A and $\{1\}$ are pure filters of A . Also we know that if A is a nontrivial BL-algebra, then $\text{Rad}(A)$ i.e. the intersection of all maximal filters of A is a proper filter. Now let A be a semisimple BL-algebra, that is $\text{Rad}(A) = \{1\}$ then $\text{Rad}(A)$ is a pure filter.

Proposition 4.6. (Leustean [9]) For each proper filter F of a local BL-algebra A , $\sigma(F) = \{1\}$.

Corollary 4.12. Let A be a local BL-algebra. Then $\{1\}$ is the unique proper pure filter of A .

Proof. It is an immediate consequence of Proposition 4.6 and Corollary 4.11 since $\{1\}$ is the only filter F satisfying $\sigma(F) = \{1\}$. We conclude that in each local BL-algebra A , the only pure filters are A and $\{1\}$. \square

Corollary 4.13. Let A be a BL-chain. Then A and $\{1\}$ are the only pure filters.

Proof. Since each BL-chain is a local BL-algebra ([11]), the Corollary follows from Corollary 4.11 \square

Corollary 4.14. In each local BL-algebra A , the stable topology is trivial.

Proof. Let $T = U(F)$ for some filter F of A be a stable open set of $\text{Spec}(A)$. Then by Corollary 4.12, we have $T = U(\{1\})$ or $T = U(A)$. Therefore $T = \emptyset$ or $T = \text{Spec}(A)$. \square

Corollary 4.15. Let A and B be BL-algebras, $h : A \longrightarrow B$ a surjective BL-morphism and F be a proper filter of A such that $\sigma(F) = F$. Then we have $\sigma(h(F)) = h(\sigma(F)) = h(F)$. In other words, $U(h(F))$ is stable and hence $h(F)$ is a pure filter of B .

Proof. Suppose that F is a proper filter of A . By Proposition 2.1, since h is surjective, $h(F)$ is a proper filter of B . Now by Proposition 4.5, $\sigma(h(F)) \subseteq h(F)$. But $\sigma(F) = F$. Therefore, $\sigma(h(F)) \subseteq h(\sigma(F))$. Conversely, let $x \in h(\sigma(F))$. Then $x = h(k)$ for some $k \in \sigma(F)$, i. e. $y \wedge z = 0$ for some $y \in k^\perp$ and some $z \in F$. Take $l = h(y)$ and $s = h(z)$. It is easy to see that $s \in h(F)$ and $l \in h(k)^\perp$. Since $y \in k^\perp$, $y \vee k = 1$ and $l \vee s = h(y) \vee h(k) = h(y \vee k) = h(1) = 1$. On the other hand, $l \wedge s = h(y) \wedge h(z) = h(y \wedge z) = h(0) = 0$. This means that $x = h(k) \in \sigma(h(F))$. \square

We like to give another proof for Corollary 4.15 from Corollary 4.9, i. e. it is enough to show that $U(h(F)) = \bigcup_{l \in h(F)} V(l^\perp)$.

Let P be a prime filter of B , $P \in U(h(F))$. Then $h(F) \not\subseteq P$, i. e. $h(x) \notin P$ for some $l = h(x) \in h(F)$. Thus $l^\perp = h(x)^\perp \subseteq P$, i. e. $P \in V(l^\perp)$ and hence $P \in \bigcup_{l \in h(F)} V(l^\perp)$. Conversely, let $P \in \text{Spec}(B)$ and $P \notin U(h(F))$ but $P \in \bigcup_{l \in h(F)} V(l^\perp)$, that is $h(F) \subseteq P$ and $l^\perp \subseteq P$ for some $l \in h(F)$. Since $l \in h(F)$, $l = h(x)$ for some $x \in F$. By Proposition 2.1, $h^{-1}(P) \in \text{Spec}(A)$. Now we claim that $x^\perp \subseteq h^{-1}(P)$. Suppose that $v \in x^\perp$. Then $v \vee x = 1$ and hence $h(v \vee x) = h(v) \vee h(x) = 1$. Therefore, $h(v) \in h(x)^\perp$ and by above $h(v) \in P$, i. e. $v \in h^{-1}(P)$. Thus $h^{-1}(P) \in V(x^\perp) \subseteq \bigcup_{m \in F} V(m^\perp)$. Since $\sigma(F) = F$, by Corollary 4.11, $U(F)$ is stable and by Corollary 4.9 we conclude that $\bigcup_{m \in F} V(m^\perp) = U(F)$. Thus, $h^{-1}(P) \in U(F)$ and $F \not\subseteq h^{-1}(P)$. This means that $h(F) \not\subseteq P$ and $P \in U(h(F))$, which contradicts our assumption.

ACKNOWLEDGEMENT

We thank the two referees for their valuable remarks and suggestions. Also, the editors deserve special thanks for helpful comments regarding the improvement of the paper.

(Received January 29, 2008.)

REFERENCES

-
- [1] L. P. Belluce, A. Di Nola, and S. Sessa: The prime spectrum of an MV-algebra. *Math. Logic Quart.* 40 (1994), 331–346.
 - [2] L. P. Belluce and S. Sessa: The stable topology for MV-algebras. *Quaestiones Math.* 23 (2000), 3, 269–277.
 - [3] D. Busneage and D. Piciu: On the lattice of deductive system of a BL-algebra. *Central European Journal of Mathematics* 2 (2003), 221–237.
 - [4] A. Di Nola, G. Georgescu, and A. Iorgulescu: Pseudo-BL-algebra, Part II. *Multiple Valued Logic* 8 (2002), 717–750.
 - [5] G. Georgescu and L. Leustean: Semilocal and maximal BL-algebras. Preprint.

- [6] P. Hájek: *Metamathematics of Fuzzy Logic*, Trends in Logic. (Studia Logica Library 4.) Kluwer Academic Publishers, Dordrecht 1998.
- [7] P. T. Johnstone: *Stone Spaces*. (Cambridge Studies in Advanced Mathematics.) Cambridge University Press, Cambridge 1982.
- [8] L. Leustean: The prime and maximal spectra and the reticulation of BL-algebras. *Central European Journal of Mathematics* 1 (2003), 382–397.
- [9] L. Leustean: *Representations of Many-Valued Algebras*. PhD. Thesis, University of Bucharest 2004.
- [10] E. Turunen: *Mathematics Behind Fuzzy Logic*. Advances in Soft Computing. Physica-Verlag, Heidelberg 1999.
- [11] E. Turunen and S. Sessa: Local BL-algebras. *Multi-Valued Log.* 6 (2001), 1–2, 229–249.

*Esfandiar Eslami, Department of Mathematics & Fuzzy Systems and Applications Center of Excellence, Shahid Bahonar University of Kerman, Kerman. Iran.
e-mail: eeslami@mail.uk.ac.ir*

*Farhad Kh. Haghani, Department of Mathematics, Shahid Bahonar University of Kerman, Kerman, and Islamic Azad University, Shahrekord Branch, Shahrekord. Iran.
e-mail: haghani1351@yahoo.com*