

# PASSIVITY ANALYSIS OF UNCERTAIN STOCHASTIC NEURAL NETWORK WITH LEAKAGE AND DISTRIBUTED DELAYS UNDER IMPULSIVE PERTURBATIONS

SENTHIL RAJ, RAJA RAMACHANDRAN, SAMIDURAI RAJENDIRAN, JINDE CAO AND XIAODI LI

In this paper, the problem of passivity analysis for a class of uncertain stochastic neural networks with mixed delays and impulsive control is investigated. The mixed delays include constant delay in the leakage term, discrete and distributed delays. The discrete delays are assumed to be time-varying and belong to a given interval, which means that the lower and upper bounds of interval time-varying delays are available. By using Lyapunov stability theory, stochastic analysis, linear matrix inequality techniques and introducing some free-weighting matrices, several novel sufficient conditions are derived to guarantee the passivity of the suggested system in the sense of mean square under two cases: with known or unknown parameters. It is believed that these results are significant and useful for the design and applications of impulsive stochastic neural networks. Finally, two numerical examples are provided to show the effectiveness of the theoretical results.

*Keywords:* distributed delays, leakage delay, passivity impulses, stochastic disturbances

*Classification:* 34Dxx

## 1. INTRODUCTION

Neural networks can imitate the human brain, and they have been used for a wide variety of applications, for example, target tracking, machine learning, system identification and so on [7, 25, 30]. Moreover, as we know, the applications of neural networks heavily depend on their dynamic behaviors. On the other hand, time delays are always unavoidably encountered in the implementation of neural networks due to the finite switching speed of neurons and amplifiers. Therefore, increasing attention has been paid to the problem of neural networks with various delays have been reported in [3, 4, 18, 12, 33]. Some criteria have been proposed to ensure the fixed-time synchronization for memristive neural network [3]. The  $H_\infty$  filtering problem for delayed discrete-time switched neural networks has been considered in [4].

On the other hand, due to modeling and measurement errors, neural network is often disturbed by stochastic factors and the parameter uncertainties. Because in real nervous

system, synaptic transmission is a noisy process and the connection weights of the neurons depend on certain resistance and capacitance values which include uncertainties. Hence, their presence must be considered in realistic dynamics and some results related to this problem have recently published in [8, 11, 14, 29].

The concept of passivity has played an important role in the analysis of the stability of dynamical systems, nonlinear control, and other research areas. The essence of the passivity theory is that the passive properties of a system can keep the system internal stability. So it gives a way to study nonlinear system only by means of the general characteristics of the input-output dynamics. Recently, passivity analysis problem for various neural networks was widely investigated in the literature [5, 15, 20, 34, 37]. Cao and Li have investigated the stability of memristive neural networks with leakage delay, and the uncertainties was also considered [15]. In [5], Chen et al. presented, both delay-independent and delay-dependent passivity conditions for stochastic neural network in the sense of mean square. Very recently Raja et al.[28], discussed the passivity analysis for stochastic BAM neural networks with time-varying structured uncertainties.

As we know, time delay in the stabilizing negative feedback term has a tendency to destabilize a system [6, 16]. Like the traditional time delays, the leakage delays also have a great impact on the dynamics of neural networks and many works appeared in the literature, see [13, 17, 22, 31]. Based on this work, [31] pay attention to the passivity analysis of uncertain neural networks. In [22], authors studied the equilibrium point of two classes fuzzy neural networks with delays in leakage terms; By use of the topological degree theory, delay-dependent stability conditions of neural networks of neutral type with time delay in the leakage term was proposed in [13]. Therefore, it is considerable to investigate the passivity analysis of neural networks with time delays in the leakage term and very little existing works appeared in the literature [1, 31, 36]. The passivity properties of uncertain neural networks with leakage delay and time-varying delay has been studied in [31]. In [1], authors investigated the problem for passivity analysis of neutral type neural networks with Markovian jumping parameters and time delays in the leakage term. Unfortunately, in these works, authors neglected the effects of stochastic disturbances, which has also an important effect on the passivity analysis of neural networks. But in [36], Zhao et al. presented the passivity problem for stochastic neural networks with time-varying delays and leakage delay using Lyapunov functional and free-weighting matrix method.

In addition, many physical systems undergo unexpected changes at certain moments due to instantaneous perturbations, which leads to impulsive effects [19, 21]. It is worth pointing out that neural networks are often subject to impulsive perturbations that in turn affect dynamical behaviors of the system. It frequently occurs in fields such as economics, mechanics, electronics, telecommunications, medicine and biology, etc. Therefore, it is necessary to consider the impulsive effects to the passivity problem of stochastic neural networks to reflect more realistic dynamics and several interesting results have been reported for continuous-time and discrete-time neural networks [26, 27, 19, 35]. More recently, in [27], Raja et al. derived the dissipativity results for a class of uncertain discrete-time stochastic neural networks with impulsive parameters. However, to the best of our knowledge, the passivity analysis problem for stochastic neural network with the effects of leakage delays and impulsive perturbations has not

been investigated in the previous literature. This motivates our present study.

In this paper, we deal with the passivity problem of impulsive stochastic neural networks with leakage, discrete and distributed delays. Then by using Lyapunov functional, Free weighting matrix method and stochastic analysis techniques, some sufficient conditions that dependent on the delays for passivity are obtained in terms of LMIs, which can be readily verified by using standard numerical software. Finally, two numerical examples are given to illustrate the effectiveness of the proposed criteria.

**Notations.** Throughout this paper,  $\mathbb{R}^n$  and  $\mathbb{R}^{n \times m}$  denote the  $n$ -dimensional Euclidean space and the set of all  $n \times m$  real matrices, respectively. The notation  $X \geq 0$  (respectively,  $X > 0$ ), where  $X$  is symmetric matrices, means that  $X$  is positive semi-definite (respectively, positive definite). Let  $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \geq 0}, \mathcal{P})$  be the complete probability space with a filtration  $\{\mathcal{F}_t\}_{t \geq 0}$  satisfying the usual conditions (i.e. the filtration contains all  $\mathcal{P}$ - null sets and is right continuous).  $\omega(t)$  be a scalar Brownian motion defined on the probability space.  $\mathbb{E}[\cdot]$  is the mathematical expectation operator with respect to the given probability measure  $\mathcal{P}$ .

## 2. PROBLEM FORMULATION

Consider the following uncertain stochastic neural networks with both discrete and distributed time-varying delays described by

$$\begin{aligned} dx(t) &= \left[ -Ax(t - \delta) + Bg(x(t)) + Cg(x(t - \tau(t))) + D \int_{t-d(t)}^t g(x(s)) ds + u(t) \right] dt \\ &\quad + \sigma(t, x(t), x(t - \tau(t)), x(t - d(t))) d\omega(t), \quad t \neq t_k \\ y(t) &= g(x(t)), \\ \Delta x(t_k) &= -I_k \left\{ x(t_k^-) - A \int_{t_k - \delta}^{t_k} x(s) ds \right\}, \quad t = t_k, \quad k \in \mathbb{Z}_+, \end{aligned} \quad (1)$$

where  $x(t) = [x_1(t), x_2(t), \dots, x_n(t)]^T \in \mathbb{R}^n$  is the state vector associated with the neurons,  $g(x(t)) = [g_1(x_1(t)), g_2(x_2(t)), \dots, g_n(x_n(t))]^T$  is the activation function,  $u(t) = [u_1(t), u_2(t), \dots, u_n(t)]^T$  is the input,  $y(t) = [y_1(t), y_2(t), \dots, y_n(t)]$  is the output.  $\sigma \in \mathbb{R}^{n \times q}$  is the diffusion coefficient vector and  $w(t) = (w_1(t), w_1(t), \dots, w_q(t))^T$  is a  $q$ -dimensional Brownian motion defined on a complete probability space

$(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \geq 0}, \mathcal{P})$  be the complete probability space with a filtration  $\{\mathcal{F}_t\}_{t \geq 0}$  satisfying the usual conditions (i.e. the filtration contains all  $\mathcal{P}$ - null sets and is right continuous). The matrix  $A = \text{diag}(a_1, a_2, \dots, a_n)$  is a diagonal matrix with positive entries  $a_i > 0$ .  $B, C, D$  are the interconnection matrices representing the weight coefficients of the neurons.  $I_k \in \mathbb{R}^{n \times n}$ ,  $k \in \mathbb{Z}_+$  denotes the impulsive matrix. The discrete delay  $\tau(t)$  and distribute delay  $d(t)$  satisfies

$$0 \leq \tau_1 \leq \tau(t) \leq \tau_2, \quad \dot{\tau}(t) \leq \mu, \quad 0 < d(t) \leq d \quad (2)$$

where  $\tau_1, \tau_2, d$  and  $\mu$  are constants. The initial condition associated with model (1) is given by

$$x(t) = \phi(t), \quad \forall t \in [-\max\{\delta, \tau_1, \tau_2, d\}, 0].$$

Throughout this paper, it is assumed that the activation functions satisfy the following assumptions:

**(H1)** For any  $j \in 1, 2, \dots, n$ ,  $f_j(0) = 0$  and there exist constants  $F_j^-$  and  $F_j^+$  such that

$$F_j^- \leq \frac{f_j(\alpha_1) - f_j(\alpha_2)}{\alpha_1 - \alpha_2} \leq F_j^+ \quad (3)$$

for all  $\alpha_1 \neq \alpha_2$ .

**(H2)** Assume that  $\sigma : \mathcal{R}^n \times \mathcal{R}^n \times \mathcal{R}^+ \times S \rightarrow \mathcal{R}^n$  is locally Lipschitz continuous and satisfies the linear growth condition [24]. Moreover,  $\sigma$  satisfies

$$\text{trace}[\sigma^T(x_1, x_2, t, i)\sigma(x_1, x_2, t, i)] \leq x_1^T \Sigma_{1i} x_1 + x_2^T \Sigma_{2i} x_2 \quad (4)$$

for all  $x_1, x_2 \in \mathcal{R}^n$  and  $x(t) = i, i \in S$ , where  $\Sigma_{1i}$  and  $\Sigma_{2i}$  are known positive constant matrices with appropriate dimensions.

**(H3)** The impulsive time instant  $t_k$  satisfy  $0 = t_0 < t_1 < \dots < t_k \rightarrow \infty$  and  $\inf_{k \in \mathbb{Z}_+} \{t_k - t_{k-1}\} > 0$ .

**Definition 1.** The stochastic neural networks (1) is said to be stochastically passive from input  $u(t)$  to output  $y(t)$ , if there exists a scalar  $\gamma \geq 0$  such that the following inequality holds:

$$2\mathbb{E} \left[ \int_0^{t_f} y^T(s)u(s) ds \right] \geq -\gamma \mathbb{E} \left[ \int_0^{t_f} u^T(s)u(s) ds \right] \quad (5)$$

for the solution of (1) with  $x(0) = 0$ . We introducing the following lemmas which are useful in the proof of the main results.

**Lemma 2.1.** (Gu [9]) For any positive definite matrix  $M \in \mathbb{R}^{n \times n}$ , scalars  $h_2 > h_1 > 0$ , vector function  $w : [h_1, h_2] \rightarrow \mathbb{R}^n$  such that the integrations concerned are well defined, the following inequality holds:

$$\begin{aligned} -(h_2 - h_1) \int_{t-h_2}^{t-h_1} w^T(s)Mw(s) ds &\leq - \left( \int_{t-h_2}^{t-h_1} w(s) ds \right)^T M \left( \int_{t-h_2}^{t-h_1} w(s) ds \right) \\ -\frac{1}{2}(h_2^2 - h_1^2) \int_{-h_2}^{-h_1} \int_{t+\theta}^t w^T(s)Mw(s) dsd\theta &\leq - \left( \int_{-h_2}^{-h_1} \int_{t+\theta}^t w^T(s) dsd\theta \right) M \\ &\quad \times \left( \int_{-h_2}^{-h_1} \int_{t+\theta}^t w(s) dsd\theta \right) \end{aligned}$$



where

$$\begin{aligned}
\Psi_1 &= (\Psi_{ij})_{20 \times 20} \\
\Psi_{11} &= -P_1 A - A^T P_1 + P_2 + \delta^2 P_3 + P_4 + L_1 + \tau_1^2 Q_1 + (\tau_2 - \tau_1)^2 Q_2 - F_1 L + M_1 \\
&\quad + M_1^T + [\lambda_1 + \tau_1 \lambda_2 + (\tau_2 - \tau_1) \lambda_3] \Sigma_1^T \Sigma_1, \quad \Psi_{12} = M_2^T - N_1 + U_1, \\
\Psi_{13} &= M_3^T - M_1 + N_1, \quad \Psi_{14} = M_4^T - U_1, \quad \Psi_{15} = P_1 B + L_2 + F_2 L + Z_1 B + M_5^T, \\
\Psi_{16} &= P_1 C + Z_1 C + M_6^T, \quad \Psi_{17} = M_7^T, \quad \Psi_{18} = M_8^T, \quad \Psi_{19} = -Z_1 A, \quad \Psi_{110} = A^T P_1 A, \\
\Psi_{112} &= -Z_1, \quad \Psi_{113} = P_1 + Z_1, \quad \Psi_{114} = P_1 D + Z_1 D, \\
\Psi_{22} &= -(1 - \mu) R_1 - F_1 S - N_2 - N_2^T + U_2 + U_2^T + [\lambda_1 + \tau_1 \lambda_2 + (\tau_2 - \tau_1) \lambda_3] \Sigma_2^T \Sigma_2, \\
\Psi_{23} &= -M_2 + N_2 - N_3^T + U_3^T, \quad \Psi_{24} = -N_4^T + U_4^T - U_2, \quad \Psi_{25} = -N_5^T + U_5^T, \\
\Psi_{26} &= -(1 - \mu) R_2 + F_2 S - N_6^T + U_6^T, \quad \Psi_{27} = -N_7^T + U_7^T, \quad \Psi_{28} = -N_8^T + U_8^T, \\
\Psi_{33} &= -L_1 + R_1 + S_1 - M_3 - M_3^T + N_3 + N_3^T, \quad \Psi_{34} = -M_4^T + N_4^T - U_3, \\
\Psi_{35} &= -M_5^T + N_5^T, \quad \Psi_{36} = -M_6^T + N_6^T, \quad \Psi_{37} = -L_2 + R_2 + S_2 - M_7^T + N_7^T, \\
\Psi_{38} &= -M_8^T + N_8^T, \quad \Psi_{44} = -S_1 - U_4 - U_4^T, \quad \Psi_{45} = -U_5^T, \quad \Psi_{46} = -U_6^T, \quad \Psi_{47} = -U_7^T, \\
\Psi_{48} &= -S_2 - U_8^T, \quad \Psi_{55} = d^2 P_5 + L_3 + \tau_1^2 Q_3 + (\tau_2 - \tau_1)^2 Q_4 - L, \quad \Psi_{510} = -B^T P_1 A, \\
\Psi_{512} &= B^T Z_2^T, \quad \Psi_{513} = -I, \quad \Psi_{66} = -(1 - \mu) R_3 - S, \quad \Psi_{610} = -C^T P_1 A, \quad \Psi_{612} = C^T Z_2^T, \\
\Psi_{77} &= -L_3 + R_3 + S_3, \quad \Psi_{88} = -S_3, \quad \Psi_{99} = -P_2, \quad \Psi_{912} = -A^T Z_2^T, \quad \Psi_{1010} = -P_3, \\
\Psi_{1013} &= -A^T P_1, \quad \Psi_{1014} = -A^T P_1 D, \quad \Psi_{1111} = -P_4 + [\lambda_1 + \tau_1 \lambda_2 + (\tau_2 - \tau_1) \lambda_3] \Sigma_3^T \Sigma_3, \\
\Psi_{1212} &= \tau_1 T_1 + (\tau_2 - \tau_1) T_2 + \frac{\tau_1^4}{4} T_3 + \frac{(\tau_2^2 - \tau_1^2)^2}{4} T_4 - Z_2 - Z_2^T, \quad \Psi_{1213} = Z_2, \\
\Psi_{1214} &= Z_2 D, \quad \Psi_{1313} = -\gamma I, \quad \Psi_{1414} = -P_5, \quad \Psi_{1515} = -Q_1, \quad \Psi_{1616} = -Q_2, \\
\Psi_{1717} &= -Q_3, \quad \Psi_{1818} = -Q_4, \quad \Psi_{1919} = -T_3, \quad \Psi_{2020} = -T_4
\end{aligned}$$

$$\begin{aligned}
M^T &= [ M_1^T \ M_2^T \ M_3^T \ M_4^T \ M_5^T \ M_6^T \ M_7^T \ M_8^T \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 ] \\
N^T &= [ N_1^T \ N_2^T \ N_3^T \ N_4^T \ N_5^T \ N_6^T \ N_7^T \ N_8^T \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 ] \\
U^T &= [ U_1^T \ U_2^T \ U_3^T \ U_4^T \ U_5^T \ U_6^T \ U_7^T \ U_8^T \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 \ 0 ]
\end{aligned}$$

*Proof.* For simplicity, we denote

$$g(t) = -Ax(t - \delta) + Bg(x(t)) + Cg(x(t - \tau(t))) + D \int_{t-d(t)}^t g(x(s)) ds + u(t), \quad (11)$$

$$\alpha(t) = \sigma(t, x(t), x(t - \tau(t)), x(t - d(t))), \quad (12)$$

then system (1) can be rewritten as

$$dx(t) = g(t)dt + \alpha(t)dw(t). \quad (13)$$

Choose a Lyapunov functional candidate for the system (1) to be

$$V(x_t) = \sum_{i=1}^9 V_i(x_t) \quad (14)$$

where

$$\begin{aligned} V_1(x_t) &= \left[ x(t) - A \int_{t-\delta}^t x(s) ds \right]^T P_1 \left[ x(t) - A \int_{t-\delta}^t x(s) ds \right] \\ V_2(x_t) &= \int_{t-\delta}^t x^T(s) P_2 x(s) ds + \delta \int_{-\delta}^0 \int_{t+\theta}^t x^T(s) P_3 x(s) ds d\theta \\ V_3(x_t) &= \int_{t-d(t)}^t x^T(s) P_4 x(s) ds + d \int_{-d}^0 \int_{t+\theta}^t g^T(x(s)) P_5 g(x(s)) ds d\theta \\ V_4(x_t) &= \int_{t-\tau_1}^t \varphi^T(s) \mathcal{L} \varphi(s) ds + \int_{t-\tau(t)}^{t-\tau_1} \varphi^T(s) \Re \varphi(s) ds \\ &\quad + \int_{t-\tau_2}^{t-\tau_1} \varphi^T(s) \Im \varphi(s) ds \\ V_5(x_t) &= \tau_1 \int_{-\tau_1}^0 \int_{t+\theta}^t x^T(s) Q_1 x(s) ds d\theta + (\tau_2 - \tau_1) \int_{-\tau_2}^{-\tau_1} \int_{t+\theta}^t x^T(s) Q_2 x(s) ds d\theta \\ V_6(x_t) &= \tau_1 \int_{-\tau_1}^0 \int_{t+\theta}^t g^T(x(s)) Q_3 g(x(s)) ds d\theta \\ &\quad + (\tau_2 - \tau_1) \int_{-\tau_2}^{-\tau_1} \int_{t+\theta}^t g^T(x(s)) Q_4 g(x(s)) ds d\theta \\ V_7(x_t) &= \int_{-\tau_1}^0 \int_{t+\theta}^t g^T(s) T_1 g(s) ds d\theta + \int_{-\tau_2}^{-\tau_1} \int_{t+\theta}^t g^T(s) T_2 g(s) ds d\theta \\ V_8(x_t) &= \frac{\tau_1^2}{2} \int_{-\tau_1}^0 \int_{\theta}^0 \int_{t+\lambda}^t g^T(s) T_3 g(s) ds d\lambda d\theta \\ &\quad + \frac{\tau_2^2 - \tau_1^2}{2} \int_{-\tau_2}^{-\tau_1} \int_{\theta}^0 \int_{t+\lambda}^t g^T(s) T_4 g(s) ds d\lambda d\theta \\ V_9(x_t) &= \int_{-\tau_1}^0 \int_{t+\theta}^t \text{tr}(\alpha^T(s) X_1 \alpha(s)) ds d\theta + \int_{-\tau_2}^{-\tau_1} \int_{t+\theta}^t \text{tr}(\alpha^T(s) X_2 \alpha(s)) ds d\theta \end{aligned}$$

and

$$\varphi(s) = \begin{bmatrix} x(s) \\ g(x(s)) \end{bmatrix}.$$

Then, it can be obtained by *Itô's* formula that

$$dV(x_t, t) = \mathcal{L}V(x_t, t)dt + 2x^T(t)P_1\alpha(t) d\omega(t) \quad (15)$$

where

$$\begin{aligned} \mathcal{L}V_1(x_t, t) &= 2 \left[ x(t) - A \int_{t-\delta}^t x(s) ds \right]^T P_1 \left[ -Ax(t) + Bg(x(t)) + Cg(x(t-\tau(t))) \right. \\ &\quad \left. + D \int_{t-d(t)}^t g(x(s)) ds + u(t) \right] + \text{tr}(\alpha^T(t)P_1\alpha(t)) \end{aligned} \quad (16)$$

$$\mathcal{L}V_2(x_t, t) = x^T(t)[P_2 + \delta^2 P_3]x(t) - x^T(t-\delta)P_2x(t-\delta) - \delta \int_{t-\delta}^t x^T(s)P_3x(s) ds \quad (17)$$

$$\begin{aligned} \mathcal{L}V_3(x_t, t) &= x^T(t)P_4x(t) - x^T(t-d(t))P_4x(t-d(t)) + d^2 g^T(x(t))P_5g(x(t)) \\ &\quad - d \int_{t-d}^t g^T(x(s))P_5g(x(s)) ds \end{aligned} \quad (18)$$

$$\begin{aligned} \mathcal{L}V_4(x_t, t) &= \varphi^T(t)\mathcal{L}\varphi(t) - \varphi^T(t-\tau_1)\mathcal{L}\varphi(t-\tau_1) + \varphi^T(t-\tau_1)\Re\varphi(t-\tau_1) \\ &\quad - (1-\mu)\varphi^T(t-\tau(t))\Re\varphi(t-\tau(t)) + \varphi^T(t-\tau_1)\Im\varphi(t-\tau_1) \\ &\quad - \varphi^T(t-\tau_2)\Im\varphi(t-\tau_2) \\ &\leq \begin{bmatrix} x(t) \\ g(x(t)) \end{bmatrix}^T \begin{bmatrix} L_1 & L_2 \\ L_2^T & L_3 \end{bmatrix} \begin{bmatrix} x(t) \\ g(x(t)) \end{bmatrix} - \begin{bmatrix} x(t-\tau_1) \\ g(x(t-\tau_1)) \end{bmatrix}^T \begin{bmatrix} L_1 & L_2 \\ L_2^T & L_3 \end{bmatrix} \begin{bmatrix} x(t-\tau_1) \\ g(x(t-\tau_1)) \end{bmatrix} \\ &\quad + \begin{bmatrix} x(t-\tau_1) \\ g(x(t-\tau_1)) \end{bmatrix}^T \begin{bmatrix} R_1 & R_2 \\ R_2^T & R_3 \end{bmatrix} \begin{bmatrix} x(t-\tau_1) \\ g(x(t-\tau_1)) \end{bmatrix} \\ &\quad - (1-\mu) \begin{bmatrix} x(t-\tau(t)) \\ g(x(t-\tau(t))) \end{bmatrix}^T \begin{bmatrix} R_1 & R_2 \\ R_2^T & R_3 \end{bmatrix} \begin{bmatrix} x(t-\tau(t)) \\ g(x(t-\tau(t))) \end{bmatrix} \\ &\quad + \begin{bmatrix} x(t-\tau_1) \\ g(x(t-\tau_1)) \end{bmatrix}^T \begin{bmatrix} S_1 & S_2 \\ S_2^T & S_3 \end{bmatrix} \begin{bmatrix} x(t-\tau_1) \\ g(x(t-\tau_1)) \end{bmatrix} \\ &\quad - \begin{bmatrix} x(t-\tau_2) \\ g(x(t-\tau_2)) \end{bmatrix}^T \begin{bmatrix} S_1 & S_2 \\ S_2^T & S_3 \end{bmatrix} \begin{bmatrix} x(t-\tau_2) \\ g(x(t-\tau_2)) \end{bmatrix} \end{aligned} \quad (19)$$

$$\begin{aligned} \mathcal{L}V_5(x_t, t) &= x^T(t)[\tau_1^2 Q_1 + (\tau_2 - \tau_1)^2 Q_2]x(t) - \tau_1 \int_{t-\tau_1}^t x^T(s)Q_1x(s) ds \\ &\quad - (\tau_2 - \tau_1) \int_{t-\tau_2}^{t-\tau_1} x^T(s)Q_2x(s) ds \end{aligned} \quad (20)$$

$$\begin{aligned} \mathcal{L}V_6(x_t, t) &= g^T(x(t))[\tau_1^2 Q_3 + (\tau_2 - \tau_1)^2 Q_4]g(x(t)) - \tau_1 \int_{t-\tau_1}^t g^T(x(s))Q_3g(x(s)) ds \\ &\quad - (\tau_2 - \tau_1) \int_{t-\tau_2}^{t-\tau_1} g^T(x(s))Q_4g(x(s)) ds \end{aligned} \quad (21)$$

$$\begin{aligned} \mathcal{L}V_7(x_t, t) &= g^T(t)[\tau_1 T_1 + (\tau_2 - \tau_1)T_2]g(t) - \int_{t-\tau_1}^t g^T(s)T_1g(s) ds - \int_{t-\tau_2}^{t-\tau_1} g^T(s)T_2g(s) ds \\ &= g^T(t)[\tau_1 T_1 + (\tau_2 - \tau_1)T_2]g(t) - \int_{t-\tau_1}^t g^T(s)T_1g(s) ds \\ &\quad - \int_{t-\tau(t)}^{t-\tau_1} g^T(s)T_2g(s) ds - \int_{t-\tau_2}^{t-\tau(t)} g^T(s)T_2g(s) ds \end{aligned} \quad (22)$$



$$\begin{aligned} \mathcal{L}V_8(x_t, t) &= \frac{\tau_1^4}{4} g^T(t) T_3 g(t) - \frac{\tau_1^2}{2} \int_{-\tau_1}^0 \int_{t+\theta}^t g^T(s) T_3 g(s) \, ds d\theta + \frac{(\tau_2^2 - \tau_1^2)^2}{4} g^T(t) T_4 g(t) \\ &\quad - \left( \frac{\tau_2^2 - \tau_1^2}{2} \right) \int_{-\tau_2}^{-\tau_1} \int_{t+\theta}^t g^T(s) T_4 g(s) \, ds d\theta \end{aligned} \quad (23)$$

$$\begin{aligned} \mathcal{L}V_9(x_t, t) &= \tau_1 \operatorname{tr}(\alpha^T(t) X_1 \alpha(t)) + (\tau_2 - \tau_1) \operatorname{tr}(\alpha^T(t) X_2 \alpha(t)) \\ &\quad - \int_{t-\tau_1}^t \operatorname{tr}(\alpha^T(s) X_1 \alpha(s)) \, ds - \int_{t-\tau_2}^{t-\tau_1} \operatorname{tr}(\alpha^T(s) X_2 \alpha(s)) \, ds \\ &\leq \tau_1 \lambda_2 \operatorname{tr}(\alpha^T(t) X_1 \alpha(t)) + (\tau_2 - \tau_1) \lambda_3 \operatorname{tr}(\alpha^T(t) X_2 \alpha(t)) \\ &\quad - \int_{t-\tau_1}^t \operatorname{tr}(\alpha^T(s) X_1 \alpha(s)) \, ds \\ &\quad - \int_{t-\tau(t)}^{t-\tau_1} \operatorname{tr}(\alpha^T(s) X_2 \alpha(s)) \, ds - \int_{t-\tau_2}^{t-\tau(t)} \operatorname{tr}(\alpha^T(s) X_2 \alpha(s)) \, ds. \end{aligned} \quad (24)$$

From Lemma 2.1, one can obtain

$$- \delta \int_{t-\delta}^t x^T(s) P_3 x(s) \, ds \leq - \left( \int_{t-\delta}^t x(s) \, ds \right)^T P_3 \left( \int_{t-\delta}^t x(s) \, ds \right) \quad (25)$$

$$- d \int_{t-d}^t g^T(x(s)) P_5 g(x(s)) \, ds \leq - \left( \int_{t-d(t)}^t g(x(s)) \, ds \right)^T P_5 \left( \int_{t-d(t)}^t g(x(s)) \, ds \right) \quad (26)$$

$$- \tau_1 \int_{t-\tau_1}^t x^T(s) Q_1 x(s) \, ds \leq - \left( \int_{t-\tau_1}^t x(s) \, ds \right)^T Q_1 \left( \int_{t-\tau_1}^t x(s) \, ds \right) \quad (27)$$

$$- (\tau_2 - \tau_1) \int_{t-\tau_2}^{t-\tau_1} x^T(s) Q_2 x(s) \, ds \leq - \left( \int_{t-\tau_2}^{t-\tau_1} x(s) \, ds \right)^T Q_2 \left( \int_{t-\tau_2}^{t-\tau_1} x(s) \, ds \right) \quad (28)$$

$$- \tau_1 \int_{t-\tau_1}^t g^T(x(s)) Q_3 g(x(s)) \, ds \leq - \left( \int_{t-\tau_1}^t g(x(s)) \, ds \right)^T Q_3 \left( \int_{t-\tau_1}^t g(x(s)) \, ds \right) \quad (29)$$

$$\begin{aligned} - (\tau_2 - \tau_1) \int_{t-\tau_2}^{t-\tau_1} g^T(x(s)) Q_4 g(x(s)) \, ds &\leq - \left( \int_{t-\tau_2}^{t-\tau_1} g(x(s)) \, ds \right)^T Q_4 \\ &\quad \times \left( \int_{t-\tau_2}^{t-\tau_1} g(x(s)) \, ds \right) \end{aligned} \quad (30)$$

$$\begin{aligned} - \frac{\tau_1^2}{2} \int_{-\tau_1}^0 \int_{t+\theta}^t g^T(s) T_3 g(s) \, ds d\theta &\leq - \left( \int_{-\tau_1}^0 \int_{t+\theta}^t g(s) \, ds d\theta \right)^T \\ &\quad \times T_3 \left( \int_{-\tau_1}^0 \int_{t+\theta}^t g(s) \, ds d\theta \right) \end{aligned} \quad (31)$$

$$\begin{aligned} - \left( \frac{\tau_2^2 - \tau_1^2}{2} \right) \int_{-\tau_2}^{-\tau_1} \int_{t+\theta}^t g^T(s) T_4 g(s) \, ds d\theta &\leq - \left( \int_{-\tau_2}^{-\tau_1} \int_{t+\theta}^t g(s) \, ds d\theta \right)^T \\ &\quad \times T_4 \left( \int_{-\tau_2}^{-\tau_1} \int_{t+\theta}^t g(s) \, ds d\theta \right). \end{aligned} \quad (32)$$

For positive diagonal matrices  $L$  and  $S$ , we can get from Assumption (H1) that

$$0 \leq \begin{bmatrix} x(t) \\ g(x(t)) \end{bmatrix}^T \begin{bmatrix} -F_1 L & F_2 L \\ F_2 L & -L \end{bmatrix} \begin{bmatrix} x(t) \\ g(x(t)) \end{bmatrix}, \quad (33)$$

$$0 \leq \begin{bmatrix} x(t - \tau(t)) \\ g(x(t - \tau(t))) \end{bmatrix}^T \begin{bmatrix} -F_1 S & F_2 S \\ F_2 S & -S \end{bmatrix} \begin{bmatrix} x(t - \tau(t)) \\ g(x(t - \tau(t))) \end{bmatrix}. \quad (34)$$

From (11) – (13) the following equations are true for any matrices  $M, N, U, Z_i$  ( $i = 1, 2, 3, 4$ ) we have the following equations:

$$0 = 2\zeta^T(t)M \left[ x(t) - x(t - \tau_1) - \int_{t-\tau_1}^t f(s) ds - \int_{t-\tau_1}^t \alpha(s) d\omega(s) \right] \quad (35)$$

$$0 = 2\zeta^T(t)N \left[ x(t - \tau_1) - x(t - \tau(t)) - \int_{t-\tau(t)}^{t-\tau_1} f(s) ds - \int_{t-\tau(t)}^{t-\tau_1} \alpha(s) d\omega(s) \right] \quad (36)$$

$$0 = 2\zeta^T(t)U \left[ x(t - \tau(t)) - x(t - \tau_2) - \int_{t-\tau_2}^{t-\tau(t)} f(s) ds - \int_{t-\tau_2}^{t-\tau(t)} \alpha(s) d\omega(s) \right] \quad (37)$$

$$0 = 2[x^T(t)Z_1 + g^T(t)Z_2] \times [-Ax(t - \delta) + Bg(x(t)) + Cg(x(t - \tau(t)))] \\ + D \int_{t-d(t)}^t g(x(s)) ds + u(t) - g(t) \quad (38)$$

where

$$\zeta^T(t) = \begin{bmatrix} x^T(t) & x^T(t - \tau(t)) & x^T(t - \tau_1) & x^T(t - \tau_2) & g^T(x(t)) & g^T(x(t - \tau(t))) \\ g^T(x(t - \tau_1)) & g^T(x(t - \tau_2)) & x^T(t - \delta) & \int_{t-\delta}^t x^T(s) ds & x^T(t - d(t)) \\ g^T(t) & u^T(t) \left( \int_{t-d(t)}^t g(x(s)) ds \right)^T & \left( \int_{t-\tau_1}^t x(s) ds \right)^T & \left( \int_{t-\tau_2}^{t-\tau_1} x(s) ds \right)^T \\ \left( \int_{t-\tau_1}^t g(x(s)) ds \right)^T & \left( \int_{t-\tau_2}^{t-\tau_1} g(x(s)) ds \right)^T & \left( \int_{-\tau_1}^0 \int_{t+\theta}^t g(s) ds d\theta \right)^T \\ \left( \int_{-\tau_2}^{-\tau_1} \int_{t+\theta}^t g(s) ds d\theta \right)^T \end{bmatrix}$$

From the formula (35) – (37), we have

$$-2\zeta^T(t)M \int_{t-\tau_1}^t \alpha(s) d\omega(s) \leq \zeta^T(t)MX_1^{-1}M^T\zeta(t) + \left( \int_{t-\tau_1}^t \alpha(s) d\omega(s) \right)^T X_1 \\ \times \left( \int_{t-\tau_1}^t \alpha(s) d\omega(s) \right) \quad (39)$$

$$\begin{aligned}
-2\zeta^T(t)N \int_{t-\tau(t)}^{t-\tau_1} \alpha(s) d\omega(s) &\leq \zeta^T(t)NX_2^{-1}N^T\zeta(t) + \left( \int_{t-\tau(t)}^{t-\tau_1} \alpha(s) d\omega(s) \right)^T X_2 \\
&\quad \times \left( \int_{t-\tau(t)}^{t-\tau_1} \alpha(s) d\omega(s) \right) \quad (40)
\end{aligned}$$

$$\begin{aligned}
-2\zeta^T(t)U \int_{t-\tau_2}^{t-\tau(t)} \alpha(s) d\omega(s) &\leq \zeta^T(t)UX_2^{-1}U^T\zeta(t) + \left( \int_{t-\tau_2}^{t-\tau(t)} \alpha(s) d\omega(s) \right)^T X_2 \\
&\quad \times \left( \int_{t-\tau_2}^{t-\tau(t)} \alpha(s) d\omega(s) \right). \quad (41)
\end{aligned}$$

On the other hand, from the Itô isometry in [24], we can obtain

$$\mathbb{E} \left\{ \left[ \int_{t-\tau_1}^t \alpha(s) d\omega(s) \right]^T X_1 \left[ \int_{t-\tau_1}^t \alpha(s) d\omega(s) \right] \right\} = \mathbb{E} \left\{ \int_{t-\tau_1}^t \text{tr}[\alpha^T(s)X_1\alpha(s)] ds \right\} \quad (42)$$

$$\mathbb{E} \left\{ \left[ \int_{t-\tau(t)}^{t-\tau_1} \alpha(s) d\omega(s) \right]^T X_2 \left[ \int_{t-\tau(t)}^{t-\tau_1} \alpha(s) d\omega(s) \right] \right\} = \mathbb{E} \left\{ \int_{t-\tau(t)}^{t-\tau_1} \text{tr}[\alpha^T(s)X_2\alpha(s)] ds \right\} \quad (43)$$

$$\mathbb{E} \left\{ \left[ \int_{t-\tau_2}^{t-\tau(t)} \alpha(s) d\omega(s) \right]^T X_2 \left[ \int_{t-\tau_2}^{t-\tau(t)} \alpha(s) d\omega(s) \right] \right\} = \mathbb{E} \left\{ \int_{t-\tau_2}^{t-\tau(t)} \text{tr}[\alpha^T(s)X_2\alpha(s)] ds \right\} \quad (44)$$

Substituting (16)–(44) into (15), and by the mathematical expectation

$$\begin{aligned}
&\mathbb{E} \{ \mathcal{L}V(x_t, t) - 2y^T(t)u(t) - \gamma u^T(t)u(t) \} \\
&\leq \mathbb{E} \zeta^T(t) \{ \Psi_1 + \tau_1 MT_1^{-1}M^T + (\tau_2 - \tau_1)NT_2^{-1}N^T + (\tau_2 - \tau_1)UT_2^{-1}U^T \\
&\quad + MX_1^{-1}M^T + NX_2^{-1}N^T + UX_2^{-1}U^T \} \zeta(t) \\
&\quad - \int_{t-\tau_1}^t \left[ \zeta^T(t)M + g^T(s)T_1 \right] T_1^{-1} \left[ M^T\zeta(t) + T_1g(s) \right] ds \\
&\quad - \int_{t-\tau(t)}^{t-\tau_1} \left[ \zeta^T(t)N + g^T(s)T_2 \right] T_2^{-1} \left[ N^T\zeta(t) + T_2g(s) \right] ds \\
&\quad - \int_{t-\tau_2}^{t-\tau(t)} \left[ \zeta^T(t)U + g^T(s)T_2 \right] T_2^{-1} \left[ U^T\zeta(t) + T_2g(s) \right] ds \quad (45)
\end{aligned}$$

where  $\Psi_1$  is defined in (10).

Since last three terms in (45) are less than 0, and we can obtain

$$\begin{aligned}
&\mathbb{E} \{ \mathcal{L}V(x_t, t) - 2y^T(t)u(t) - \gamma u^T(t)u(t) \} \\
&\leq \mathbb{E} \zeta^T(t) \{ \Psi_1 + \tau_1 MT_1^{-1}M^T + (\tau_2 - \tau_1)NT_2^{-1}N^T + (\tau_2 - \tau_1)UT_2^{-1}U^T \\
&\quad + MX_1^{-1}M^T + NX_2^{-1}N^T + UX_2^{-1}U^T \} \zeta(t). \quad (46)
\end{aligned}$$

Now we consider the change of  $V(x_t)$  at impulse time  $t = t_k$ ,  $k \in \mathbb{Z}_+$ . From (1) we have

$$x(t_k) - A \int_{t_k-\delta}^{t_k} x(s) ds = x(t_k^-) - I_k \left[ x(t_k^-) - A \int_{t_k-\delta}^{t_k} x(s) ds \right] - A \int_{t_k-\delta}^{t_k} x(s) ds$$

$$= (I - I_k) \left[ x(t_k^-) - A \int_{t_k - \delta}^{t_k} x(s) ds \right]. \quad (47)$$

Moreover, it follows that from (9) that

$$\begin{aligned} & \begin{bmatrix} P_1 & (I - I_k)^T P_1 \\ * & P_1 \end{bmatrix} \geq 0 \\ \Leftrightarrow & \begin{bmatrix} I & -(I - I_k)^T \\ 0 & I \end{bmatrix} \begin{bmatrix} P_1 & (I - I_k)^T P_1 \\ * & P_1 \end{bmatrix} \begin{bmatrix} I & 0 \\ -(I - I_k) & I \end{bmatrix} \geq 0 \\ \Leftrightarrow & \begin{bmatrix} P_1 - (I - I_k)^T P_1 (I - I_k) & 0 \\ * & P_1 \end{bmatrix} \geq 0 \\ \Leftrightarrow & P_1 - (I - I_k)^T P_1 (I - I_k) \geq 0. \end{aligned} \quad (48)$$

Together with (47) and (48), it yields

$$\begin{aligned} V_1(x(t_k)) &= \left[ x(t_k) - A \int_{t_k - \delta}^{t_k} x(s) ds \right]^T P_1 \left[ x(t_k) - A \int_{t_k - \delta}^{t_k} x(s) ds \right] \\ &= \left[ x(t_k^-) - A \int_{t_k - \delta}^{t_k} x(s) ds \right]^T (I - I_k)^T P_1 (I - I_k) \left[ x(t_k^-) - A \int_{t_k - \delta}^{t_k} x(s) ds \right] \\ &\leq \left[ x(t_k^-) - A \int_{t_k - \delta}^{t_k} x(s) ds \right]^T P_1 \left[ x(t_k^-) - A \int_{t_k - \delta}^{t_k} x(s) ds \right] \\ V_1(x(t_k)) &= V_1(x(t_k^-)). \end{aligned}$$

Moreover it is obvious that  $V_2(t_k) = V_2(t_k^-)$ ,  $V_3(t_k) = V_3(t_k^-)$ ,  $V_4(t_k) = V_4(t_k^-)$ ,  $V_5(t_k) = V_5(t_k^-)$ ,  $V_6(t_k) = V_6(t_k^-)$ ,  $V_7(t_k) = V_7(t_k^-)$ ,  $V_8(t_k) = V_8(t_k^-)$ ,  $V_9(t_k) = V_9(t_k^-)$ . which implies that

$$V(x(t_k)) = V(x(t_k^-)), \quad k \in \mathbb{Z}_+. \quad (49)$$

It follows from (46) that

$$\begin{aligned} & \mathbb{E}dV(x_t, t) - 2\mathbb{E}y^T(t)u(t) - \gamma\mathbb{E}u^T(t)u(t) \\ &= \mathbb{E}\mathcal{L}V(x_t, t) - 2\mathbb{E}y^T(t)u(t) - \gamma\mathbb{E}u^T(t)u(t) \\ &\leq \mathbb{E}\zeta^T(t) \left\{ \Psi_1 + \tau_1 M T_1^{-1} M^T + (\tau_2 - \tau_1) N T_2^{-1} N^T + (\tau_2 - \tau_1) U T_2^{-1} U^T \right. \\ &\quad \left. + M X_1^{-1} M^T + N X_2^{-1} N^T + U X_2^{-1} U^T \right\} \zeta(t). \end{aligned}$$

Let

$$\hat{\Psi} = \Psi_1 + \tau_1 M T_1^{-1} M^T + (\tau_2 - \tau_1) N T_2^{-1} N^T + (\tau_2 - \tau_1) U T_2^{-1} U^T$$

$$+MX_1^{-1}M^T + NX_2^{-1}N^T + UX_2^{-1}U^T.$$

By applying Schur complement [2], it is easy to see that  $\hat{\Psi}$  is equivalent to (10), then we can obtain

$$\mathbb{E}dV(x_t, t) - 2\mathbb{E}y^T(t)u(t) - \gamma u^T(t)u(t) \leq 0. \quad (50)$$

By integrating (50) over the time period from 0 to  $t_f$ , we have

$$2\mathbb{E} \int_0^{t_f} y^T(s)u(s) ds \geq \mathbb{E}V(x_0, 0) - \gamma \mathbb{E} \int_0^{t_f} u^T(s)u(s) ds \geq -\gamma \mathbb{E} \int_0^{t_f} u^T(s)u(s) ds. \quad (51)$$

From Definition 1, we know that the stochastic neural network (1) is passive in the sense of expectation. This completes the proof.  $\square$

**Remark 1.** In the proof of Theorem 3.1, we introduce a new estimation on the upper bound of the time derivative of  $V(t)$ . For this, we introduce the following inequalities,

$$\begin{aligned} \tau_1 \zeta^T(t)MT_1^{-1}M^T \zeta(t) - \int_{t-\tau_1}^t \zeta^T(s)MT_1^{-1}M^T \zeta(s) ds &\geq 0, \\ (\tau_2 - \tau_1)NT_2^{-1}N^T \zeta(t) - \int_{t-\tau(t)}^{t-\tau_1} \zeta^T(s)NT_2^{-1}N^T \zeta(s) ds &\geq 0, \\ (\tau_2 - \tau_1)UT_2^{-1}U^T \zeta(t) - \int_{t-\tau_2}^{t-\tau(t)} \zeta^T(s)UT_2^{-1}U^T \zeta(s) ds &\geq 0, \end{aligned}$$

where it is employed in [10]; and  $\tau(t) - \tau_1$ ,  $\tau_2 - \tau(t)$  were enlarged to  $\tau_2 - \tau_1$ . It is easy to see that this treatment is more conservative than the expression in the proof of Theorem 3.1.

**Remark 2.** It should be pointed out that the range of the time-varying delays in [36] is varying from 0 to upper bounds. However, in many practical cases [5, 36, 37], the time delays may typically exist on intervals, where the lower bounds of the time-varying delays are not restricted to be 0. In this work, the time-varying delays are assumed to be intervals, which means that the lower and upper bounds of interval time-varying delay is available, where  $\tau(t) \in [\tau_1, \tau_2]$ . On the other hand distributed delays, parameter uncertainties and impulsive perturbations are considered; see Theorem 3.1.

**Remark 3.** In order to reduce the conservativeness, when obtaining the derivative of  $V_7(x_t, t)$ ,  $V_9(x_t, t)$  the integral terms  $\int_{t-\tau_2}^{t-\tau_1} g^T(s)T_2g(s) ds$ ,  $\int_{t-\tau_2}^{t-\tau_1} \alpha^T(s)X_2\alpha(s) ds$  is divided into two parts as  $\int_{t-\tau(t)}^{t-\tau_1} g^T(s)T_2g(s) ds$ ,  $\int_{t-\tau_2}^{t-\tau(t)} g^T(s)T_2g(s) ds$  and  $\int_{t-\tau(t)}^{t-\tau_1} \alpha^T(s)X_2\alpha(s) ds$ ,  $\int_{t-\tau_2}^{t-\tau(t)} \alpha^T(s)X_2\alpha(s) ds$  respectively, which is mainly based on the information about  $\tau_1 \leq \tau(t) \leq \tau_2$ , which may leads to less conservative results.

**Remark 4.** In Assumption (H1), the constants  $F_j^-$  and  $F_j^+$   $j = 1, 2, \dots, n$  are allowed to be positive, negative or zero. However, in [5, 26, 35], the Lipschitz constants are only allowed to be positive. Hence, Assumption(H1), first proposed by Liu et al.in [23], is weaker than the assumption in [5, 26, 35].

#### 4. PASSIVITY FOR UNCERTAIN STOCHASTIC NEURAL NETWORKS WITH IMPULSES

In this section, we extend the previous passivity condition to the following uncertain stochastic neural network:

$$\begin{aligned}
dx(t) &= \left[ -(A + \Delta A(t))x(t - \delta) + (B + \Delta B(t))g(x(t)) + (C + \Delta C(t))g(x(t - \tau(t))) \right. \\
&\quad \left. + (D + \Delta D(t)) \int_{t-d(t)}^t g(x(s)) ds + u(t) \right] dt \\
&\quad + \sigma(t, x(t), x(t - \tau(t)), x(t - d(t))) d\omega(t), \quad t \neq t_k \\
y(t) &= g(x(t)) \\
\Delta x(t_k) &= -I_k \left\{ x(t_k^-) - A \int_{t_k - \delta}^{t_k} x(s) ds \right\}, \quad t = t_k, \quad k \in \mathbb{Z}_+, \quad (52)
\end{aligned}$$

where  $\Delta A(t), \Delta B(t), \Delta C(t), \Delta D(t)$  are the time varying uncertainties of the form:

$$\begin{bmatrix} \Delta A(t) & \Delta B(t) & \Delta C(t) & \Delta D(t) \end{bmatrix} = HF(t) \begin{bmatrix} G_1 & G_2 & G_3 & G_4 \end{bmatrix} \quad (53)$$

where  $H, G_i (i = 1, 2, 3, 4)$  are known real constant matrices,  $F(t)$  is the time-varying uncertain matrices, which satisfies  $F^T(t)F(t) \leq I$ .

**Theorem 4.1.** Assume that assumptions (H1)–(H3) holds. For given scalars  $\tau_1, \tau_2, \mu$  and  $d$ , the stochastic neural network described by (52) is stochastically passive in the sense of Definition 1, for any time varying delay  $\tau(t)$  and  $d(t)$  satisfying (2), if there exist constant scalars  $\lambda_i > 0 (i = 1, 2, 3), \gamma > 0$ , positive definite symmetric matrices  $P_i (i = 1, 2, \dots, 5), Q_i, T_i (i = 1, 2, \dots, 4), X_1, X_2, \angle = \begin{bmatrix} L_1 & L_2 \\ L_2^T & L_3 \end{bmatrix}, \Re = \begin{bmatrix} R_1 & R_2 \\ R_2^T & R_3 \end{bmatrix}, \Im = \begin{bmatrix} S_1 & S_2 \\ S_2^T & S_3 \end{bmatrix}$ , positive diagonal matrices  $L, S$ , real matrices  $Z_1, Z_2, M_i, N_i, U_i (i = 1, 2, \dots, 8)$  such that the following LMI holds:

$$P_1 < \lambda_1 I, \quad (54)$$

$$X_1 < \lambda_2 I, \quad (55)$$

$$X_2 < \lambda_3 I, \quad (56)$$

$$\begin{bmatrix} P_1 & (I - I_k)^T P_1 \\ * & P_1 \end{bmatrix} \geq 0, \quad k \in \mathbb{Z}_+, \quad (57)$$

and

$$\Psi = \begin{pmatrix} \Psi_1 & \sqrt{\tau_1} M & \sqrt{\tau_2 - \tau_1} N & \sqrt{\tau_2 - \tau_1} U & M & N & U & \Gamma_{d1} & \Gamma_{d2} & \Gamma_{d3} \\ * & -T_1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ * & * & -T_2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ * & * & * & -T_2 & 0 & 0 & 0 & 0 & 0 & 0 \\ * & * & * & * & -X_1 & 0 & 0 & 0 & 0 & 0 \\ * & * & * & * & * & -X_2 & 0 & 0 & 0 & 0 \\ * & * & * & * & * & * & -X_2 & 0 & 0 & 0 \\ * & * & * & * & * & * & * & -\frac{1}{4} I & 0 & 0 \\ * & * & * & * & * & * & * & * & -\frac{1}{4} I & 0 \\ * & * & * & * & * & * & * & * & * & -\frac{1}{4} I \end{pmatrix} < 0, \quad (58)$$

where

$$\begin{aligned}\Psi_{55} &= d^2 P_5 + L_3 + \tau_1^2 Q_3 + (\tau_2 - \tau_1)^2 Q_4 - L + 4G_2^T G_2, \\ \Psi_{66} &= -(1 - \mu)R_3 - S + 4G_3^T G_3, \\ \Psi_{99} &= -P_2 + 4G_1^T G_1, \quad \Psi_{14_{14}} = -P_5 + 4G_4^T G_4.\end{aligned}$$

$$\begin{aligned}\Gamma_{d1} &= \text{col} [ P_1 H + Z_1 H \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 ] \\ \Gamma_{d2} &= \text{col} [ 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad Z_2 H \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 ] \\ \Gamma_{d3} &= \text{col} [ 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad A^T P_1 H \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 ]\end{aligned}$$

and other terms are same as defined in Theorem 3.1.

*Proof.* It is not difficult to check that system (52) is equivalent to the following form:

$$\begin{aligned}d \left[ x(t) - A \int_{t-\delta}^t x(s) ds \right] &= \left\{ -Ax(t) - \Delta A(t)x(t-\delta) + [B + \Delta B(t)]g(x(t)) \right. \\ &\quad + [C + \Delta C(t)]g(x(t-\tau(t))) + [D + \Delta D(t)] \\ &\quad \left. \times \int_{t-d(t)}^t g(x(s)) ds + u(t) \right\} dt \\ &\quad + \sigma(t, x(t), x(t-\tau(t)), x(t-d(t))) d\omega(t).\end{aligned}\quad (59)$$

Then, from Theorem 3.1, we only need to estimate the following equalities:

$$\begin{aligned}\mathcal{L}V_1(x_t, t) &= 2 \left[ x(t) - A \int_{t-\delta}^t x(s) ds \right]^T P_1 \left[ -Ax(t) - Ax(t-\delta) \right. \\ &\quad + (B + \Delta B)g(x(t)) + (C + \Delta C)g(x(t-\tau(t))) \\ &\quad \left. + (D + \Delta D) \int_{t-d(t)}^t g(x(s)) ds + u(t) \right] + \alpha^T(t) P_1 \alpha(t) \\ 0 &= 2[x^T(t)Z_1 + f^T(t)Z_2] \times [- (A + \Delta A)x(t-\delta) + (B + \Delta B)g(x(t)) \\ &\quad + (C + \Delta C)g(x(t-\tau(t))) + (D + \Delta D) \int_{t-d(t)}^t g(x(s)) ds \\ &\quad + u(t) - g(t)].\end{aligned}\quad (60)$$

$$\begin{aligned}0 &= 2[x^T(t)Z_1 + f^T(t)Z_2] \times [- (A + \Delta A)x(t-\delta) + (B + \Delta B)g(x(t)) \\ &\quad + (C + \Delta C)g(x(t-\tau(t))) + (D + \Delta D) \int_{t-d(t)}^t g(x(s)) ds \\ &\quad + u(t) - g(t)].\end{aligned}\quad (61)$$

Replace  $\Delta A$ ,  $\Delta B$ ,  $\Delta C$ ,  $\Delta D$  with  $HF(t)G_1$ ,  $HF(t)G_2$ ,  $HF(t)G_3$ ,  $HF(t)G_4$  respectively, and using Lemma 2.3 in (60) and (61), we can obtain

$$\begin{aligned}-2x^T(t)P_1\Delta A(t)x(t-\delta) &= -2x^T(t)P_1HF(t)G_1x(t-\delta) \\ &\leq x^T(t)P_1HF(t)F^T(t)H^T P_1x(t) \\ &\quad + x^T(t-\delta)G_1^T G_1x(t-\delta) \\ &\leq x^T(t)P_1HH^T P_1x(t) \\ &\quad + x^T(t-\delta)G_1^T G_1x(t-\delta)\end{aligned}\quad (62)$$

$$\begin{aligned}
2x^T(t)P_1\Delta B(t)g(x(t)) &= 2x^T(t)P_1HF(t)G_2g(x(t)) \\
&\leq x^T(t)P_1HF(t)F^T(t)H^T P_1x(t) \\
&\quad + g^T(x(t))G_2^T G_2g(x(t)) \\
&\leq x^T(t)P_1HH^T P_1x(t) \\
&\quad + g^T(x(t))G_2^T G_2g(x(t)) \tag{63}
\end{aligned}$$

$$\begin{aligned}
2x^T(t)P_1\Delta C(t)g(x(t-\tau(t))) &= 2x^T(t)P_1HF(t)G_3g(x(t-\tau(t))) \\
&\leq x^T(t)P_1HF(t)F^T(t)H^T P_1x(t) \\
&\quad + g^T(x(t-\tau(t)))G_3^T G_3g(x(t-\tau(t))) \\
&\leq x^T(t)P_1HH^T P_1x(t) \\
&\quad + g^T(x(t-\tau(t)))G_3^T G_3g(x(t-\tau(t))) \tag{64}
\end{aligned}$$

$$\begin{aligned}
2x^T(t)P_1\Delta D(t) \int_{t-d(t)}^t g(x(s)) ds &= 2x^T(t)P_1HF(t)G_4 \int_{t-d(t)}^t g(x(s)) ds \\
&\leq x^T(t)P_1HF(t)F^T(t)H^T P_1x(t) \\
&\quad + \left( \int_{t-d(t)}^t g(x(s)) ds \right)^T G_4^T G_4 \\
&\quad \times \left( \int_{t-d(t)}^t g(x(s)) ds \right) \\
&\leq x^T(t)P_1HH^T P_1x(t) \\
&\quad + \left( \int_{t-d(t)}^t g(x(s)) ds \right)^T G_4^T G_4 \\
&\quad \times \left( \int_{t-d(t)}^t g(x(s)) ds \right) \tag{65}
\end{aligned}$$

$$\begin{aligned}
2\left( \int_{t-\delta}^t x(s) ds \right)^T A^T P_1\Delta A(t)x(t-\delta) &= 2\left( \int_{t-\delta}^t x(s) ds \right)^T A^T P_1HF(t)G_1x(t-\delta) \\
&\leq \left( \int_{t-\delta}^t x(s) ds \right)^T A^T P_1HF(t)F^T(t)H^T P_1A \\
&\quad \times \left( \int_{t-\delta}^t x(s) ds \right) \\
&\quad + x^T(t-\delta)G_1^T G_1x(t-\delta) \\
&\leq \left( \int_{t-\delta}^t x(s) ds \right)^T A^T P_1HH^T P_1A \\
&\quad \times \left( \int_{t-\delta}^t x(s) ds \right) \\
&\quad + x^T(t-\delta)G_1^T G_1x(t-\delta) \tag{66}
\end{aligned}$$

$$2\left( \int_{t-\delta}^t x(s) ds \right)^T A^T P_1\Delta B(t)g(x(t)) = 2\left( \int_{t-\delta}^t x(s) ds \right)^T A^T P_1HF(t)G_2g(x(t))$$



$$\begin{aligned}
&\leq \left( \int_{t-\delta}^t x(s) ds \right)^T A^T P_1 H F(t) F^T(t) H^T P_1 A \\
&\quad \times \left( \int_{t-\delta}^t x(s) ds \right) \\
&\quad + g^T(x(t)) G_2^T G_2 g(x(t)) \\
&\leq \left( \int_{t-\delta}^t x(s) ds \right)^T A^T P_1 H H^T P_1 A \\
&\quad \times \left( \int_{t-\delta}^t x(s) ds \right) \\
&\quad + g^T(x(t)) G_2^T G_2 g(x(t)) \tag{67}
\end{aligned}$$

$$\begin{aligned}
&2 \left( \int_{t-\delta}^t x(s) ds \right)^T A^T P_1 \Delta C(t) g(x(t-\tau(t))) \\
&= 2 \left( \int_{t-\delta}^t x(s) ds \right)^T A^T P_1 H F(t) G_3 g(x(t-\tau(t))) \\
&\leq \left( \int_{t-\delta}^t x(s) ds \right)^T A^T P_1 H F(t) F^T(t) H^T P_1 A \left( \int_{t-\delta}^t x(s) ds \right) \\
&\quad + g^T(x(t-\tau(t))) G_3^T G_3 g(x(t-\tau(t))) \\
&\leq \left( \int_{t-\delta}^t x(s) ds \right)^T A^T P_1 H H^T P_1 A \left( \int_{t-\delta}^t x(s) ds \right) + g^T(x(t-\tau(t))) \\
&\quad \times G_3^T G_3 g(x(t-\tau(t))) \tag{68}
\end{aligned}$$

$$\begin{aligned}
&2 \left( \int_{t-\delta}^t x(s) ds \right)^T A^T P_1 \Delta D(t) \left( \int_{t-d(t)}^t g(x(s)) ds \right) \\
&= 2 \left( \int_{t-\delta}^t x(s) ds \right)^T A^T P_1 H F(t) G_4 \\
&\quad \times \left( \int_{t-d(t)}^t g(x(s)) ds \right) \\
&\leq \left( \int_{t-\delta}^t x(s) ds \right)^T A^T P_1 H F(t) F^T(t) H^T P_1 A \left( \int_{t-\delta}^t x(s) ds \right) \\
&\quad + \left( \int_{t-d(t)}^t g(x(s)) ds \right)^T G_4^T G_4 \left( \int_{t-d(t)}^t g(x(s)) ds \right) \\
&\leq \left( \int_{t-\delta}^t x(s) ds \right)^T A^T P_1 H H^T P_1 A \left( \int_{t-\delta}^t x(s) ds \right) \\
&\quad + \left( \int_{t-d(t)}^t g(x(s)) ds \right)^T G_4^T G_4 \left( \int_{t-d(t)}^t g(x(s)) ds \right) \tag{69}
\end{aligned}$$

$$-2x^T(t) Z_1 \Delta A(t) x(t-\delta) = -2x^T(t) Z_1 H F(t) G_1 x(t-\delta)$$

$$\begin{aligned}
&\leq x^T(t)Z_1HF(t)F^T(t)H^TZ_1^Tx(t) \\
&\quad + x^T(t-\delta)G_1^TG_1x(t-\delta) \\
&\leq x^T(t)Z_1HH^TZ_1^Tx(t) \\
&\quad + x^T(t-\delta)G_1^TG_1x(t-\delta) \tag{70}
\end{aligned}$$

$$\begin{aligned}
2x^T(t)Z_1\Delta B(t)g(x(t)) &= 2x^T(t)Z_1HF(t)G_2g(x(t)) \\
&\leq x^T(t)Z_1HF(t)F^T(t)H^TZ_1^Tx(t) \\
&\quad \times + g^T(x(t))G_2^TG_2g(x(t)) \\
&\leq x^T(t)Z_1HH^TZ_1^Tx(t) \\
&\quad \times + g^T(x(t))G_2^TG_2g(x(t)) \tag{71}
\end{aligned}$$

$$\begin{aligned}
2x^T(t)Z_1\Delta C(t)g(x(t-\tau(t))) &= 2x^T(t)Z_1HF(t)G_3g(x(t-\tau(t))) \\
&\leq x^T(t)Z_1HF(t)F^T(t)H^TZ_1^Tx(t) \\
&\quad + g^T(x(t-\tau(t)))G_3^TG_3 \\
&\quad \times g(x(t-\tau(t))) \\
&\leq x^T(t)Z_1HH^TZ_1^Tx(t) \\
&\quad + g^T(x(t-\tau(t)))G_3^TG_3 \\
&\quad \times g(x(t-\tau(t))) \tag{72}
\end{aligned}$$

$$\begin{aligned}
2x^T(t)Z_1\Delta D(t)\int_{t-d(t)}^t g(x(s)) ds &= 2x^T(t)Z_1HF(t)G_4\int_{t-d(t)}^t g(x(s)) ds \\
&\leq x^T(t)Z_1HF(t)F^T(t)H^TZ_1^Tx(t) \\
&\quad + \left(\int_{t-d(t)}^t g(x(s)) ds\right)^T G_4^TG_4 \\
&\quad \times \left(\int_{t-d(t)}^t g(x(s)) ds\right) \\
&\leq x^T(t)Z_1HH^TZ_1^Tx(t) + \left(\int_{t-d(t)}^t g(x(s)) ds\right)^T \\
&\quad \times G_4^TG_4\left(\int_{t-d(t)}^t g(x(s)) ds\right) \tag{73}
\end{aligned}$$

$$\begin{aligned}
-2f^T(t)Z_2\Delta A(t)x(t-\delta) &= -2f^T(t)Z_2HF(t)G_1x(t-\delta) \\
&\leq f^T(t)Z_2HF(t)F^T(t)H^TZ_2^Tf(t) \\
&\quad \times + x^T(t-\delta)G_1^TG_1x(t-\delta) \\
&\leq f^T(t)Z_2HH^TZ_2^Tf(t) \\
&\quad + x^T(t-\delta)G_1^TG_1x(t-\delta) \tag{74}
\end{aligned}$$

$$\begin{aligned}
2f^T(t)Z_2\Delta B(t)g(x(t)) &= 2f^T(t)Z_2HF(t)G_2g(x(t)) \\
&\leq f^T(t)Z_2HF(t)F^T(t)H^TZ_2^Tf(t) \\
&\quad + g^T(x(t))G_2^TG_2g(x(t)) \\
&\leq f^T(t)Z_2HH^TZ_2^Tf(t)
\end{aligned}$$

$$\begin{aligned} & \times + g^T(x(t))G_2^T G_2 g(x(t)) \end{aligned} \quad (75)$$

$$\begin{aligned} 2f^T(t)Z_2\Delta C(t)g(x(t-\tau(t))) &= 2f^T(t)Z_2HF(t)G_3g(x(t-\tau(t))) \\ &\leq f^T(t)Z_2HF(t)F^T(t)H^T Z_2^T f(t) \\ &\quad + g^T(x(t-\tau(t)))G_3^T G_3g(x(t-\tau(t))) \\ &\leq f^T(t)Z_2HH^T Z_2^T f(t) \\ &\quad + g^T(x(t-\tau(t)))G_3^T G_3g(x(t-\tau(t))) \end{aligned} \quad (76)$$

$$\begin{aligned} 2f^T(t)Z_2\Delta D(t) \int_{t-d(t)}^t g(x(s)) ds &= 2f^T(t)Z_2HF(t)G_4 \int_{t-d(t)}^t g(x(s)) ds \\ &\leq f^T(t)Z_2HF(t)F^T(t)H^T Z_2^T f(t) \\ &\quad + \left( \int_{t-d(t)}^t g(x(s)) ds \right)^T G_4^T G_4 \\ &\quad \times \left( \int_{t-d(t)}^t g(x(s)) ds \right) \\ &\leq f^T(t)Z_2HH^T Z_2^T f(t) \\ &\quad + \left( \int_{t-d(t)}^t g(x(s)) ds \right)^T G_4^T G_4 \\ &\quad \times \left( \int_{t-d(t)}^t g(x(s)) ds \right). \end{aligned} \quad (77)$$

Then along the same line as for Theorem 3.1, we can obtain the desired result by applying Lemma 2.2 and (62)–(77). This completes the proof of Theorem 4.1.  $\square$

**Remark 5.** From proof of above Theorems 3.1 and 4.1, we can see that the novelty of the Lyapunov functional contains, quadratic Lyapunov–Krasovskii functional terms in  $V_4(x_t, t)$ , and triple-integral terms in  $V_8(x_t, t)$  are introduced, which can be expected to reduce the conservatism. More specifically to improve the feasible region for the corresponding system, by taking the states  $\int_{t-\tau_1}^t x^T(s) ds$ ,  $\int_{t-\tau_2}^{t-\tau_1} x^T(s) ds$ ,  $\int_{t-\tau_1}^t g^T(x(s)) ds$ ,  $\int_{t-\tau_2}^{t-\tau_1} g^T(x(s)) ds$ ,  $\int_{-\tau_2}^{-\tau_1} \int_{t+\theta}^t g(s) ds d\theta$ ,  $\int_{-\tau_2}^{-\tau_1} \int_{t+\theta}^t \alpha^T(s) ds d\theta$ , the passivity conditions in Theorems 3.1 and 4.1 sufficiently utilize more information on state variables, which can yield less conservatism.

**Remark 6.** In [5, 20], the authors discussed the passivity analysis of stochastic neural networks with time varying delay. But in this paper, we have studied passivity analysis of stochastic neural networks with leakage and distributed delays using impulse control. Moreover, different from the previous literature, our results are derived by constructing a new Lyapunov–Krasovskii functional with triple integral terms with bounding technique. In addition, some free weighting matrices are introduced in Theorem 3.1 for getting feasible solution. To ascertain the passivity for the stochastic neural network with time delays in the leakage terms, Theorem 4.1 further presents sufficient conditions for the correspondent system with unknown parameters (58). Hence, in our knowledge, passivity problem of stochastic interval neural network with distributed delays in the

leakage terms using impulsive perturbations has never been studied in the previous literature and it is essentially new.

## 5. NUMERICAL EXAMPLES

In this section, we are analyzing examples showing the effectiveness of the proposed methods.

**Example 1.** Consider the stochastic neural network with time varying delays and impulses in (1). The parametric coefficients are

$$A = \begin{bmatrix} 5 & 0 \\ 0 & 7 \end{bmatrix}, \quad B = \begin{bmatrix} 0.3 & -4 \\ 0.1 & 0.3 \end{bmatrix}, \quad C = \begin{bmatrix} 0.4 & 0.2 \\ 0.1 & 0.6 \end{bmatrix}, \quad D = \begin{bmatrix} 0.4 & -0.3 \\ 0.1 & 0.6 \end{bmatrix},$$

$$\Sigma_1 = \begin{bmatrix} 0.3 & 0.1 \\ 0.5 & -0.1 \end{bmatrix}, \quad \Sigma_2 = \begin{bmatrix} -0.1 & 0.2 \\ 0 & 0.1 \end{bmatrix}, \quad \Sigma_3 = \begin{bmatrix} 0.4 & 0.3 \\ 0.2 & 0 \end{bmatrix}, \quad I_k = \begin{bmatrix} 0.5 & 0 \\ 0 & 0.5 \end{bmatrix}.$$

The activation functions are taken as follows:

$$g_1(x) = \frac{1}{20}(|x+1| + |x-1|), \quad g_2(x) = \frac{1}{10}(|x+1| + |x-1|).$$

It can be verified that Assumption (H1) is satisfied with  $F_1^- = -0.1$ ,  $F_1^+ = 0.1$ ,  $F_2^- = -0.2$ ,  $F_2^+ = 0.2$ . Thus

$$F_1 = \begin{bmatrix} -0.01 & 0 \\ 0 & -0.04 \end{bmatrix}, \quad F_2 = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix},$$

Our main purpose in this example is to estimate the maximum allowable upper bound delay  $\tau_2$ ,  $d$  for given lower bound  $\tau_1$ . For  $\tau_1 = 1.6$ ,  $\mu = 0.5$ ,  $\delta = 0.1$ , by solving LMIs (6)–(10) in Theorem 3.1 using MATLAB LMI toolbox, one can obtain the feasible solution for any time delay satisfying  $0 < d(t) \leq 1.5243$  and  $0 \leq 1.6 < \tau(t) \leq 3.2458$ . For example, if we take  $\mu = 0.5$ ,  $\delta = 0.1$ ,  $d = 0.8$ ,  $\tau_1 = 1.6$  and  $\tau_2 = 2.8$  we obtain the following feasible solutions

$$P_1 = \begin{bmatrix} 175.1815 & 0.1912 \\ 0.1912 & 220.2178 \end{bmatrix}, \quad P_2 = \begin{bmatrix} 101.2050 & 0.0368 \\ 0.0368 & 138.3154 \end{bmatrix}, \quad P_3 = 10^4 \times \begin{bmatrix} 2.2897 & 0.0987 \\ 0.0987 & 7.5980 \end{bmatrix},$$

$$P_4 = \begin{bmatrix} 54.1964 & 22.1242 \\ 22.1242 & 33.6705 \end{bmatrix}, \quad P_5 = 10^3 \times \begin{bmatrix} 1.1855 & -0.0208 \\ -0.0208 & 0.6355 \end{bmatrix}, \quad L_1 = \begin{bmatrix} 72.6224 & -5.3054 \\ -5.3054 & 93.0195 \end{bmatrix},$$

$$L_2 = \begin{bmatrix} 110.0120 & 36.3356 \\ 36.3356 & 106.1753 \end{bmatrix}, \quad L_3 = 10^3 \times \begin{bmatrix} 2.1376 & 0.2181 \\ 0.2181 & 1.0399 \end{bmatrix}, \quad R_1 = \begin{bmatrix} 38.0286 & -5.9779 \\ -5.9779 & 59.6507 \end{bmatrix},$$

$$R_2 = \begin{bmatrix} 71.5361 & 13.2683 \\ 13.2683 & 55.1780 \end{bmatrix}, \quad R_3 = \begin{bmatrix} 659.6061 & 116.7109 \\ 116.7109 & 362.1591 \end{bmatrix}, \quad S_1 = \begin{bmatrix} 23.0397 & 1.2640 \\ 1.2640 & 23.5236 \end{bmatrix},$$

$$\begin{aligned}
S_2 &= \begin{bmatrix} 63.7244 & 14.1845 \\ 14.1845 & 55.4936 \end{bmatrix}, S_3 = \begin{bmatrix} 909.2693 & 88.9898 \\ 88.9898 & 459.1695 \end{bmatrix}, Q_1 = \begin{bmatrix} 7.1802 & -0.0377 \\ -0.0377 & 7.1061 \end{bmatrix}, \\
Q_2 &= \begin{bmatrix} 12.6557 & -0.0667 \\ -0.0667 & 12.5245 \end{bmatrix}, Q_3 = \begin{bmatrix} 432.3567 & 35.8151 \\ 35.8151 & 154.1902 \end{bmatrix}, Q_4 = \begin{bmatrix} 655.5405 & 51.1455 \\ 51.1455 & 257.6060 \end{bmatrix}, \\
T_1 &= \begin{bmatrix} 0.2454 & 0.0069 \\ 0.0069 & 0.02521 \end{bmatrix}, T_2 = \begin{bmatrix} 0.4118 & 0.0088 \\ 0.0088 & 0.4212 \end{bmatrix}, T_3 = \begin{bmatrix} 0.2455 & 0.0039 \\ 0.0039 & 0.2494 \end{bmatrix}, \\
T_4 &= \begin{bmatrix} 0.1305 & 0.0009 \\ 0.0009 & 0.1314 \end{bmatrix}, X_1 = \begin{bmatrix} 18.1162 & -0.1125 \\ -0.1125 & 17.9124 \end{bmatrix}, X_2 = \begin{bmatrix} 27.0286 & -0.1404 \\ -0.1404 & 26.7514 \end{bmatrix}, \\
L &= 10^3 \times \begin{bmatrix} 6.1240 & 0 \\ 0 & 5.1268 \end{bmatrix}, S = \begin{bmatrix} 870.8465 & 0 \\ 0 & 287.0032 \end{bmatrix}, M_1 = \begin{bmatrix} -0.2052 & 0.0148 \\ 0.0148 & -0.1210 \end{bmatrix}, \\
M_2 &= 10^3 \times \begin{bmatrix} 0.0658 & -0.6740 \\ -0.6740 & -0.2752 \end{bmatrix}, M_3 = \begin{bmatrix} 0.0930 & 0.0038 \\ 0.0038 & 0.0981 \end{bmatrix}, \\
M_5 &= \begin{bmatrix} 0.2946 & 0.2981 \\ 0.2981 & -0.3589 \end{bmatrix}, M_6 = \begin{bmatrix} -0.2283 & 0.0041 \\ 0.0041 & 0.0092 \end{bmatrix}, M_7 = \begin{bmatrix} -0.0116 & -0.0066 \\ -0.0066 & 0.0051 \end{bmatrix}, \\
M_8 &= \begin{bmatrix} 0.0026 & 0.0018 \\ 0.0018 & -0.0013 \end{bmatrix}, N_1 = \begin{bmatrix} -0.0720 & 0.0054 \\ 0.0054 & -0.0113 \end{bmatrix}, N_2 = \begin{bmatrix} 0.2607 & 0.0065 \\ 0.0065 & 0.2681 \end{bmatrix}, \\
N_3 &= \begin{bmatrix} -0.2602 & -0.0076 \\ -0.0076 & -0.2680 \end{bmatrix}, N_4 = \begin{bmatrix} 0.6438 & -0.4825 \\ -0.4825 & 0.3821 \end{bmatrix}, N_5 = \begin{bmatrix} 0.2124 & 0.1873 \\ 0.1873 & -0.0754 \end{bmatrix}, \\
N_6 &= \begin{bmatrix} -0.1449 & -0.0053 \\ -0.0053 & -0.0131 \end{bmatrix}, N_7 = \begin{bmatrix} -0.0108 & -0.0033 \\ -0.0033 & 0.0006 \end{bmatrix}, N_8 = \begin{bmatrix} 0.0017 & 0.0010 \\ 0.0010 & -0.0004 \end{bmatrix}, \\
U_1 &= \begin{bmatrix} -0.0187 & 0.0132 \\ 0.0132 & -0.0102 \end{bmatrix}, U_2 = \begin{bmatrix} -0.2606 & -0.0070 \\ -0.0070 & -0.2684 \end{bmatrix}, U_3 = \begin{bmatrix} -0.1075 & 0.0972 \\ 0.0972 & 0.4607 \end{bmatrix}, \\
U_4 &= \begin{bmatrix} 0.2589 & 0.0067 \\ 0.0067 & 0.2669 \end{bmatrix}, U_5 = \begin{bmatrix} 0.0138 & 0.0581 \\ 0.0581 & -0.2713 \end{bmatrix}, U_6 = \begin{bmatrix} -0.0379 & 0.0139 \\ 0.0139 & 0.0248 \end{bmatrix}, \\
U_7 &= \begin{bmatrix} -0.0026 & -0.0004 \\ -0.0004 & 0.0039 \end{bmatrix}, U_8 = \begin{bmatrix} -0.0080 & -0.0018 \\ -0.0018 & -0.0088 \end{bmatrix}, Z_1 = \begin{bmatrix} -1.6385 & 0.6544 \\ 0.6544 & -0.3471 \end{bmatrix}, \\
Z_2 &= \begin{bmatrix} 2.5340 & 0.0210 \\ 0.0210 & 2.4667 \end{bmatrix}, M_4 = 10^3 \times \begin{bmatrix} 0.8794 & -0.7704 \\ -0.7704 & 0.8027 \end{bmatrix}, \lambda_1 = 215.1348, \\
\lambda_2 &= 247.3246, \lambda_3 = 198.5241, \gamma = 405.2402. \text{ For given initial state } x(t) = [0.5 \quad -0.5]^T, \\
\text{Figure 1 gives the state trajectory of the neural network (1) under zero input, which} \\
\text{shows that the neural network is stable.}
\end{aligned}$$




Fig. 1. State trajectory of the NNs in Example 1.

**Example 2.** Consider the uncertain stochastic neural network with time varying delays and impulses (52) with the following parameters

$$A = \begin{bmatrix} 4 & 0 \\ 0 & 6 \end{bmatrix}, \quad B = \begin{bmatrix} 0.2 & -3.5 \\ 0.1 & 0.3 \end{bmatrix}, \quad C = \begin{bmatrix} 0.4 & 0.2 \\ 0.1 & 0.7 \end{bmatrix}, \quad D = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix},$$

$$\Sigma_1 = \begin{bmatrix} 0 & 0.1 \\ 0.1 & -0.1 \end{bmatrix}, \quad \Sigma_2 = \begin{bmatrix} -0.1 & 0.1 \\ 0 & 0.1 \end{bmatrix}, \quad \Sigma_3 = \begin{bmatrix} 0.1 & 0.1 \\ 0.1 & 0 \end{bmatrix}, \quad H = \begin{bmatrix} 0.01 \\ 0.01 \end{bmatrix},$$

$$G_1 = \begin{bmatrix} 0.2 & 0 \\ 0 & -0.2 \end{bmatrix}, \quad G_2 = \begin{bmatrix} -0.1 & -0.2 \\ 0.2 & 0.1 \end{bmatrix}, \quad G_3 = \begin{bmatrix} 0.1 & -0.1 \\ 0.05 & 0.1 \end{bmatrix}, \quad G_4 = \begin{bmatrix} 0.1 & 0.2 \\ -0.1 & -0.1 \end{bmatrix}$$

$$I_k = \begin{bmatrix} 0.3 & 0 \\ 0 & 0.3 \end{bmatrix},$$

The activation functions are taken as follows:

$$g_1(x) = \frac{1}{20}(|x+1| + |x-1|), \quad g_2(x) = \frac{1}{10}(|x+1| + |x-1|).$$

It can be verified that Assumption (H1) is satisfied with  $F_1^- = -0.1$ ,  $F_1^+ = 0.1$ ,  $F_2^- = -0.2$ ,  $F_2^+ = 0.2$ . Thus

$$F_1 = \begin{bmatrix} -0.01 & 0 \\ 0 & -0.04 \end{bmatrix}, \quad F_2 = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix},$$

By solving the LMIs (54)–(58) in Theorem 4.1 using MATLAB LMI toolbox, one can obtain the feasible solution for any time delay satisfying  $0 < d(t) \leq 1.6241$  and  $0 \leq 1.0 \leq \tau(t) \leq 3.5783$  when  $\mu = 0.2, \delta = 0.1$ . Suppose, if we take  $\mu = 0.2, \delta = 0.1, \tau_1 = 1.0, d = 0.3, \tau_2 = 2.3$ , we can obtain the following feasible solutions

$$P_1 = \begin{bmatrix} 123.1307 & -2.6762 \\ -2.6762 & 126.2225 \end{bmatrix}, \quad P_2 = \begin{bmatrix} 47.1324 & -0.9150 \\ -0.9150 & 56.3039 \end{bmatrix}, \quad P_3 = 10^4 \times \begin{bmatrix} 0.8462 & 0.0301 \\ 0.0301 & 2.8263 \end{bmatrix},$$

$$\begin{aligned}
P_4 &= \begin{bmatrix} 14.1132 & 0.7140 \\ 0.7140 & 11.0384 \end{bmatrix}, & P_5 &= 10^3 \times \begin{bmatrix} 1.0766 & 0.0020 \\ 0.0020 & 0.5875 \end{bmatrix}, & L_1 &= \begin{bmatrix} 43.0610 & -1.6700 \\ -1.6700 & 43.8648 \end{bmatrix}, \\
L_2 &= \begin{bmatrix} 60.7614 & 18.9541 \\ 18.9541 & 60.9004 \end{bmatrix}, & L_3 &= \begin{bmatrix} 906.9030 & 97.3614 \\ 97.3614 & 484.7922 \end{bmatrix}, & R_1 &= \begin{bmatrix} 20.5262 & -1.3446 \\ -1.3446 & 25.1153 \end{bmatrix}, \\
R_2 &= \begin{bmatrix} 38.7288 & 6.8815 \\ 6.8815 & 32.1568 \end{bmatrix}, & R_3 &= \begin{bmatrix} 269.8072 & 46.8835 \\ 46.8835 & 156.1916 \end{bmatrix}, & S_1 &= \begin{bmatrix} 15.3430 & 0.2558 \\ 0.2558 & 13.4679 \end{bmatrix}, \\
S_2 &= \begin{bmatrix} 34.5891 & 7.1417 \\ 7.1417 & 30.1517 \end{bmatrix}, & S_3 &= \begin{bmatrix} 386.7042 & 41.7548 \\ 41.7548 & 221.4267 \end{bmatrix}, & Q_1 &= \begin{bmatrix} 11.9561 & -0.4032 \\ -0.4032 & 9.9872 \end{bmatrix}, \\
Q_2 &= \begin{bmatrix} 7.1335 & -0.2417 \\ -0.2417 & 5.953 \end{bmatrix}, & Q_3 &= \begin{bmatrix} 363.5368 & 29.8191 \\ 29.8191 & 171.1898 \end{bmatrix}, & Q_4 &= \begin{bmatrix} 274.7815 & 25.0485 \\ 25.0485 & 113.2372 \end{bmatrix}, \\
T_1 &= \begin{bmatrix} 0.3657 & -0.0004 \\ -0.0004 & 0.2838 \end{bmatrix}, & T_2 &= \begin{bmatrix} 0.4195 & -0.0008 \\ -0.0008 & 0.3240 \end{bmatrix}, & T_3 &= \begin{bmatrix} 1.3479 & -0.0040 \\ -0.0040 & 1.0312 \end{bmatrix}, \\
T_4 &= \begin{bmatrix} 0.0933 & -0.0002 \\ -0.0002 & 0.0760 \end{bmatrix}, & X_1 &= \begin{bmatrix} 11.9820 & -0.4040 \\ -0.4040 & 9.9964 \end{bmatrix}, & X_2 &= \begin{bmatrix} 17.7797 & -0.5943 \\ -0.5943 & 14.8716 \end{bmatrix}, \\
L &= 10^3 \times \begin{bmatrix} 2.3688 & 0 \\ 0 & 2.3437 \end{bmatrix}, & S &= \begin{bmatrix} 509.7439 & 0 \\ 0 & 216.5831 \end{bmatrix}, & M_1 &= \begin{bmatrix} -0.4444 & -0.0119 \\ -0.0119 & -0.2591 \end{bmatrix}, \\
M_2 &= \begin{bmatrix} 0.2137 & 0.4291 \\ 0.4291 & 0.3288 \end{bmatrix}, & M_3 &= \begin{bmatrix} 0.3304 & -0.0016 \\ -0.0016 & 0.2543 \end{bmatrix}, & M_4 &= \begin{bmatrix} -0.5125 & -0.3566 \\ -0.3566 & -0.6523 \end{bmatrix}, \\
M_5 &= \begin{bmatrix} 0.1783 & 0.1182 \\ 0.1182 & 0.0496 \end{bmatrix}, & M_6 &= \begin{bmatrix} -0.2253 & -0.0570 \\ -0.0570 & -0.0205 \end{bmatrix}, & M_7 &= \begin{bmatrix} 0.0125 & 0.0003 \\ 0.0003 & -0.0008 \end{bmatrix}, \\
M_8 &= \begin{bmatrix} 0.0022 & 0.0028 \\ 0.0028 & 0.0002 \end{bmatrix}, & N_1 &= \begin{bmatrix} -0.0753 & -0.0085 \\ -0.0085 & -0.0014 \end{bmatrix}, & N_2 &= \begin{bmatrix} 0.2967 & -0.0009 \\ -0.0009 & 0.2269 \end{bmatrix}, \\
N_3 &= \begin{bmatrix} -0.2987 & -0.0004 \\ -0.0004 & -0.2275 \end{bmatrix}, & N_4 &= \begin{bmatrix} -0.1970 & -0.1884 \\ -0.1884 & -0.3900 \end{bmatrix}, & N_5 &= \begin{bmatrix} 0.1288 & 0.0692 \\ 0.06920 & 0.0412 \end{bmatrix}, \\
N_6 &= \begin{bmatrix} -0.1526 & -0.0380 \\ -0.0380 & -0.0211 \end{bmatrix}, & N_7 &= \begin{bmatrix} -0.0014 & 0.0036 \\ 0.0036 & -0.0009 \end{bmatrix}, & N_8 &= \begin{bmatrix} 0.0015 & 0.0019 \\ 0.0019 & 0.0002 \end{bmatrix},
\end{aligned}$$

$$U_1 = \begin{bmatrix} -0.0153 & 0.0066 \\ 0.0066 & 0.0009 \end{bmatrix}, \quad U_2 = \begin{bmatrix} -0.2964 & 0.0009 \\ 0.0009 & -0.2268 \end{bmatrix}, \quad U_3 = \begin{bmatrix} -0.0896 & -0.2758 \\ -0.2758 & 0.8246 \end{bmatrix},$$

$$U_4 = \begin{bmatrix} 0.2944 & -0.0009 \\ -0.0009 & 0.2252 \end{bmatrix}, \quad U_5 = \begin{bmatrix} -0.0060 & 0.0274 \\ 0.0274 & -0.0932 \end{bmatrix}, \quad U_6 = \begin{bmatrix} -0.0210 & 0.0037 \\ 0.0037 & 0.0290 \end{bmatrix},$$

$$U_7 = \begin{bmatrix} 0.0015 & -0.0003 \\ -0.0003 & -0.0008 \end{bmatrix}, \quad U_8 = \begin{bmatrix} -0.0147 & -0.0026 \\ -0.0026 & -0.0112 \end{bmatrix}, \quad Z_1 = \begin{bmatrix} -1.6241 & 0.1507 \\ 0.1507 & -0.3396 \end{bmatrix},$$

$$Z_2 = \begin{bmatrix} 1.9428 & -0.0044 \\ -0.0044 & 1.4590 \end{bmatrix}, \quad \lambda_1 = 218.3413, \quad \lambda_2 = 254.9511, \quad \lambda_3 = 218.1895,$$

$$\gamma = 451.8541.$$

## 6. CONCLUSION

In this paper we have studied the passivity issue for a new class of impulsive stochastic neural networks with time delays in the leakage terms and mixed time delays are studied under two cases: with known or unknown parameters. In order to prove the passivity for the suggested system, many techniques such as Lyapunov stability theory, stochastic analysis and linear matrix inequalities techniques have been successfully used in this paper. Finally, numerical examples have been provided to demonstrate the validity of the approach. By utilizing the proposed idea of this paper, future works will focus on stabilization for various dynamic systems with time-delays such as switched generalized neural networks and Memristor-Based Recurrent Neural Networks, Complex valued neural networks, Chaotic Lur'e Systems. The corresponding results will appear in the near future.

(Received May 10, 2017)

## REFERENCES

- 
- [1] P. Balasubramaniam, G. Nagamani, and R. Rakkiyappan: Passivity analysis for neural networks of neutral type with Markovian jumping parameters and time delay in the leakage term. *Comm. Nonlinear Sci. Numerical Simul.* *16* (2011), 4422–4437. DOI:10.1016/j.cnsns.2011.03.028
  - [2] S. Boyd, L. Ghaoui, E. Feron, and V. Balakrishnan: *Linear Matrix Inequalities in System and Control Theory*. SIAM, Philadelphia 1994. DOI:10.1137/1.9781611970777
  - [3] J. Cao and R. Li: Fixed-time synchronization of delayed memristor-based recurrent neural networks. *Science China Inform. Sci.* *60* (2017), 032201. DOI:10.1007/s11432-016-0555-2
  - [4] J. Cao, R. Rakkiyappan, K. Maheswari, and A. Chandrasekar: Exponential  $H_\infty$  filtering analysis for discrete-time switched neural networks with random delays using sojourn probabilities. *Science China Inform. Sci.* *59*(2016), 3, 387–402. DOI:10.1007/s11431-016-6006-5



- [5] Y. Chen, H. Wang, A. Xue, and R. Lu: Passivity analysis of stochastic time-delay neural networks. *Nonlinear Dynamics* 61 (2010), 71–82. DOI:10.1007/s11071-009-9632-7
- [6] K. Gopalsamy: *Stability and Oscillations in Delay Differential Equations of Population Dynamics*. Kluwer Academic Publishers, Dordrecht 1992. DOI:10.1007/978-94-015-7920-9
- [7] S. Haykin: *Neural Networks: a Comprehensive Foundation* (revised ed.) Upper Saddle River, Prentice-Hall, NJ 1998.
- [8] M. Hu, J. Cao, and A. Hu: Exponential stability of discrete-time recurrent neural networks with time-varying delays in the leakage terms and linear fractional uncertainties. *IMA J. Math. Control Inform.* 31 (2014), 345–362. DOI:10.1093/imamci/dnt014
- [9] K. Gu: An integral inequality in the stability problem of time delay systems. In: *Proc. 39th IEEE Conference on Decision Control* 2000, pp. 2805–2810. DOI:10.1109/cdc.2000.914233
- [10] Y. He, Q. Wang, C. Lin, and M. Wu: Delay-range-dependent stability for systems with time-varying delay. *Automatica* 43 (2007), 371–376. DOI:10.1016/j.automatica.2006.08.015
- [11] O. Kwon, S. Lee and Ju H. Park: Improved delay-dependent exponential stability for uncertain stochastic neural networks with time-varying delays. *Physics Lett. A* 374 (2010), 1232–1241. DOI:10.1016/j.physleta.2010.01.007
- [12] O. Kwon, M. Park, Ju.H. Park, S. Lee, and E. Cha: Improved approaches to stability criteria for neural networks with time-varying delays. *J. Franklin Inst.* 350 (2013), 2710–2735. DOI:10.1016/j.jfranklin.2013.06.014
- [13] X. Li and J. Cao: Delay-dependent stability of neural networks of neutral typewith time delay in the leakage term. *Nonlinearity* 23 (2010), 1709–1726. DOI:10.1088/0951-7715/23/7/010
- [14] R. Li and J. Cao: Dissipativity analysis of memristive neural networks with time-varying delays and randomly occurring uncertainties. *Math. Methods Appl. Sci.* 39 (2016), 11, 2896–2915. DOI:10.1002/mma.3738
- [15] R. Li and J. Cao: Stability analysis of reaction-diffusion uncertain memristive neural networks with time-varying delays and leakage term. *Appl. Math. Comput.* 278 (2016), 54–69. DOI:10.1016/j.amc.2016.01.016
- [16] X. Li and X. Fu: Effect of leakage time-varying delay on stability of nonlinear differential systems. *J. Franklin Inst.* 350 (2013), 1335–1344. DOI:10.1016/j.jfranklin.2012.04.007
- [17] X. Li and R. Rakkiyappan: Stability results for Takagi–Sugeno fuzzy uncertain BAM neural networks with time delays in the leakage term. *Neural Computing Appl.* 22 (2013), S203–S219. DOI:10.1007/s00521-012-0839-z
- [18] X. Li and S. Song: Impulsive control for existence, uniqueness and global stability of periodic solutions of recurrent neural networks with discrete and continuously distributed delays. *IEEE Trans. Neural Networks Learning Systems* 24 (2013), 868–877. DOI:10.1109/tnnls.2012.2236352
- [19] X. Li and S. Song: Stabilization of delay systems: Delay-dependent impulsive control. *IEEE Trans. Automat. Control* 62 (2017), 406–411. DOI:10.1109/tac.2016.2530041
- [20] H. Li, C. Wang, P. Shi, and H. Gao: New passivity results for uncertain discrete-time stochastic neural networks with mixed time delays. *Neurocomputing* 73 (2010), 3291–3299. DOI:10.1016/j.neucom.2010.04.019
- [21] X. Li and J. Wu: Stability of nonlinear differential systems with state-dependent delayed impulses. *Automatica* 64 (2016), 63–69. DOI:10.1016/j.automatica.2015.10.002

- [22] Y. Li, L. Yang, and L. Sun: Existence and exponential stability of an equilibrium point for fuzzy BAM neural networks with time-varying delays in leakage terms on time scales. *Advances Diff. Equations* 2013 (2013), 218. DOI:10.1186/1687-1847-2013-218
- [23] Y. Liu, Z.D.Wang, and X.H. Liu: Global exponential stability of generalized recurrent neural networks with discrete and distributed delays. *Neural Networks* 19 (2006), 5, 667–675. DOI:10.1016/j.neunet.2005.03.015
- [24] X. Mao: *Stochastic Differential Equations with their Applications*. Horwood, Chichester 1997.
- [25] A. Michel and D. Liu: *Qualitative Analysis and Synthesis of Recurrent Neural Networks*. Marcel Dekker, New York 2002.
- [26] L. Pan and J. Cao: Robust stability for uncertain stochastic neural network with delay and impulses. *Neurocomputing* 94 (2012), 102–110. DOI:10.1016/j.neucom.2012.04.013
- [27] R. Raja, U. Karthik Raja, R. Samidurai, and A. Leelamani: Dissipativity of discrete-time BAM stochastic neural networks with Markovian switching and impulses. *J. Franklin Inst.* 350 (2013), 3217–3247. DOI:10.1016/j.jfranklin.2013.08.003
- [28] R. Raja, U. Karthik Raja, R. Samidurai, and A. Leelamani: Passivity analysis for uncertain discrete time stochastic BAM neural networks with time-varying delays. *Neural Computing Appl.* 25 (2014), 751–766. DOI:10.1007/s00521-014-1545-9
- [29] R. Raja and R. Samidurai: New delaydependent robust asymptotic stability for uncertain stochastic recurrent neural networks with multiple time varying delays. *J. Franklin Inst.* 349 (2012), 2108–2123. DOI:10.1016/j.jfranklin.2012.03.007
- [30] J. Rubio: Interpolation neural network model of a manufactured wind turbine. *Neural Computing Appl.* 28 (2017), 2017–2028. DOI:10.1007/s00521-015-2169-4
- [31] Q. Song and J. Cao: Passivity of uncertain neural networks with both leakage delay and time-varying delay. *Nonlinear Dynamics* 67 (2012), 1695–1707. DOI:10.1007/s11071-011-0097-0
- [32] Q. Song and J. Cao: Passivity of uncertain neural networks with both leakage delay and time-varying delay. *Nonlinear Dynamics* 67 (2012), 1695–1707. DOI:10.1007/s11071-011-0097-0
- [33] Z. Tu, J. Cao, A. Alsaedi, and T Hayat: Global dissipativity analysis for delayed quaternion-valued neural networks. *Neural Networks* 89 (2017), 97–104. DOI:10.1016/j.neunet.2017.01.006
- [34] Z. Wu, Ju H. Park, H. Su, and J. Chu: New results on exponential passivity of neural networks with time-varying delays. *Nonlinear Analysis: Real World Appl.* 13 (2012), 1593–1599. DOI:10.1016/j.nonrwa.2011.11.017
- [35] C. Yang and T. Huang: Improved stability criteria for a class of neural networks with variable delays and impulsive perturbations. *Appl. Math. Comput.* 243 (2014), 923–935. DOI:10.1016/j.amc.2014.06.045
- [36] Z. Zhao, Q. Song, and S. He: Passivity analysis of stochastic neural networks with time-varying delays and leakage delay. *Neurocomputing* 47 (2015), 1–10. DOI:10.1016/j.neucom.2012.08.049
- [37] C. Zheng, C. Gong, and Z. Wang: New passivity conditions with fewer slack variables for uncertain neural networks with mixed delays. *Neurocomputing* 118 (2013), 237–244. DOI:10.1016/j.neucom.2013.02.032

*Senthil Raj, Department of Mathematics, Thiruvalluvar University, Vellore 632 115. India.*

*e-mail: senthilraj106@gmail.com*

*Raja Ramachandran, Ramanujan Centre for Higher Mathematics, Alagappa University, Karaikudi 630 004. India.*

*e-mail: rajarchm2012@gmail.com*

*Samidurai Rajendiran, Department of Mathematics, Thiruvalluvar University, Vellore 632 115. India.*

*e-mail: samidurair@gmail.com*

*Jinde Cao, School of Mathematics, Southeast University, Nanjing 211 189. P. R. China.*

*e-mail: jdcao@seu.edu.cn*

*Xiaodi Li, School of Mathematics and Statistics, Shandong Normal University, Ji'nan, 250014. P. R. China.*

*e-mail: sodymath@163.com*