

EPI-CONVERGENCE IN DISTRIBUTION OF NORMAL INTEGRANDS WITH APPLICATIONS TO SETS OF ϵ -OPTIMAL SOLUTIONS

DIETMAR FERGER

We derive necessary and sufficient conditions for epi-convergence in distribution of normal integrands. As a basic tool for the proof a new characterisation for distributional convergence of random closed sets is used. Our approach via the epi-topology allows us to show that, if a net of normal integrands epi-converges in distribution, then the pertaining sets of ϵ -optimal solutions converge in distribution in the underlying hyperspace endowed with the upper Fell topology. Under some boundedness and uniqueness assumptions the convergence even holds for the Fell topology. Finally, measurable selections converge weakly to a Choquet-capacity.

Keywords: weak convergence, epi-topology, hyperspaces, Fell topologies, random closed sets, capacity functionals

Classification: 60B05, 60B10, 26E25

1. INTRODUCTION

Let (E, \mathcal{G}) be a locally compact second countable Hausdorff-space (lcsH) with \mathcal{F} and \mathcal{K} the pertaining families of all closed sets and all compact sets, respectively. E is called the *carrier space*. We want to equip \mathcal{F} with some topology. For that purpose introduce for every subset $A \subseteq E$ the systems $\mathcal{M}(A) := \{F \in \mathcal{F} : F \cap A = \emptyset\}$ of all *missing sets* and $\mathcal{H}(A) := \{F \in \mathcal{F} : F \cap A \neq \emptyset\}$ of all *hitting sets* of A . Put

$$\mathcal{S} := \{\mathcal{M}(K) : K \in \mathcal{K}\} \cup \{\mathcal{H}(G) : G \in \mathcal{G}\} \subseteq 2^{\mathcal{F}}.$$

Then the topology on \mathcal{F} generated by \mathcal{S} is called *Fell topology* and denoted by τ_F . It goes back to J. Fell [4]. Convergence in the Fell topology is the same as convergence in the sense of Painlevé–Kuratowski, confer Theorem C.7 in Molchanov [13]. It induces the Borel- σ -algebra $\mathcal{B}_F := \sigma(\tau_F)$, the smallest σ -algebra on \mathcal{F} containing the Fell topology. If $(\Omega, \mathcal{A}, \mathbb{P})$ is some probability space, then by definition a *random closed set* (in E on Ω) is a map $C : \Omega \rightarrow \mathcal{F}$, which is $\mathcal{A} - \mathcal{B}_F$ measurable. Occasionally, we will write $\mathcal{F} \equiv \mathcal{F}(E)$ and $\tau_F \equiv \tau_F(E)$ in order to underline the basic carrier space E .

Besides random closed sets we will investigate so-called *normal integrands*. To explain this notion first consider the collection $S \equiv S(E)$ of all lower semicontinuous (lsc)

functions $f : E \rightarrow \overline{\mathbb{R}}$, where $\overline{\mathbb{R}} = [-\infty, \infty]$ is the extended real line:

$$S := \{f : E \rightarrow \overline{\mathbb{R}}; f \text{ lsc}\}.$$

Similarly as for \mathcal{F} the function space S will be endowed with a topology. With this in mind consider for each $A \subseteq E$ the functional $I_A : S \rightarrow \overline{\mathbb{R}}$ defined by

$$I_A(f) := \inf_{x \in A} f(x).$$

Let the *epi-topology* τ_e be the coarsest topology on S with respect to which I_K is lsc for all $K \in \mathcal{K}$ and I_G is upper semicontinuous (usc) for all $G \in \mathcal{G}$. Convergence in the epi-topology is equivalent to epi-convergence. This follows from Theorem 5.3.2 in Molchanov [13]. If $\mathcal{B}_e := \sigma(\tau_e)$ is the corresponding Borel- σ -algebra on S , then a mapping $Z : \Omega \rightarrow S$, that is $\mathcal{A} - \mathcal{B}_e$ measurable is called *normal integrand (on Ω)*. According to Lemma 2.5 of Ferger [6] our notion of normal integrand coincides with that of Molchanov [13] and Rockafellar and Wets [16].

Given a directed set (J, \leq) and a family of probability spaces $(\Omega_\alpha, \mathcal{A}_\alpha, \mathbb{P}_\alpha)$, $\alpha \in J$, we consider random closed sets C_α in E and normal integrands Z_α on Ω_α . In our paper we first give an equivalent condition for distributional convergence of a net $(C_\alpha)_{\alpha \in J}$ to a limit C :

$$C_\alpha \xrightarrow{\mathcal{D}} C \quad \text{in } (\mathcal{F}, \tau_F). \quad (1)$$

Notice that the convergence in (1) by definition is the same as convergence of the distributions $Q_\alpha := \mathbb{P}_\alpha \circ C_\alpha^{-1}$ to $\mathbb{P} \circ C^{-1} =: Q$ in the weak topology:

$$Q_\alpha \xrightarrow{w} Q \quad \text{on } (\mathcal{F}, \tau_F). \quad (2)$$

It follows from the definition of Topsøe [20] or Gänssler and Stute [10] that weak convergence (2) holds for instance if and only if

$$\liminf_{\alpha} Q_\alpha(\mathbf{O}) \geq Q(\mathbf{O}) \quad \forall \mathbf{O} \in \tau_F.$$

A short introduction of the weak topology and weak convergence of probability measures on arbitrary topological spaces can be found in Ferger [7]. Once the above announced criterion for (1) or (2), respectively, is available it will be used to describe distributional convergence of (Z_α) to Z , i. e.

$$Z_\alpha \xrightarrow{\mathcal{D}} Z \quad \text{in } (S, \tau_e)$$

or what is the same

$$P_\alpha \xrightarrow{w} P \quad \text{on } (S, \tau_e),$$

where $P = \mathbb{P} \circ Z^{-1}$ and $P_\alpha = \mathbb{P}_\alpha \circ Z_\alpha^{-1}$ are the distributions of Z and Z_α under \mathbb{P} and \mathbb{P}_α , respectively. This is possible, because there is a strong relation between the two spaces (\mathcal{F}, τ_F) and (S, τ_e) , which can be expressed by *epigraphs*. Here, for any function $f : E \rightarrow \overline{\mathbb{R}}$ the *epigraph* of f is the set

$$\text{epi}(f) := \{(x, a) \in E \times \mathbb{R} : f(x) \leq a\}.$$

It is well-known that the function f is lsc if and only if $\text{epi}(f)$ is a closed subset of $E \times \mathbb{R}$ endowed with the product-topology. Let $\mathcal{E} = \{\text{epi}(f) : f \in S\} \subseteq \mathcal{F}(E \times \mathbb{R})$ be the family of all epigraphs and σ be the subspace topology on \mathcal{E} . Then Attouch [1], p.254-255, shows that (\mathcal{E}, σ) is compact and that the map

$$\phi : (S, \tau_e) \rightarrow (\mathcal{E}, \sigma) \text{ defined by } \phi(f) = \text{epi}(f), \quad (3)$$

is a homeomorphism.

The paper is organized as follows: In the next section we derive a new necessary and sufficient condition for weak convergence of probability measures on the hyperspace \mathcal{F} equipped with the Fell topology. This result is then used in section 3 to find a new equivalent characterization for weak convergence of probability measures on the space S of all lower semicontinuous functions endowed with the epi-topology. In section 4 we show that epi-convergence in distribution of normal integrands entails the convergence of the corresponding ϵ -optimal solutions as random closed sets. In particular, it follows that single solutions converge in distribution to the almost sure unique minimizing point of the limit normal integrand. If uniqueness is not given, then the solutions converge to the entire set, say C , of all minimizers. The latter means that the distributions converge weakly to the capacity-functional of C .

2. WEAK CONVERGENCE OF PROBABILITY MEASURES ON (\mathcal{F}, τ_F)

In this section we give a necessary and sufficient condition for $Q_\alpha \rightarrow_w Q$ on (\mathcal{F}, τ_F) . As (E, \mathcal{G}) is lcscH it is metrisable. By Theorem 2 of Vaughan [21] there exists an equivalent metric d such that every bounded set is relatively compact. Further, by second countability there exists a countable and dense subset $E_0 \subseteq E$. For a general subset A of E , A^0, \overline{A} and ∂A denote the interior, the closure and the boundary of A . Let $B_d(x, r) \equiv B(x, r)$ and $\overline{B}_d(x, r) \equiv \overline{B}(x, r)$ be the open and closed ball with center at $x \in E$ and radius $r \in \mathbb{R}$. (Observe that e.g. $\overline{B}_d(x, r) = \emptyset$ for $r < 0$.) Notice that every closed ball is bounded and therefore is compact. Given a subset $D \subseteq [0, \infty)$ we introduce the family

$$\mathcal{U}(D) := \left\{ \bigcup_{i=1}^m \overline{B}(x_i, r_i) : m \in \mathbb{N}, x_i \in E_0, r_i \in D, 1 \leq i \leq m \right\}. \quad (4)$$

If we agree to say that a closed ball with center in E_0 and radius in D is called a *closed D-ball*, then $\mathcal{U}(D)$ is the set of all finite unions of closed D -balls. Moreover, let

$$\mathcal{K}_Q := \{K \in \mathcal{K} : Q(\partial_F \mathcal{H}(K)) = 0\} = \{K \in \mathcal{K} : Q(\mathcal{H}(K)) = Q(\mathcal{H}(K^0))\}, \quad (5)$$

where $\partial_F \mathbf{A}$ gives the boundary of a set $\mathbf{A} \subseteq \mathcal{F}$ with respect to the Fell topology. Here, the second equality follows from Lemma 4.3 in Ferger [7]. Deviating from the usual naming convention, we call $K \in \mathcal{K}_Q$ a *Q-continuity set*.

Next, we specify the set D . For that purpose consider

$$R_\pm(x, r) := \{s > 0 : \overline{B}(x, r \pm s) \text{ is a } Q\text{-continuity set}\}, r > 0.$$

It follows from (3.3) of Ferger [8] that the sets $\mathcal{H}(\overline{B}(x, r+s), s > 0)$, are pairwise disjoint, whence the complement $R_+(x, r)^c = \{s > 0 : Q(\partial_F \mathcal{H}(\overline{B}(x, r+s))) > 0\}$ is denumerable. Similarly, $R_-(x, r)^c$ is denumerable as well. (Here, notice that $\mathcal{H}(\overline{B}(x, r-s)) = \mathcal{H}(\emptyset) = \emptyset$. Therefore $\overline{B}(x, r-s)$ is a Q -continuity set for all $s > r$.) Thus the set

$$\bigcup_{x \in E_0, r \in \mathbb{Q}_+} R_+(x, r)^c \cup \bigcup_{x \in E_0, r \in \mathbb{Q}_+} R_-(x, r)^c$$

is still denumerable and so

$$R := \bigcap_{x \in E_0, r \in \mathbb{Q}_+} R_+(x, r) \cap \bigcap_{x \in E_0, r \in \mathbb{Q}_+} R_-(x, r)$$

lies dense in $[0, \infty)$. (Here \mathbb{Q}_+ denotes the set of all positive rational numbers.) In particular, there exists a sequence $(s_k)_{k \in \mathbb{N}}$ in R such that $s_k \downarrow 0$. Finally, we define

$$D := \{r + s_k : r \in \mathbb{Q}_+, k \in \mathbb{N}\} \cup \{r - s_k : r \in \mathbb{Q}_+, k \in \mathbb{N}, k \geq n_r\},$$

where $n_r \in \mathbb{N}$ is such that $r - s_k > 0$ for all $n \geq n_r$. Notice, that $D = D(E_0, Q)$ depends on E_0 and Q . It is countable and lies dense in $[0, \infty)$, i. e. $[0, \infty) \subseteq \overline{D}$. By construction, every closed D -ball is a Q -continuity set.

Theorem 2.1. The following two statements are equivalent:

$$(1) \quad Q_\alpha \longrightarrow_w Q \quad \text{on } (\mathcal{F}, \tau_F).$$

$$(2)$$

$$\lim_\alpha Q_\alpha(\mathcal{H}(U)) = Q(\mathcal{H}(U)) \quad \text{for all } U \in \mathcal{U}(D). \quad (6)$$

Proof. Assume that (2) holds. Let

$$\mathcal{D} := \{B(x, r) : x \in E_0, r \in \mathbb{Q}_+\} \quad \text{and} \quad \overline{\mathcal{D}} := \{\overline{B}(x, r) : x \in E_0, r \in \mathbb{Q}_+\}.$$

Then the family

$$\left\{ \mathbf{B} = \bigcap_{i=1}^m \mathcal{M}(C_i) \cap \bigcap_{j=1}^k \mathcal{H}(D_j) : m \in \mathbb{N}, k \in \mathbb{N}_0, C_1, \dots, C_m \in \overline{\mathcal{D}}, D_1, \dots, D_k \in \mathcal{D} \right\} \quad (7)$$

is known to be a countable base for τ_F , confer Schneider and Weil [19] or Ferger [5]. Consequently:

$$\text{Every open } \mathbf{O} \in \tau_F \text{ is a countable union of such base-sets } \mathbf{B}. \quad (8)$$

We will show that in turn every base-set \mathbf{B} can be represented as a countable union as follows:

$$\mathbf{B} = \bigcup_{k, l} \mathcal{M}(U_k) \cap \mathcal{H}(B_{l_1}) \cap \dots \cap \mathcal{H}(B_{l_k}), \quad (9)$$

where the union extends over all $\underline{k} = (k_1, \dots, k_m) \in \mathbb{N}^m$ and all $\underline{l} = (l_1, \dots, l_k) \in \mathbb{N}^k$. Moreover, all involved sets $U_{\underline{k}}$ are elements of $\mathcal{U}(D)$ and $B_{l_1}, \dots, B_{l_k} \in \mathcal{U}(D)$ are actually single closed D -balls. For the proof of (9) consider a general set \mathbf{B} from the countable base (7):

$$\mathbf{B} = \bigcap_{i=1}^m \mathcal{M}(C_i) \cap \bigcap_{j=1}^k \mathcal{H}(D_j). \quad (10)$$

Here, for each $1 \leq i \leq m$ there are $x_i \in E_0$ and $r_i \in \mathbb{Q}_+$ such that $C_i = \overline{B}(x_i, r_i)$. Then $C_i^{(k)} := \overline{B}(x_i, r_i + s_k)$ is a closed D -ball for every $k \in \mathbb{N}$. Furthermore, $C_i^{(k)} \downarrow C_i, k \rightarrow \infty$. By Lemma 4.5 in Ferger [7] we obtain that $\mathcal{H}(C_i) = \bigcap_{k \in \mathbb{N}} \mathcal{H}(C_i^{(k)})$, whence by complementation it follows that

$$\mathcal{M}(C_i) = \bigcup_{k \in \mathbb{N}} \mathcal{M}(C_i^{(k)}) \quad \text{for all } 1 \leq i \leq m. \quad (11)$$

For every $1 \leq j \leq k$ the sets D_j are equal to $B(z_j, t_j)$, where $z_j \in E_0$ and $t_j \in \mathbb{Q}_+$. Similarly as above $D_j^{(l)} = \overline{B}(z_j, t_j - s_l)$ is a closed D -ball for each $l \in \mathbb{N}$ (possibly empty, namely when $s_l > t_j$.) Clearly, $D_j = \bigcup_{l \in \mathbb{N}} D_j^{(l)}$ and therefore

$$\mathcal{H}(D_j) = \bigcup_{l \in \mathbb{N}} \mathcal{H}(D_j^{(l)}) \quad \text{for all } 1 \leq j \leq k. \quad (12)$$

If we substitute (11) and (12) into (10), then the distributive law for sets gives us:

$$\mathbf{B} = \bigcup_{\underline{k}, \underline{l}} \mathcal{M}(C_1^{(k_1)}) \cap \dots \cap \mathcal{M}(C_m^{(k_m)}) \cap \mathcal{H}(D_1^{(l_1)}) \cap \dots \cap \mathcal{H}(D_k^{(l_k)}). \quad (13)$$

Notice that $\mathcal{M}(C_1^{(k_1)}) \cap \dots \cap \mathcal{M}(C_m^{(k_m)}) = \mathcal{M}(\bigcup_{i=1}^m C_i^{(k_i)})$. So, putting $U_{\underline{k}} := \bigcup_{i=1}^m C_i^{(k_i)}$ and $B_{l_j} := D_j^{(l_j)}, 1 \leq j \leq k$, gives the representation (9). Moreover, conclude that $U_{\underline{k}} \in \mathcal{U}(D)$ for all $\underline{k} \in \mathbb{N}^m$ and recall that B_{l_1}, \dots, B_{l_k} are closed D -balls for all $\underline{l} \in \mathbb{N}^k$ as announced.

Introduce the family

$$\underline{\mathcal{C}} := \{\mathcal{M}(U) \cap \mathcal{H}(B_1) \cap \dots \cap \mathcal{H}(B_k) : U \in \mathcal{U}(D), k \in \mathbb{N}_0, B_1, \dots, B_k \text{ are closed } D\text{-balls}\}.$$

It follows from (8) and (9) that every open set $\mathbf{O} \in \tau_F$ is a countable union of sets from $\underline{\mathcal{C}}$. One easily verifies that $\underline{\mathcal{C}} \subseteq \underline{\mathcal{B}}_F$ is a π -system, i. e. closed under finite intersections. Moreover,

$$Q_\alpha(\mathbf{C}) \rightarrow Q(\mathbf{C}) \quad \text{for all } \mathbf{C} \in \underline{\mathcal{C}}. \quad (14)$$

We prove this by induction on $k \in \mathbb{N}_0$. For $k = 0$ the sets in $\underline{\mathcal{C}}$ simplify to $\mathbf{C} = \mathcal{M}(U)$. Thus the convergence in (14) holds as a consequence of assumption (2) and the complementation rule for probabilities. Next, consider $k \geq 1$. Check that

$$\begin{aligned} & \mathcal{M}(U) \cap \mathcal{H}(B_1) \cap \dots \cap \mathcal{H}(B_k) \\ &= [\mathcal{M}(U) \cap \mathcal{H}(B_1) \cap \dots \cap \mathcal{H}(B_{k-1})] \setminus [\mathcal{M}(U \cup B_k) \cap \mathcal{H}(B_1) \cap \dots \cap \mathcal{H}(B_{k-1})]. \end{aligned}$$

Therefore, (14) follows from the induction hypothesis, because clearly $U \cup B_k \in \mathcal{U}(D)$.

Now, as in the proof of Theorem 2.2 in Billingsley [3] we can deduce that $\liminf_{\alpha} Q_{\alpha}(\mathbf{O}) \geq Q(\mathbf{O})$ for all open $\mathbf{O} \in \tau_F$, which as we know is equivalent to (1).

Finally, assume that (1) holds. Then

$$\lim_{\alpha} Q_{\alpha}(\mathcal{H}(K)) = Q(\mathcal{H}(K)) \quad \text{for all } K \in \mathcal{K}_Q$$

by Theorem 1.7.7 of Molchanov [13]. But $(\star) \mathcal{U}(D) \subseteq \mathcal{K}_Q$, because, as we have seen, every closed ball is compact and so is each finite union of them. Moreover, for closed D -balls B_1, \dots, B_m it follows that

$$0 \leq Q(\partial_F \mathcal{H}(\cup_{i=1}^m B_i)) = Q(\partial_F \cup_{i=1}^m \mathcal{H}(B_i)) \leq Q(\cup_{i=1}^m \partial_F \mathcal{H}(B_i)) \leq \sum_{i=1}^m Q(\partial_F \mathcal{H}(B_i)) = 0,$$

whence $\cup_{i=1}^m B_i \in \mathcal{K}_Q$ and thus (\star) holds. This shows the validity of (2). \square

If $(P_{\alpha})_{\alpha \in \mathbb{N}}$ is a sequence, then there are comparable results in the literature, where $\mathcal{U}(D)$ is replaced by another family, say \mathcal{V} . In Kallenberg [12] and Norberg [14] \mathcal{V} is a *separating class*, confer Molchanov [13] for its definition. In Salinetti and Wets [17] E is a finite dimensional linear space and $\mathcal{V} = \mathcal{U}(\mathbb{Q}_+) \cap \mathcal{K}_Q$ and in Pfugl [15] $E = \mathbb{R}^d$ and \mathcal{V} is equal to the family of all finite unions of compact rectangles with rational endpoints. Here, as it is the case in Salinetti and Wets [17] the unions (and not the single rectangles) must be Q -continuity sets. A major advantage of our "convergence determining class" $\mathcal{V} = \mathcal{U}(D)$ is that it is tailor-made for describing epi-convergence in distribution. We will see that this is so, because in our result the single closed balls are Q -continuity sets.

A reformulation of Theorem 2.1 in terms of random closed sets as in (1) reads as follows: $C_{\alpha} \xrightarrow{\mathcal{D}} C$ in (\mathcal{F}, τ_F) if and only if

$$\lim_{\alpha} \mathbb{P}_{\alpha}(C_{\alpha} \cap U \neq \emptyset) = \mathbb{P}(C \cap U \neq \emptyset) \text{ for all } U \in \mathcal{U}(D).$$

Here, by (5) every closed D -ball B satisfies $\mathbb{P}(C \cap B \neq \emptyset) = \mathbb{P}(C \cap B^0 \neq \emptyset)$.

3. WEAK CONVERGENCE OF PROBABILITY MEASURES ON S EQUIPPED WITH THE EPI-TOPOLOGY

We begin with several rather simple necessary conditions for weak convergence, which will be of good use later on.

Lemma 3.1. If $P_{\alpha} \rightarrow_w P$ in (S, τ_e) , then the following statements hold:

$$\liminf_{\alpha} P_{\alpha}(\cap_{j=1}^m \{I_{G_j} < a_j\}) \geq P(\cap_{j=1}^m \{I_{G_j} < a_j\}), \quad (15)$$

$$\liminf_{\alpha} P_{\alpha}(\cap_{j=1}^m \{I_{K_j} > a_j\}) \geq P(\cap_{j=1}^m \{I_{K_j} > a_j\}), \quad (16)$$

$$\limsup_{\alpha} P_{\alpha}(\cap_{j=1}^m \{I_{K_j} \leq a_j\}) \geq P(\cap_{j=1}^m \{I_{K_j} \leq a_j\}), \quad (17)$$

$$\limsup_{\alpha} P_{\alpha}(\cap_{j=1}^m \{I_{G_j} \geq a_j\}) \geq P(\cap_{j=1}^m \{I_{G_j} \geq a_j\}) \quad (18)$$

for all $m \in \mathbb{N}, G_1, \dots, G_m \in \mathcal{G}, K_1, \dots, K_m \in \mathcal{K}$ and $a_1, \dots, a_m \in \mathbb{R}$.

Proof. Since $\bigcap_{j=1}^m \{I_{G_j} < a_j\} \in \tau_e$, and $\bigcap_{j=1}^m \{I_{K_j} > a_j\} \in \tau_e$, the first two assertions (15) and (16) follow from the Portmanteau-Theorem. Similarly, as $\bigcap_{j=1}^m \{I_{K_j} \leq a_j\}$ and $\bigcap_{j=1}^m \{I_{G_j} \geq a_j\}$ both are τ_e -closed another application of the Portmanteau-Theorem yields (17) and (18) \square

A combination of (15) – (18) leads to a result involving usual limits.

Proposition 3.2. Weak convergence $P_\alpha \rightarrow_w P$ in (S, τ_e) entails

$$\lim_{\alpha} P_\alpha(\bigcap_{j=1}^m \{I_{K_j} \leq a_j\}) = P(\bigcap_{j=1}^m \{I_{K_j} \leq a_j\}), \quad (19)$$

$$\lim_{\alpha} P_\alpha(\bigcap_{j=1}^m \{I_{K_j} > a_j\}) = P(\bigcap_{j=1}^m \{I_{K_j} > a_j\}). \quad (20)$$

Here, (19) and (20), respectively, hold for all $m \in \mathbb{N}, a_1, \dots, a_m \in \mathbb{R}$ and $K_1, \dots, K_m \in \mathcal{K}$ satisfying the continuity condition

$$P(I_{K_j} \leq a_j) = P(I_{K_j^0} < a_j), \quad j = 1, \dots, m.$$

Proof. As to the proof of (19) observe that

$$\begin{aligned} P(\bigcap_{j=1}^m \{I_{K_j^0} < a_j\}) &\leq \liminf_{\alpha} P_\alpha(\bigcap_{j=1}^m \{I_{K_j^0} < a_j\}) \quad \text{by (15)} \\ &\leq \liminf_{\alpha} P_\alpha(\bigcap_{j=1}^m \{I_{K_j} \leq a_j\}) \\ &\qquad \text{since } I_{K_j^0} \geq I_{K_j} \text{ and } (-\infty, a_j) \subseteq (-\infty, a_j] \\ &\leq \limsup_{\alpha} P_\alpha(\bigcap_{j=1}^m \{I_{K_j} \leq a_j\}) \\ &\leq P(\bigcap_{j=1}^m \{I_{K_j} \leq a_j\}) \quad \text{by (17)} \\ &= P(\bigcap_{j=1}^m \{I_{K_j^0} < a_j\}) \quad \text{see below.} \end{aligned}$$

So, the assertion (19) follows, once we have shown the last equation. For this purpose, let $M_j := \{I_{K_j^0} < a_j\}$ and $N_j := \{I_{K_j} \leq a_j\}, j = 1, \dots, m$. Then $M_j \subseteq N_j$ for all j , because $I_{K_j^0} \geq I_{K_j}$ and $(-\infty, a_j) \subseteq (\infty, a_j]$. As a consequence, $M := \bigcap_{j=1}^m M_j \subseteq \bigcap_{j=1}^m N_j =: N$ and thus:

$$\begin{aligned} 0 &\leq Q(N) - Q(M) = Q(N \setminus M) = Q(N \cap (\bigcup_{j=1}^m M_j^c)) = Q(\bigcup_{j=1}^m (N \cap M_j^c)) \\ &\leq Q(\bigcup_{j=1}^m (N_j \cap M_j^c)) \leq \sum_{j=1}^m Q(N_j \setminus M_j) = \sum_{j=1}^m Q(N_j) - Q(M_j) = 0, \end{aligned}$$

where the last equality follows from our assumption. Consequently, $Q(N) = Q(M)$ as desired. Analogously, by using (16) and (18) one proves (20). \square

Proposition 3.2 gives two necessary conditions for weak convergence. Our next result yields two necessary and sufficient conditions.

Theorem 3.3. Let $D := D(E_0 \times \mathbb{Q}, P \circ \phi^{-1})$. Then the following statements (1)–(3) are equivalent:

- (1) $P_\alpha \rightarrow_w P$ in (S, τ_e)
- (2) $\lim_\alpha P_\alpha(\cap_{j=1}^m \{I_{\overline{B}(x_j, r_j)} \leq r_j + \alpha_j\}) = P(\cap_{j=1}^m \{I_{(\overline{B}(x_j, r_j))^0} \leq r_j + \alpha_j\})$.
- (3) $\lim_\alpha P_\alpha(\cap_{j=1}^m \{I_{\overline{B}(x_j, r_j)} > r_j + \alpha_j\}) = P(\cap_{j=1}^m \{I_{(\overline{B}(x_j, r_j))^0} > r_j + \alpha_j\})$.

Here, the equalities in (2) and (3), respectively, hold for all $m \in \mathbb{N}, x_1, \dots, x_m \in E_0, r_1, \dots, r_m \in D, \alpha_1, \dots, \alpha_m \in \mathbb{Q}$. (Notice that the closed balls $\overline{B}(x_j, r_j)$ satisfy the continuity condition (21) below.)

Proof. The necessity of (2) or (3), respectively, for weak convergence (1) follows from Proposition 3.2, because closed balls are compact and $r_j \in D$ means that $\overline{B}_{d \times u}((x_j, \alpha_j), r_j)$ is a $P \circ \phi^{-1}$ -continuity set, which by Lemma 5.1 in the appendix is equivalent to

$$P(I_{\overline{B}(x_j, r_j)} \leq r_j + \alpha_j) = P(I_{(\overline{B}(x_j, r_j))^0} < r_j + \alpha_j). \quad (21)$$

As to sufficiency we will see that it is enough to show that

$$P_\alpha \circ \phi^{-1} \rightarrow_w P \circ \phi^{-1} \quad \text{on } (\mathcal{F}(E \times \mathbb{R}), \tau_F(E \times \mathbb{R})). \quad (22)$$

So, we are dealing with $(E \times \mathbb{R}, d \times u)$ as carrier space instead of (E, d) . (The product-metric is specified in the Appendix below.) Assume that (2) holds. Observe that

$$\mathcal{U}(D) = \left\{ \bigcup_{i=1}^m \overline{B}_{d \times u}((x_i, \alpha_i), r_i) : m \in \mathbb{N}, (x_i, \alpha_i) \in E_0 \times \mathbb{Q}, r_i \in D, i = 1, \dots, m \right\}.$$

Let $U = \bigcup_{i=1}^m B_i \in \mathcal{U}(D)$, i. e. each B_i is equal to the closed ball $\overline{B}_{d \times u}((x_i, \alpha_i), r_i), i = 1, \dots, m$. Put $Q_\alpha := P_\alpha \circ \phi^{-1}$ and $Q := P \circ \phi^{-1}$. The *inclusion-exclusion formula* yields:

$$Q_\alpha(\mathcal{H}(U)) = Q_\alpha(\cup_{i=1}^m (\mathcal{H}(B_i))) = \sum_{k=1}^m (-1)^{k+1} \sum_{1 \leq i_1 < \dots < i_k \leq m} Q_\alpha(\cap_{s=1}^k \mathcal{H}(B_{i_s})). \quad (23)$$

For each summand on the right side of equation (23) it follows that:

$$\begin{aligned} Q_\alpha(\cap_{s=1}^k \mathcal{H}(B_{i_s})) &= P_\alpha(\cap_{s=1}^k \{f \in S : \text{epi}(f) \in \mathcal{H}(B_{i_s})\}) \\ &= P_\alpha(\cap_{s=1}^k \{f \in S : \text{epi}(f) \cap B_{i_s} \neq \emptyset\}) \\ &= P_\alpha(\cap_{s=1}^k \{I_{\overline{B}_d(x_{i_s}, r_{i_s})} \leq r_{i_s} + \alpha_{i_s}\}) \quad \text{by (36) and (38).} \end{aligned}$$

Consequently by (2),

$$Q_\alpha(\cap_{s=1}^k \mathcal{H}(B_{i_s})) \rightarrow P(\cap_{s=1}^k \{I_{\overline{B}_d(x_{i_s}, r_{i_s})} \leq r_{i_s} + \alpha_{i_s}\}) = Q(\cap_{s=1}^k \mathcal{H}(B_{i_s}))$$

upon noticing (36) and (38) again. With (23) we obtain that $Q_\alpha(\mathcal{H}(U)) \rightarrow Q(\mathcal{H}(U))$ for all $U \in \mathcal{U}(D)$. Thus the weak convergence (22) follows from Theorem 2.1. Deduce from (22) that

$$P_\alpha \circ \phi^{-1} \rightarrow_w P \circ \phi^{-1} \quad \text{on the subspace } (\mathcal{E}, \sigma).$$

This yields $P_\alpha \rightarrow_w P$ by the Continuous Mapping Theorem, because

$$P_\alpha = (P_\alpha \circ \phi^{-1}) \circ (\phi^{-1})^{-1}$$

and $\phi^{-1} : (\mathcal{E}, \sigma) \rightarrow (S, \tau_e)$ is continuous.

Finally, assume that (3) holds. Then

$$\begin{aligned} Q_\alpha(\mathcal{M}(U)) &= Q_\alpha(\mathcal{M}(\cup_{i=1}^m B_i)) = Q_\alpha(\cap_{i=1}^m \mathcal{M}(B_i)) \\ &= P_\alpha(\cap_{i=1}^m \{f \in S : \text{epi}(f) \in \mathcal{M}(B_i)\}) \\ &= P_\alpha(\cap_{i=1}^m \{f \in S : \text{epi}(f) \cap B_i = \emptyset\}) \\ &= P_\alpha(\cap_{i=1}^m \{I_{\overline{B}_d(x_i, r_i)} > r_i + \alpha_i\}) \quad \text{by (36) and (38).} \end{aligned}$$

So, weak convergence (22) follows from Theorem 2.1 by complementation, which as shown above results in (1). \square

The reformulation of our results in terms of normal integrands is obvious. For instance, a net (Z_α) of normal integrands epi-converges in distribution to a normal integrand Z if and only if

$$\mathbb{P}_\alpha\left(\inf_{t \in \overline{B}_d(x_i, r_i)} Z_\alpha(t) > r_i + \alpha_i, i = 1, \dots, m\right) \rightarrow \mathbb{P}\left(\inf_{t \in \overline{B}_d(x_i, r_i)} Z(t) > r_i + \alpha_i, i = 1, \dots, m\right)$$

for all $m \in \mathbb{N}, x_1, \dots, x_m \in E_0, r_1, \dots, r_m \in D, \alpha_1, \dots, \alpha_m \in \mathbb{Q}$. In this form we can immediately compare it with the equivalent characterisation of Molchanov's [13] Proposition 5.3.20:

$$\mathbb{P}_\alpha\left(\inf_{t \in K_i} Z_\alpha(t) > t_i, i = 1, \dots, m\right) \rightarrow \mathbb{P}\left(\inf_{t \in K_i} Z(t) > t_i, i = 1, \dots, m\right)$$

for all $m \in \mathbb{N}, t_1, \dots, t_m \in \mathbb{R}$ and K_1, \dots, K_m belonging to a separating class of subsets of E satisfying the condition

$$P(I_{K_i} \leq t_i) = P(I_{K_i^0} < t_i).$$

Examples for separating classes are the family \mathcal{K} or the family of all finite unions of closed balls with center in E_0 and positive rational radii. In both cases our countable class $\{\overline{B}_d(x, r) : x \in E_0, r \in D\}$ is significantly smaller. Similarly, the countable set $\{r + \alpha : r \in D, \alpha \in \mathbb{Q}\}$ is a subset of the real line \mathbb{R} . Finally, Molchanov only considers sequences $(Z_n)_{n \in \mathbb{N}}$ and not more generally nets $(Z_\alpha)_{\alpha \in J}$, as we do. In the case of sequences and $E = \mathbb{R}^d$ there are further characterisations for epi-convergence in distribution. In their Theorem 3.14 Salinetti and Wets [17] show that if (Z_n) is almost surely equi-lower semicontinuous, then epi-convergence in distribution is equivalent to convergence of the finite-dimensional distributions (fidis). Gersch [11], Theorems 2.19 and 2.25, requires merely *stochastically* equi-lower semicontinuity and gives a sufficient condition, which again involve the marginals of the Z_n , but in a more complicated way in comparison to the convergence of the fidis. However, if in addition the limit process Z is stochastically uniformly lower semicontinuous, then epi-convergence in distribution and convergence of the fidis are equivalent. This equivalence also holds for convex Z and $Z_n, n \in \mathbb{N}$ as

Ferger [6] shows in Theorem 2.12 and Proposition 2.13. In statistical applications the Skorokhod-space $(D(\mathbb{R}^d), s)$ plays an important role, because many empirical processes have trajectories lying in that function space. From Proposition 2.1 of Ferger [9] it follows that, if $Z_n \xrightarrow{D} Z$ in $(D(\mathbb{R}^d), s)$, then $\bar{Z}_n \xrightarrow{D} \bar{Z}$ in $(S(\mathbb{R}^d), e)$. Here, s denotes the Skorokhod-metric and \bar{f} is the lsc regularization of a function $f \in D(\mathbb{R}^d)$.

Remark 3.4. Since $\phi : (S, \tau_e) \rightarrow (\mathcal{E}, \sigma)$ is a homeomorphism, the Continuous Mapping Theorem yields that a net (Z_α) of normal integrands epi-converges in distribution to a normal integrand Z , $Z_\alpha \xrightarrow{D} Z$ in (S, τ_e) , if and only if the pertaining $\text{epi}(Z_\alpha)$ converge in distribution to $\text{epi}(Z)$ in the space $E \times \mathbb{R}$, i. e. $\text{epi}(Z_\alpha) \xrightarrow{D} \text{epi}(Z)$ in $(\mathcal{F}(E \times \mathbb{R}), \tau_F(E \times \mathbb{R}))$. In the literature so far this was taken as the definition of epi-convergence in distribution.

Remark 3.5. An important point in our investigations is that Vaughan's metric ensures that every closed bounded set is compact. This is fulfilled if E is a finite-dimensional normed linear space. Moreover, notice that in normed linear spaces $(\bar{B}(x, r))^0 = B(x, r)$. This need not be true more generally in metric spaces even if they are lcscH. As an example consider a countable set E with at least two elements endowed with the discrete metric. This space is lcscH, but $(\bar{B}(x, 1))^0 = E^0 = E \not\subseteq \{x\} = B(x, 1)$. If E is finite, then (E, d) is even compact and thus in particular d has the property of Vaughan's metric as actually every subset is compact.

4. APPLICATIONS TO SETS OF ϵ -OPTIMAL SOLUTIONS OF NORMAL INTEGRANDS

For $f \in S(E)$ and $\epsilon \geq 0$ let

$$A(f, \epsilon) := \{t \in E : f(t) \leq I_E(f) + \epsilon\}$$

be the set of all ϵ -optimal solutions of f . This set is closed, because $A(f, \epsilon) = \{f \leq \alpha\}$ with $\alpha := I_E(f) + \epsilon \in [-\infty, \infty]$. Now, if $I_E(f) \in \mathbb{R}$, then $\alpha \in \mathbb{R}$ and $A(f, \epsilon) = \{f \leq \alpha\} \in \mathcal{F}$ by lower semicontinuity of f . If $I_E(f) = \infty$, then $A(f, \epsilon) = \{f \leq \infty\} = E \in \mathcal{F}$. Finally, in case that $I_E(f) = -\infty$ then $A(f, \epsilon) = \{f = -\infty\} = \bigcap_{n \in \mathbb{N}} \{f \leq -n\} \in \mathcal{F}$ again by lower semicontinuity of f and since \mathcal{F} is closed under intersection. So, the assignment $(f, \epsilon) \mapsto A(f, \epsilon)$ defines a map

$$A : S \times \mathbb{R}_+ \rightarrow \mathcal{F}. \quad (24)$$

For $\epsilon = 0$ one obtains $A(f, 0) = \{t \in E : f(t) = I_E(f)\} = \text{Argmin}(f)$, the set of all minimizing points of the function f .

A very nice property of epi-convergence is formulated in our next result.

Proposition 4.1. Let $(f_n)_{n \in \mathbb{N}}$ be a sequence of lsc functions, which epi-converges to some $f \in S$, i. e. $f_n \rightarrow f$ in (S, τ_e) . Moreover, assume $(\epsilon_n)_{n \in \mathbb{N}} \geq 0$ is a sequence of non-negative numbers such that $\limsup_{n \rightarrow \infty} \epsilon_n \leq \epsilon$. Then

$$\text{PK-} \limsup_{n \rightarrow \infty} A(f_n, \epsilon_n) \subseteq A(f, \epsilon). \quad (25)$$

Here, PK- \limsup denotes the *Painlevé–Kuratowski outer limit* of a sequence of sets.

Proof. Let $t \in \text{PK-}\limsup_{n \rightarrow \infty} A(f_n, \epsilon_n)$. By definition of the outer limit there exists a subsequence $(n_j)_{j \in \mathbb{N}}$ of the natural numbers and points $t_{n_j} \in A(f_{n_j}, \epsilon_{n_j})$ with $t_{n_j} \rightarrow t$ as $j \rightarrow \infty$. Assume that $t \notin A(f, \epsilon)$. Then there exists some $s \in E$ such that $f(t) > f(s) + \epsilon$, because otherwise $f(t) \leq f(s) + \epsilon$ for all $s \in E$, whence $I_E(f) \geq f(t) - \epsilon$ and thus $f(t) \leq I_E(f) + \epsilon$ in contradiction to $t \notin A(f, \epsilon)$. It follows from the characteristion of epi-convergence, confer Theorem 5.3.2 (ii) in Molchanov [13], that there exists some sequence (s_n) such that $s_n \rightarrow s$ and $f_n(s_n) \rightarrow f(s)$. Since $t_{n_j} \in A(f_{n_j}, \epsilon_{n_j})$, we have that $f_{n_j}(t_{n_j}) \leq I_E(f_{n_j}) + \epsilon_{n_j} \leq f_{n_j}(s_{n_j}) + \epsilon_{n_j}$ and so

$$f_{n_j}(s_{n_j}) \geq f_{n_j}(t_{n_j}) - \epsilon_{n_j} \quad \forall j \in \mathbb{N}. \quad (26)$$

Conclude that

$$\begin{aligned} f(t) &> f(s) + \epsilon \\ &= \lim_{n \rightarrow \infty} f_n(s_n) + \epsilon \\ &= \liminf_{j \rightarrow \infty} f_{n_j}(s_{n_j}) + \epsilon \\ &\geq \liminf_{j \rightarrow \infty} (f_{n_j}(t_{n_j}) - \epsilon_{n_j}) + \epsilon && \text{by (26)} \\ &\geq \liminf_{j \rightarrow \infty} f_{n_j}(t_{n_j}) + \liminf_{j \rightarrow \infty} (-\epsilon_{n_j}) + \epsilon \\ &= \liminf_{j \rightarrow \infty} f_{n_j}(t_{n_j}) - \limsup_{j \rightarrow \infty} \epsilon_{n_j} + \epsilon \\ &\geq \liminf_{j \rightarrow \infty} f_{n_j}(t_{n_j}) - \limsup_{n \rightarrow \infty} \epsilon_n + \epsilon \\ &\geq \liminf_{j \rightarrow \infty} f_{n_j}(t_{n_j}) && \text{by assumption on } (\epsilon_n) \\ &\geq f(t), \end{aligned}$$

where the last inequality follows from Theorem 5.3.2 (ii) in Molchanov [13] since $(f_{n_j})_{j \in \mathbb{N}}$ as a subsequence epi-converges to f as well. So, we arrive at a contradiction, which finishes our proof. \square

Let $\mathcal{T} := \{[0, r) : 0 < r \in \mathbb{R}\} \cup \{\emptyset, \mathbb{R}_+\}$ be the *left-order topology* on $\mathbb{R}_+ := [0, \infty)$. Then the assumption $\limsup_{n \rightarrow \infty} \epsilon_n \leq \epsilon$ is equivalent to $\epsilon_n \rightarrow \epsilon$ in $(\mathbb{R}_+, \mathcal{T})$. If $\epsilon = 0$, then $\epsilon_n \rightarrow 0$ in the natural topology \mathcal{T}_n on \mathbb{R}_+ since all ϵ_n are non-negative and therefore $\liminf_{n \rightarrow \infty} \epsilon_n \geq 0$. For this special case we obtain Proposition 2.9 of Attouch [1], who however considers more generally arbitrary topological spaces (E, \mathcal{G}) .

Next, let τ_{uF} be the *upper Fell topology*, which is generated by the family $\mathcal{S}_{uF} := \{\mathcal{M}(K) : K \in \mathcal{K}\}$. In the literature τ_{uF} is also known under the name *miss topology*, confer Beer [2]. Since $\mathcal{S}_{uF} \subseteq \mathcal{S}$, the upper Fell topology is weaker than the Fell topology. In Lemma 2.2, Vogel [22] shows in case $E = \mathbb{R}^d$ that a sequence $(F_n)_{n \in \mathbb{N}} \subseteq \mathcal{F}$ of closed sets in E converges to some $F \in \mathcal{F}$ in the upper Fell topology, i. e.

$$F_n \rightarrow F \text{ in } (\mathcal{F}, \tau_{uF}) \text{ if and only if } \text{PK-}\limsup_{n \rightarrow \infty} F_n \subseteq F. \quad (27)$$

This remains valid more generally for (E, \mathcal{G}) lcsCH, confer Proposition 2.18 (b) in Ferger [5]. Our next result yields continuity of the map

$$A : S \times \mathbb{R}_+ \rightarrow \mathcal{F}.$$

Corollary 4.2. Let $\tau_e \times \mathcal{T}$ be the product-topology on $S \times \mathbb{R}_+$. Then

$$A : (S \times \mathbb{R}_+, \tau_e \times \mathcal{T}) \rightarrow (\mathcal{F}, \tau_{uF}) \quad \text{is continuous.} \quad (28)$$

In particular,

$$A : (S \times \mathbb{R}_+, \tau_e \times \mathcal{T}_n) \rightarrow (\mathcal{F}, \tau_{uF}) \quad \text{is continuous.} \quad (29)$$

Proof. By Corollary 2.79 of Attouch [1] the space (S, τ_e) is second-countable, which also holds for $(\mathbb{R}_+, \mathcal{T})$ as one easily verifies. Thus the product $(S \times \mathbb{R}_+, \tau_e \times \mathcal{T})$ is second-countable as well and in particular, it is first-countable. By Theorem 7.1.3 in Singh [18] it suffices to show that A is sequentially-continuous. So, assume that $(f_n, \epsilon_n) \rightarrow (f, \epsilon)$ in $(S \times \mathbb{R}_+, \tau_e \times \mathcal{T})$. This is equivalent to $f_n \rightarrow f$ in (S, τ_e) and $\epsilon_n \rightarrow \epsilon$ in $(\mathbb{R}_+, \mathcal{T})$. But the latter in turn means that $\limsup_{n \rightarrow \infty} \epsilon_n \leq \epsilon$. Therefore, the assertion in (28) follows from Proposition 4.1 in combination with the equivalence (27). Since $\tau_e \times \mathcal{T}_n \supseteq \tau_e \times \mathcal{T}$, the second assertion follows from the first one. \square

Let Z be a normal integrand and ϵ a random variable with values in \mathbb{R}_+ both defined on some measurable space (Ω, \mathcal{A}) . A first useful application of Corollary 4.2 yields measurability of the random set $A(Z, \epsilon)$. In the proof below we will use the following notation: Given a topological space (X, \mathcal{O}) the pertaining Borel- σ -algebra $\sigma(\mathcal{O})$ is denoted by $\mathcal{B}(X)$.

Corollary 4.3. If Z and ϵ are as above, then $A(Z, \epsilon)$ is a random closed set.

Proof. It follows from (29) that A is $\mathcal{B}(S \times \mathbb{R}_+) - \underline{\mathcal{B}}_{uF}$ measurable, where $\underline{\mathcal{B}}_{uF} := \sigma(\tau_{uF})$. But $\underline{\mathcal{B}}_{uF} = \underline{\mathcal{B}}_F$, because $\underline{\mathcal{B}}_F = \sigma(\mathcal{S}_{uF})$ by Lemma 2.1.1 in Schneider and Weil [19] and $\mathcal{S}_{uF} \subseteq \tau_{uF} \subseteq \tau_F$. Since $(S \times \mathbb{R}_+, \tau_e \times \mathcal{T}_n)$ is second-countable, it follows that the Borel- σ -algebra $\mathcal{B}(S \times \mathbb{R}_+) = \mathcal{B}(S) \otimes \mathcal{B}(\mathbb{R}_+)$. Infer that A is $\mathcal{B}(S) \otimes \mathcal{B}(\mathbb{R}_+) - \underline{\mathcal{B}}_F$ measurable. Deduce from our assumption that the product map $(Z, \epsilon) : (\Omega, \mathcal{A}) \rightarrow (S \times \mathbb{R}_+, \mathcal{B}(S) \otimes \mathcal{B}(\mathbb{R}_+))$ is measurable, whence the assertion follows upon noticing that $A(Z, \epsilon) = A \circ (Z, \epsilon)$ is a composition of measurable maps. \square

A further utility of Corollary 4.2 is that it enables us to apply the Continuous Mapping Theorem (CMT) for random variables in topological spaces, confer Proposition 8.4.16 in Gänssler and Stute [10]. The random variables Z, ϵ and $Z_\alpha, \epsilon_\alpha$ occurring in our results below are defined on $(\Omega, \mathcal{A}, \mathbb{P})$ and $(\Omega_\alpha, \mathcal{A}_\alpha, \mathbb{P}_\alpha)$, respectively.

Theorem 4.4. Let (Z_α) and (ϵ_α) be nets of normal integrands and non-negative random variables, respectively. Assume that

$$(Z_\alpha, \epsilon_\alpha) \xrightarrow{\mathcal{D}} (Z, \epsilon) \quad \text{in } (S \times \mathbb{R}_+, \tau_e \times \mathcal{T}). \quad (30)$$

Then $A(Z_\alpha, \epsilon_\alpha) \xrightarrow{\mathcal{D}} A(Z, \epsilon)$ in (\mathcal{F}, τ_{uF}) . This is the same as

$$\begin{aligned} & \limsup_\alpha \mathbb{P}_\alpha \left(\bigcap_{K \in \mathcal{K}^*} \{\omega \in \Omega_\alpha : A(Z_\alpha(\omega), \epsilon_\alpha(\omega)) \cap K \neq \emptyset\} \right) \\ & \leq \mathbb{P} \left(\bigcap_{K \in \mathcal{K}^*} \{\omega \in \Omega : A(Z(\omega), \epsilon(\omega)) \cap K \neq \emptyset\} \right) \end{aligned} \quad (31)$$

for every collection $\mathcal{K}^* \subseteq \mathcal{K}$ of compact sets in E .

Proof. Assumption (30) and the CMT applied to A in Corollary 4.2 gives the first assertion of the theorem. The second one follows from Proposition 2.1 of Ferger [7]. \square

A legitimate question is what are sufficient conditions for the validity of (30)? We provide a few answers in:

Remark 4.5. (1) Assume that there is a component-wise convergence, i. e.

$$Z_\alpha \xrightarrow{\mathcal{D}} Z \text{ in } (S, \tau_e) \text{ and } \epsilon_\alpha \xrightarrow{\mathcal{D}} \epsilon \text{ in } (\mathbb{R}_+, \mathcal{T}_n). \quad (32)$$

If ϵ is almost surely constant, then

$$(Z_\alpha, \epsilon_\alpha) \xrightarrow{\mathcal{D}} (Z, \epsilon) \text{ in } (S \times \mathbb{R}_+, \tau_e \times \mathcal{T}_n). \quad (33)$$

This follows from Slutsky's Theorem, confer Proposition 8.6.4 in Gänssler and Stute [10]. Now, (33) implies (30), because $\tau_e \times \mathcal{T}_n \supseteq \tau_e \times \mathcal{T}$.

(2) Suppose that Z_α and ϵ_α are \mathbb{P}_α -independent for each α . If component-wise convergence (32) holds, where Z and ϵ are \mathbb{P} -independent, then again (33) holds by Theorem 2.8 in Billingsley [3]. As we know this is enough for (30). A special case for this is when the net (ϵ_α) is deterministic and convergent with (deterministic) limit ϵ .

(3) It should be mentioned that Slutsky's Theorem and Theorem 2.8 in Billingsley [3] are formulated only for sequences. However, by using Proposition 8.4.9 in Gänssler and Stute [10] one can see that their proofs can easily be transferred to nets .

The question arises as to the requirements under which Fell-convergence in distribution is obtained. The answer involves the family $\mathbf{F}_{0,1} := \{\emptyset\} \cup \{\{x\} : x \in E\}$ of sets with at most one element.

Theorem 4.6. Assume that (30) holds with $\epsilon = 0$. If for every $\eta > 0$ there exists a compact $K \subseteq E$ such that

$$\liminf_\alpha \mathbb{P}_\alpha (\{\omega \in \Omega_\alpha : \emptyset \neq A(Z_\alpha(\omega), \epsilon_\alpha(\omega)) \subseteq K\}) \geq 1 - \eta$$

and if $\mathbb{P}(\{\omega \in \Omega : Z(\omega) \in \mathbf{F}_{0,1}\}) = 1$, i. e. Z has at most one minimizing point almost surely, then

$$A(Z_\alpha, \epsilon_\alpha) \xrightarrow{\mathcal{D}} \text{Argmin}(Z) \text{ in } (\mathcal{F}, \tau_F).$$

Proof. First, notice that by Lemma 4.6 of Ferger [7] the set $\mathbf{F}_{0,1}$ is closed in (\mathcal{F}, τ_F) and consequently $\{Z \in \mathbf{F}_{0,1}\} \in \mathcal{A}$, the domain of \mathbb{P} . It follows from Theorem 4.4 that $A(Z_\alpha, \epsilon_\alpha) \xrightarrow{\mathcal{D}} \text{Argmin}(Z)$ in (\mathcal{F}, τ_{uF}) . Thus, an application of Theorem 2.9 of Ferger [7] yields the assertion. \square

For Argmin-sets the assumption (30) simplifies significantly, because here $\epsilon_\alpha = 0$ for all $\alpha \in J$, which is a special case of the special case in Remark 4.5 (2).

Corollary 4.7. Suppose that $Z_\alpha \xrightarrow{\mathcal{D}} Z$ in (S, τ_e) , where Z has at most one minimizing point almost surely. If for every $\eta > 0$ there exists a compact $K \subseteq E$ such that

$$\liminf_{\alpha} \mathbb{P}_\alpha(\{\omega \in \Omega_\alpha : \emptyset \neq \text{Argmin}(Z_\alpha(\omega)) \subseteq K\}) \geq 1 - \eta,$$

then

$$\text{Argmin}(Z_\alpha) \xrightarrow{\mathcal{D}} \text{Argmin}(Z) \text{ in } (\mathcal{F}, \tau_F).$$

In applications one is often more interested in single ϵ -optimal solutions. So, for each $\alpha \in J$ let $\xi_\alpha : (\Omega_\alpha, \mathcal{A}_\alpha, \mathbb{P}_\alpha) \rightarrow (E, \mathcal{B}(E))$ be a measurable map. Such a random variable in E is called *measurable selection* of $A(Z_\alpha, \epsilon_\alpha)$ if $\xi_\alpha \in A(Z_\alpha, \epsilon_\alpha)$ \mathbb{P}_α -almost-surely. Since $A(Z_\alpha, \epsilon_\alpha)$ is a random closed set by Corollary 4.3, it follows from the Fundamental Selection Theorem, confer Molchanov [13] on p. 77, that measurable selections exist.

Theorem 4.8. Assume that (30) holds and that for every $\eta > 0$ there exists a compact $K \subseteq E$ with

$$\liminf_{\alpha} \mathbb{P}_\alpha(\{\omega \in \Omega_\alpha : \xi_\alpha(\omega) \in K\}) \geq 1 - \eta.$$

Then

$$\limsup_{\alpha} \mathbb{P}_\alpha(\{\omega \in \Omega_\alpha : \xi_\alpha(\omega) \in F\}) \leq T(F) \quad \text{for all closed sets } F \text{ in } E, \quad (34)$$

where $T : \mathcal{B}(E) \rightarrow [0, 1]$ is the *capacity functional* of $A(Z, \epsilon)$, i. e.

$$T(B) = \mathbb{P}(A(Z, \epsilon) \cap B \neq \emptyset) \quad \text{for all Borel-sets } B \in \mathcal{B}(E).$$

If $\epsilon = 0$ and $\text{Argmin}(Z) \subseteq \{\xi\}$ \mathbb{P} -almost surely for some random variable ξ on $(\Omega, \mathcal{A}, \mathbb{P})$ with values in E , then

$$\xi_\alpha \xrightarrow{\mathcal{D}} \xi \quad \text{in } (E, \mathcal{G}).$$

Proof. $A(Z_\alpha, \epsilon_\alpha) \xrightarrow{\mathcal{D}} A(Z, \epsilon)$ in (\mathcal{F}, τ_{uF}) by Theorem 4.4. Now, the assertion follows from Corollary 3.7 of Ferger [7]. \square

Remark 4.9. Note that (34) looks exactly like the corresponding characterisation in the Portmanteau theorem, except for the fact that T is generally not a probability measure, but only a Choquet-capacity. On the other hand, T uniquely determines the distribution of the random closed set $A(Z, \epsilon)$. It is therefore reasonable to say that the points ξ_α converge in distribution to the set $A(Z, \epsilon)$.

Remark 4.10. If we choose $F := E \in \mathcal{F}$ in (34), then we obtain that $T(E) = 1$. But $T(E) = \mathbb{P}(A(Z, \epsilon) \neq \emptyset)$, whence in particular $\text{Argmin}(Z) = A(Z, 0)$ is actually equal to $\{\xi\}$ with probability one.

5. APPENDIX

Recall that d is Vaughan's metric on E . Let u be the usual euclidian distance on \mathbb{R} . We endow the product $E \times \mathbb{R}$ with the product-metric $d \times u$ defined by $d \times u((x, \alpha), (y, \beta)) := \max\{d(x, y), |\alpha - \beta|\}$ for points (x, α) and (y, β) in $E \times \mathbb{R}$. It is well-known that this metric (among many others) induces the product-topology on $E \times \mathbb{R}$. The reason for our special choice lies in that

$$B_{d \times u}((x, \alpha), r) = B_d(x, r) \times (\alpha - r, \alpha + r), \quad (35)$$

so open balls are open rectangles. Similarly,

$$\overline{B}_{d \times u}((x, \alpha), r) = \overline{B}_d(x, r) \times [\alpha - r, \alpha + r]. \quad (36)$$

In particular, $\overline{B}_{d \times u}((x, \alpha), r)$ is compact.

Lemma 5.1. Let P be a probability measure on (S, \mathcal{B}_e) and ϕ be the homeomorphism (3). If $Q := P \circ \phi^{-1}$, then for each $(x, \alpha) \in E \times \mathbb{R}$ and $r > 0$ the pertaining closed ball $\overline{B}_{d \times u}((x, \alpha), r)$ is a Q -continuity set, if and only if

$$P(I_{\overline{B}_d(x, r)} \leq r + \alpha) = P(I_{(\overline{B}_d(x, r))^0} < r + \alpha).$$

Proof. We know that every closed ball $\overline{B}_{d \times u}((x, \alpha), r)$ is compact, whence by (5) it is a Q -continuity set if and only if

$$\begin{aligned} & P(\{f \in S : \text{epi}(f) \in \mathcal{H}(\overline{B}_d(x, r) \times [\alpha - r, \alpha + r])\}) \\ &= P(\{f \in S : \text{epi}(f) \in \mathcal{H}((\overline{B}_d(x, r) \times [\alpha - r, \alpha + r])^0)\}). \end{aligned} \quad (37)$$

Now,

$$\text{epi}(f) \cap (\overline{B}_d(x, r) \times [\alpha - r, \alpha + r]) \neq \emptyset \Leftrightarrow I_{\overline{B}_d(x, r)}(f) \leq r + \alpha. \quad (38)$$

To see this, let $(y, t) \in E \times \mathbb{R}$ such that $t \geq f(y)$, $y \in \overline{B}_d(x, r)$ and $t \in [\alpha - r, \alpha + r]$. Then $I_{\overline{B}_d(x, r)}(f) \leq f(y) \leq t \leq r + \alpha$. Conversely, assume that $I_{\overline{B}_d(x, r)}(f) \leq r + \alpha$. Since $\overline{B}_d(x, r)$ is compact and f is lsc, there exists some $z \in \overline{B}_d(x, r)$ with $f(z) = I_{\overline{B}_d(x, r)}(f)$. Consequently, $(z, r + \alpha) \in \text{epi}(f)$. Moreover, $(z, r + \alpha) \in \overline{B}_d(x, r) \times [\alpha - r, \alpha + r]$ as can be seen immediately. Thus, we have shown (38). Further,

$$\text{epi}(f) \cap (\overline{B}_d(x, r) \times [\alpha - r, \alpha + r])^0 \neq \emptyset \Leftrightarrow I_{(\overline{B}_d(x, r))^0}(f) < r + \alpha. \quad (39)$$

For the proof of necessity \Rightarrow , first observe that $(\overline{B}_d(x, r) \times [\alpha - r, \alpha + r])^0 = (\overline{B}_d(x, r))^0 \times (\alpha - r, \alpha + r)$. Let $(y, t) \in E \times \mathbb{R}$ with $t \geq f(y)$, $y \in (\overline{B}_d(x, r))^0$ and $t \in (\alpha - r, \alpha + r)$. Then $I_{(\overline{B}_d(x, r))^0}(f) \leq f(y) \leq t < r + \alpha$.

As to sufficiency \Leftarrow , put $i := I_{(\overline{B}_d(x, r))^0}(f)$. We know that $i < \alpha + r$. Then there exists some $u \in (\alpha - r, \alpha + r)$ with $i < u$, because otherwise $i \geq \alpha + r - \epsilon$ for all $\epsilon \in (0, 2r)$ and taking the limit $\epsilon \downarrow 0$ yields that $i \geq \alpha + r$, a contradiction. Therefore, there exists some $v \in (\overline{B}_d(x, r))^0$ such that $f(v) < u$, whence

$$\begin{aligned} (v, u) &\in \text{epi}(f) \cap (\overline{B}_d(x, r))^0 \times (\alpha - r, \alpha + r) \\ &= \text{epi}(f) \cap (\overline{B}_d(x, r) \times [\alpha - r, \alpha + r])^0 \neq \emptyset \end{aligned}$$

as desired.

From (37) in combination with (38) and (39) the assertion follows. \square

(Received July 22, 2025)

REFERENCES

- [1] H. Attouch: Variational Convergence for Functions and Operators. Applicable Mathematics Series, Pitmann, London 1984.
- [2] G. Beer: Topologies on Closed and Closed Convex Sets. Kluwer Academic Publishers, Dordrecht 1993.
- [3] P. Billingsley: Convergence of Probability Measures. Second Edition. John Wiley and Sons, New York 1999.
- [4] J. Fell: A Hausdorff topology for the closed subsets of a locally compact non-Hausdorff space. *Proc. Amer. Math. Soc.* 13 (1962), 472–476. DOI:10.1090/S0002-9939-1962-0139135-6
- [5] D. Ferger: Weak convergence on topological hyperspaces and random closed sets. Lecture notes, Technische Universität Dresden 2024.
- [6] D. Ferger: A continuous Mapping Theorem for the Argmin-set functional with applications to convex stochastic processes. *Kybernetika* 57 (2021), 426–445. DOI:10.14736/kyb-2021-3-0426
- [7] D. Ferger: Weak convergence of probability measures on hyperspaces with the upper Fell topology. *Bull. Iranian. Math. Soc.* 50 (2024), 79. DOI:10.1007/s41980-024-00927-3
- [8] D. Ferger: Weak convergence of probability measures to Choquet capacity functionals. *Turkish J. Math.* 42 (2018), 1747–1764. DOI:10.3906/mat-1705-106
- [9] D. Ferger: Arginf-sets of multivariate càdlàg processes and their convergence in hyperspace topologies. *Theory Stoch. Process.* 20 (2015), 36, 13–41.
- [10] P. Gänssler and W. Stute: Wahrscheinlichkeitstheorie. Springer–Verlag, Berlin, Heidelberg 1977.
- [11] O. Gersch: Convergence in Distribution of Random Closed Sets and Applications in Stability Theory of Stochastic Optimisation. PhD thesis, Technische Universität Ilmenau 2007.
- [12] O. Kallenberg: Foundations of Modern Probability. Third Edition, Volume 2. Springer Nature Switzerland AG 2021.
- [13] I. Molchanov: Theory of random sets. Second Edition. Springer–Verlag London 2017.
- [14] T. Norberg: Convergence and existence of random set distributions. *Th. Probab. Appl.* 12 (1984), 726–732.
- [15] G. Ch. Pflug: Asymptotic dominance and confidence for solutions of stochastic programs. *Czechoslovak J. Oper. Res.* 1 (1992), 21–30.
- [16] R. T. Rockafellar and R. J.-B. Wets: Variational Analysis. Springer–Verlag, Berlin, Heidelberg 1998.
- [17] G. Salinetti and R. J.-B. Wets: On the convergence in distribution of measurable multifunctions (random sets), normal integrands, stochastic processes and stochastic infima. *Math. Oper. Res.* 11 (1986), 385–419.

- [18] T. B. Singh: Introduction to Topology. Springer Nature Singapore 2019.
- [19] R. Schneider and W. Weil: Stochastic and Integral Geometry. Springer–Verlag, Berlin, Heidelberg 2008.
- [20] F. Topsøe: Topology and Measure. Lecture Notes in Mathematics 133, Springer–Verlag, Berlin–Heidelberg–New York 1970.
- [21] H. E. Vaughan: On Locally Compact Metrisable Spaces. Bull. Amer. Math. Soc. 43 (1937), 532–535. DOI:10.1090/S0002-9904-1937-06593-1
- [22] S. Vogel: Semiconvergence in distribution of random closed sets with application to random optimization problems. Ann. Oper. Res. 142 (2006), 269–282. DOI:10.1007/s10479-006-6172-0

Dietmar Ferger, Technische Universität Dresden, Fakultät Mathematik, Zellescher Weg 12-14, D-01069 Dresden. Germany.

e-mail: dietmar.ferger@tu-dresden.de